



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:17 AM GMT

PDB ID : 3A5Z
Title : Crystal structure of Escherichia coli GenX in complex with elongation factor P
Authors : Sumida, T.; Yanagisawa, T.; Ishii, R.; Yokoyama, S.
Deposited on : 2009-08-17
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	
1	C	328	
1	E	328	
1	G	328	
2	B	191	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	191	
2	F	191	
2	H	191	

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 14965 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative lysyl-tRNA synthetase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	323	2584	1631	450	487	16	0	0	0
1	C	324	2590	1634	451	489	16	0	0	0
1	E	323	2584	1631	450	487	16	0	0	0
1	G	323	2582	1630	448	487	17	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	EXPRESSION TAG	UNP C3SGA2
A	-1	SER	-	EXPRESSION TAG	UNP C3SGA2
A	0	HIS	-	EXPRESSION TAG	UNP C3SGA2
C	-2	GLY	-	EXPRESSION TAG	UNP C3SGA2
C	-1	SER	-	EXPRESSION TAG	UNP C3SGA2
C	0	HIS	-	EXPRESSION TAG	UNP C3SGA2
E	-2	GLY	-	EXPRESSION TAG	UNP C3SGA2
E	-1	SER	-	EXPRESSION TAG	UNP C3SGA2
E	0	HIS	-	EXPRESSION TAG	UNP C3SGA2
G	-2	GLY	-	EXPRESSION TAG	UNP C3SGA2
G	-1	SER	-	EXPRESSION TAG	UNP C3SGA2
G	0	HIS	-	EXPRESSION TAG	UNP C3SGA2

- Molecule 2 is a protein called Elongation factor P.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	134	1064	680	174	206	4	0	0	0
2	D	175	1368	873	225	266	4	0	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	F	59	Total 468	298	82	87	1	0	0	0
2	H	175	Total 1368	873	225	266	4	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	EXPRESSION TAG	UNP C3SGD7
B	-1	SER	-	EXPRESSION TAG	UNP C3SGD7
B	0	HIS	-	EXPRESSION TAG	UNP C3SGD7
D	-2	GLY	-	EXPRESSION TAG	UNP C3SGD7
D	-1	SER	-	EXPRESSION TAG	UNP C3SGD7
D	0	HIS	-	EXPRESSION TAG	UNP C3SGD7
F	-2	GLY	-	EXPRESSION TAG	UNP C3SGD7
F	-1	SER	-	EXPRESSION TAG	UNP C3SGD7
F	0	HIS	-	EXPRESSION TAG	UNP C3SGD7
H	-2	GLY	-	EXPRESSION TAG	UNP C3SGD7
H	-1	SER	-	EXPRESSION TAG	UNP C3SGD7
H	0	HIS	-	EXPRESSION TAG	UNP C3SGD7


- Molecule 3 is 5'-O-[(L-LYSYLAMINO)SULFONYL]ADENOSINE (three-letter code: KAA) (formula: C₁₆H₂₆N₈O₇S).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			32	16	8	7	1		
3	C	1	Total	C	N	O	S	0	0
			32	16	8	7	1		
3	E	1	Total	C	N	O	S	0	0
			32	16	8	7	1		
3	G	1	Total	C	N	O	S	0	0
			32	16	8	7	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	52	Total	O	0	0
			52	52		
4	B	4	Total	O	0	0
			4	4		
4	C	56	Total	O	0	0
			56	56		
4	D	13	Total	O	0	0
			13	13		
4	E	62	Total	O	0	0
			62	62		
4	F	2	Total	O	0	0
			2	2		
4	G	29	Total	O	0	0
			29	29		
4	H	11	Total	O	0	0
			11	11		



- Molecule 2: Elongation factor P



- Molecule 2: Elongation factor P



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	105.93Å 102.96Å 119.94Å 90.00° 99.40° 90.00°	Depositor
Resolution (Å)	45.68 – 2.50 45.68 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.7 (45.68-2.50) 93.8 (45.68-2.50)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.84 (at 2.51Å)	Xtrriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.226 , 0.268 0.225 , 0.267	Depositor DCC
R_{free} test set	4162 reflections (5.04%)	DCC
Wilson B-factor (Å ²)	54.2	Xtrriage
Anisotropy	0.331	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 41.5	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtrriage
Outliers	1 of 82571 reflections (0.001%)	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	14965	wwPDB-VP
Average B, all atoms (Å ²)	63.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 19.10% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: KAA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.44	0/2641	0.65	0/3578
1	C	0.42	0/2647	0.64	0/3586
1	E	0.42	0/2641	0.66	1/3578 (0.0%)
1	G	0.37	0/2637	0.59	0/3570
2	B	0.35	0/1088	0.61	0/1476
2	D	0.37	0/1396	0.59	0/1895
2	F	0.33	0/476	0.63	0/637
2	H	0.35	0/1396	0.60	0/1895
All	All	0.40	0/14922	0.63	1/20215 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	E	250	HIS	N-CA-C	-5.08	97.28	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	TYR	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2584	0	2522	103	0
1	C	2590	0	2527	96	0
1	E	2584	0	2522	92	0
1	G	2582	0	2526	126	0
2	B	1064	0	1028	42	0
2	D	1368	0	1344	58	0
2	F	468	0	475	15	0
2	H	1368	0	1344	50	0
3	A	32	0	26	2	0
3	C	32	0	26	2	0
3	E	32	0	26	1	0
3	G	32	0	26	7	0
4	A	52	0	0	2	0
4	B	4	0	0	1	0
4	C	56	0	0	0	0
4	D	13	0	0	0	0
4	E	62	0	0	4	0
4	F	2	0	0	0	0
4	G	29	0	0	0	0
4	H	11	0	0	1	0
All	All	14965	0	14392	566	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

The worst 5 of 566 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:116:THR:HB	2:B:124:SER:HB3	1.48	0.95
1:C:206:LYS:HD2	1:C:206:LYS:H	1.28	0.94
2:F:44:ARG:HE	2:F:51:ARG:HH12	1.03	0.94
1:E:243:ILE:HD13	1:E:307:LEU:HD21	1.50	0.93
1:G:33:ARG:HG3	1:G:33:ARG:HH21	1.35	0.91

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	321/328 (98%)	303 (94%)	18 (6%)	0	100	100
1	C	322/328 (98%)	299 (93%)	21 (6%)	2 (1%)	30	50
1	E	321/328 (98%)	304 (95%)	16 (5%)	1 (0%)	46	68
1	G	319/328 (97%)	283 (89%)	31 (10%)	5 (2%)	12	21
2	B	132/191 (69%)	113 (86%)	14 (11%)	5 (4%)	4	5
2	D	171/191 (90%)	154 (90%)	17 (10%)	0	100	100
2	F	57/191 (30%)	49 (86%)	5 (9%)	3 (5%)	2	2
2	H	171/191 (90%)	151 (88%)	18 (10%)	2 (1%)	16	29
All	All	1814/2076 (87%)	1656 (91%)	140 (8%)	18 (1%)	19	34

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	66	SER
1	G	166	LYS
2	B	6	SER
2	F	9	PHE
1	G	228	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	277/281 (99%)	256 (92%)	21 (8%)	16	30

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	278/281 (99%)	260 (94%)	18 (6%)	21	39
1	E	277/281 (99%)	257 (93%)	20 (7%)	18	33
1	G	277/281 (99%)	256 (92%)	21 (8%)	16	30
2	B	113/157 (72%)	105 (93%)	8 (7%)	18	34
2	D	148/157 (94%)	141 (95%)	7 (5%)	32	56
2	F	49/157 (31%)	47 (96%)	2 (4%)	37	63
2	H	148/157 (94%)	142 (96%)	6 (4%)	37	63
All	All	1567/1752 (89%)	1464 (93%)	103 (7%)	21	38

5 of 103 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	253	THR
1	E	53	LEU
1	G	285	GLU
2	D	41	VAL
2	D	156	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 63 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	77	ASN
1	E	94	GLN
1	G	276	GLN
2	D	87	ASN
1	E	8	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	KAA	A	990	-	27,34,34	2.67	4 (14%)	32,49,49	3.04	5 (15%)
3	KAA	C	991	-	27,34,34	2.69	4 (14%)	32,49,49	3.10	7 (21%)
3	KAA	E	992	-	27,34,34	2.70	4 (14%)	32,49,49	3.09	5 (15%)
3	KAA	G	993	-	27,34,34	2.71	4 (14%)	32,49,49	3.15	8 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	KAA	A	990	-	-	0/18/40/40	0/3/3/3
3	KAA	C	991	-	-	0/18/40/40	0/3/3/3
3	KAA	E	992	-	-	0/18/40/40	0/3/3/3
3	KAA	G	993	-	-	0/18/40/40	0/3/3/3

The worst 5 of 16 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	990	KAA	C-N8	-2.76	1.33	1.37
3	G	993	KAA	C-N8	-2.75	1.33	1.37
3	E	992	KAA	C-N8	-2.75	1.33	1.37
3	C	991	KAA	C-N8	-2.62	1.33	1.37
3	A	990	KAA	O4'-C1'	2.51	1.44	1.41

The worst 5 of 25 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	993	KAA	C-N8-S1	-9.66	110.53	124.05
3	C	991	KAA	O2S-S1-O1S	-9.62	109.24	120.77
3	G	993	KAA	N3-C2-N1	-9.62	121.53	128.89
3	C	991	KAA	N3-C2-N1	-9.60	121.55	128.89
3	E	992	KAA	O2S-S1-O1S	-9.56	109.31	120.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	990	KAA	2	0
3	C	991	KAA	2	0
3	E	992	KAA	1	0
3	G	993	KAA	7	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/328 (98%)	-0.05	0 100 100	33, 51, 73, 103	0
1	C	324/328 (98%)	0.31	17 (5%) 31 35	34, 54, 81, 97	0
1	E	323/328 (98%)	0.08	3 (0%) 85 88	31, 48, 75, 108	0
1	G	323/328 (98%)	0.64	30 (9%) 11 11	34, 73, 102, 110	0
2	B	134/191 (70%)	1.25	33 (24%) 1 1	39, 86, 123, 134	0
2	D	175/191 (91%)	0.22	10 (5%) 27 31	47, 65, 91, 107	0
2	F	59/191 (30%)	2.03	27 (45%) 0 0	48, 97, 130, 139	0
2	H	175/191 (91%)	0.42	16 (9%) 11 12	49, 73, 100, 114	0
All	All	1836/2076 (88%)	0.39	136 (7%) 17 19	31, 60, 101, 139	0

The worst 5 of 136 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	136	VAL	8.9
2	F	5	TYR	8.3
2	B	133	LEU	7.1
2	F	10	ARG	6.7
2	F	3	THR	6.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	KAA	E	992	32/32	0.98	0.18	0.47	36,39,43,45	0
3	KAA	A	990	32/32	0.98	0.16	0.23	30,39,44,44	0
3	KAA	G	993	32/32	0.97	0.19	0.22	47,56,61,62	0
3	KAA	C	991	32/32	0.97	0.19	-0.26	41,45,47,49	0

6.5 Other polymers [i](#)

There are no such residues in this entry.