wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 01:47 PM BST

PDB ID	:	1A6B
Title	:	NMR STRUCTURE OF THE COMPLEX BETWEEN THE ZINC FINGER PROTEIN NCP10 OF MOLONEY MURINE LEUKEMIA VIRUS AND A SEQUENCE OF THE PSI-PACKAGING DOMAIN OF HIV-1, 20 STRUCTURES
Authors	:	Schueler, W.; Dong, C.-Z.; Wecker, K.; Roques, B.P.
Deposited on	:	1998-02-23

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	5	20% 20% 60%
2	B	40	43% 8% 50%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 16 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	B:23-B:42 (20)	0.34	16

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 4, 5, 7, 8, 9, 10, 14, 16, 18, 20
2	3, 6, 13, 15, 17
3	2, 12
Single-model clusters	11; 19

3 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 807 atoms, of which 381 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called DNA (5'-D(*AP*CP*GP*CP*C)-3').

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	P	
1	A	5	154	47	57	19	27	4	0

- Molecule 2 is a protein called ZINC FINGER PROTEIN NCP10.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	B	40	652	196	324	72	57	3	0

- Molecule 3 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	
			Total	Zn
3	B	1	1	1

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: DNA (5'-D(*AP*CP*GP*CP*C)-3')

Chain A: 20% 20% 60%

A1 C2 G3 C4 C5

- Molecule 2: ZINC FINGER PROTEIN NCP10

Chain B: 43% 8% 50%

G14 E15 R16 R17 R18 S19 Q20 L21 D22 A27 W35 A36 P43 R44 G45 P46 R47 G48 P49 R50 P51 Q52 T53

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 16. Colouring as in section 4.1 above.

- Molecule 1: DNA (5'-D(*AP*CP*GP*CP*C)-3')

Chain A: 20% 60% 20%

A1 C2 G3 C4 C5

- Molecule 2: ZINC FINGER PROTEIN NCP10

Chain B: 35% 15% 50%

G14 E15 R16 R17 R18 S19 Q20 L21 D22 A27 K30 E31 K32 W35 A36 P43 R44 G45 P46 R47 G48 P49 R50 P51 Q52 T53

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *DYNAMICAL SIMULATED ANNEALING, ENERGY MINIMIZATION.*

Of the 50 calculated structures, 20 were deposited, based on the following criterion: *LEAST RESTRAINT VIOLATIONS, LOWEST TOTAL ENERGY.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
DISCOVER	refinement	
BRUKER UXNMR	structure solution	UXNMR
BIOSYM/MSI FELIX	structure solution	FELIX

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.32±0.02	0±0/108 (0.0±0.0%)	2.15±0.13	6±1/164 (3.8±0.7%)
2	B	0.56±0.01	0±0/170 (0.0±0.0%)	1.10±0.06	0±0/227 (0.0±0.1%)
All	All	0.93	0/5560 (0.0%)	1.62	128/7820 (1.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	1.3±0.6
2	B	0.0±0.0	0.1±0.3
All	All	0	28

There are no bond-length outliers.

5 of 15 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	4	DC	O4'-C1'-N1	12.72	116.91	108.00	3	20
1	A	2	DC	O4'-C1'-N1	11.46	116.02	108.00	4	20
1	A	1	DA	O4'-C1'-N9	10.75	115.53	108.00	5	6
1	A	3	DG	O4'-C1'-N9	9.38	114.57	108.00	11	9
1	A	2	DC	O4'-C4'-C3'	-8.58	100.85	106.00	5	2

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	3	DG	Sidechain	18
1	A	4	DC	Sidechain	7
2	B	28	TYR	Sidechain	1
2	B	23	ARG	Sidechain	1
1	A	5	DC	Sidechain	1

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	97	57	57	1±1
2	B	165	158	158	2±1
All	All	5260	4300	4300	34

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:27:ALA:HB3	2:B:36:ALA:HB2	0.52	1.80	11	12
1:A:1:DA:H2'	2:B:27:ALA:HB1	0.50	1.83	8	13
1:A:4:DC:C6	2:B:35:TRP:CH2	0.48	3.02	11	3
1:A:2:DC:C5	1:A:3:DG:C2	0.45	3.04	18	3
1:A:3:DG:C4	2:B:35:TRP:CH2	0.41	3.08	17	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
2	B	20/40 (50%)	16±2 (82±8%)	3±2 (14±8%)	1±1 (4±3%)	7 35

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	400/800 (50%)	329 (82%)	56 (14%)	15 (4%)	7 35

All 5 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	32	LYS	5
2	B	28	TYR	3
2	B	23	ARG	3
2	B	33	GLY	3
2	B	24	ASP	1

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
2	B	17/34 (50%)	15±1 (87±5%)	2±1 (13±5%)	9 50
All	All	340/680 (50%)	295 (87%)	45 (13%)	9 50

5 of 10 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	35	TRP	20
2	B	30	LYS	9
2	B	31	GLU	5
2	B	29	CYS	3
2	B	37	LYS	2

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided