wwPDB X-ray Structure Validation Summary Report (i)

Feb 1, 2016 – 12:24 AM GMT

PDB ID	:	2AAG
Title	:	Crystal Structures of the Wild-type, Mutant-P1A and Inactivated Malonate Semialdehyde Decarboxylase: A Structural Basis for the Decarboxylase and Hydratase Activities
Authors	:	Almrud, J.J.; Poelarends, G.J.; Johnson Jr., W.H.; Serrano, H.; Hackert, M.L.; Whitman, C.P.
Deposited on	:	2005-07-13
Resolution	:	1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1745 (1.86-1.86)
Clashscore	102246	1898 (1.86-1.86)
Ramachandran outliers	100387	1875 (1.86-1.86)
Sidechain outliers	100360	1875 (1.86-1.86)
RSRZ outliers	91569	1747 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	76% 17% • ••
1	B	130	% 77% 18% • •
1	C	130	% 76% 19% • ••
1	D	130	2% 78% 16% 5% •
1	E	130	2% 78% 15% 5% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	130	2% 79% 15% 5%

2 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 6770 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Malonate Semialdehyde Decarboxylase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	129	992	625	171	192	4	0	0	0
1	B	130	996	627	172	193	4	0	0	0
1	C	129	992	625	171	192	4	0	0	0
1	D	129	989	622	171	192	4	0	0	0
1	E	129	1002	631	175	192	4	0	1	0
1	F	129	992	625	171	192	4	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	LEU	ILE	ENGINEERED	UNP Q9EV83
A	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83
B	7	LEU	ILE	ENGINEERED	UNP Q9EV83
B	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83
C	7	LEU	ILE	ENGINEERED	UNP Q9EV83
C	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83
D	7	LEU	ILE	ENGINEERED	UNP Q9EV83
D	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83
E	7	LEU	ILE	ENGINEERED	UNP Q9EV83
E	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83
F	7	LEU	ILE	ENGINEERED	UNP Q9EV83
F	130	GLY	-	CLONING ARTIFACT	UNP Q9EV83

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
2	A	163	163	163	0	0
			Total	O		
2	B	147	147	147	0	0
			Total	O		
2	C	143	143	143	0	0
			Total	O		
2	D	115	115	115	0	0
			Total	O		
2	E	114	114	114	0	0
			Total	O		
2	F	125	125	125	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain A:



- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain B:



- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain C:



- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain D:



- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain E:



- Molecule 1: Malonate Semialdehyde Decarboxylase

Chain F:




P1
L2
I3
K4
L7
D13
L19
Y28
D29
D37
R38
Y39
S43
R46
D54
Y59
G60
R61
V65
Y66
I67
E78
K85
I86
R93
D100
D101
S110
D111
L124
D127
I128
V129
GLY

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α, β, γ	57.71Å 82.11Å 77.60Å 90.00° 101.15° 90.00°	Depositor
Resolution (Å)	19.21 – 1.85 19.21 – 1.82	Depositor EDS
% Data completeness (in resolution range)	96.8 (19.21-1.85) 96.4 (19.21-1.82)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$< I/\sigma(I) >$ [1]	1.16 (at 1.82Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.178 , 0.232 0.184 , 0.184	Depositor DCC
R_{free} test set	2928 reflections (5.25%)	DCC
Wilson B-factor (Å^2)	21.6	Xtriage
Anisotropy	0.016	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.34 , 46.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Outliers	0 of 60968 reflections	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	6770	wwPDB-VP
Average B, all atoms (Å^2)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.56% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#$\|Z\|$ >5	RMSZ	#$\|Z\|$ >5
1	A	1.23	4/1010 (0.4%)	1.57	17/1367 (1.2%)
1	B	1.45	6/1014 (0.6%)	1.66	22/1372 (1.6%)
1	C	1.26	6/1010 (0.6%)	1.40	22/1367 (1.6%)
1	D	1.29	5/1007 (0.5%)	1.44	19/1362 (1.4%)
1	E	1.29	7/1020 (0.7%)	1.56	21/1381 (1.5%)
1	F	1.32	8/1010 (0.8%)	1.63	24/1367 (1.8%)
All	All	1.31	36/6071 (0.6%)	1.55	125/8216 (1.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
1	C	0	1
1	D	0	1
1	E	0	1
1	F	0	1
All	All	0	6

The worst 5 of 36 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	39	TYR	CE1-CZ	-14.10	1.20	1.38
1	B	39	TYR	CE2-CZ	-13.59	1.20	1.38
1	B	39	TYR	CG-CD2	-11.97	1.23	1.39
1	B	39	TYR	CG-CD1	-10.30	1.25	1.39
1	F	65	VAL	CB-CG2	-8.06	1.35	1.52

The worst 5 of 125 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	B	61	ARG	NE-CZ-NH1	26.06	133.33	120.30
1	A	61	ARG	NE-CZ-NH1	25.35	132.98	120.30
1	F	61	ARG	NE-CZ-NH1	23.30	131.95	120.30
1	E	61	ARG	NE-CZ-NH1	21.55	131.07	120.30
1	D	61	ARG	NE-CZ-NH1	18.09	129.34	120.30

There are no chirality outliers.

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	59	TYR	Peptide
1	B	59	TYR	Peptide
1	C	59	TYR	Peptide
1	D	59	TYR	Peptide
1	E	59	TYR	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	992	0	977	25	0
1	B	996	0	980	24	0
1	C	992	0	977	21	0
1	D	989	0	969	17	0
1	E	1002	0	987	20	0
1	F	992	0	977	13	0
2	A	163	0	0	12	2
2	B	147	0	0	4	0
2	C	143	0	0	4	3
2	D	115	0	0	4	0
2	E	114	0	0	3	0
2	F	125	0	0	3	1
All	All	6770	0	5867	104	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:46:ARG:NH2	1:E:46[A]:ARG:NH2	1.89	1.18
2:A:288:HOH:O	1:C:48:GLY:HA3	0.96	1.11
1:A:108:GLU:HG2	2:A:266:HOH:O	1.61	0.98
1:B:7:LEU:H	1:B:7:LEU:HD23	1.30	0.93
1:F:7:LEU:H	1:F:7:LEU:HD23	1.32	0.92

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:165:HOH:O	2:C:138:HOH:O[2_555]	1.79	0.41
2:A:141:HOH:O	2:C:265:HOH:O[2_655]	2.01	0.19
2:C:181:HOH:O	2:F:143:HOH:O[1_455]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	127/130 (98%)	124 (98%)	3 (2%)	0	100	100
1	B	128/130 (98%)	124 (97%)	4 (3%)	0	100	100
1	C	127/130 (98%)	122 (96%)	5 (4%)	0	100	100
1	D	127/130 (98%)	123 (97%)	4 (3%)	0	100	100
1	E	128/130 (98%)	123 (96%)	5 (4%)	0	100	100
1	F	127/130 (98%)	123 (97%)	4 (3%)	0	100	100
All	All	764/780 (98%)	739 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	106/106 (100%)	98 (92%)	8 (8%)	17	4
1	B	106/106 (100%)	100 (94%)	6 (6%)	25	9
1	C	106/106 (100%)	101 (95%)	5 (5%)	32	13
1	D	105/106 (99%)	101 (96%)	4 (4%)	40	19
1	E	106/106 (100%)	101 (95%)	5 (5%)	32	13
1	F	106/106 (100%)	100 (94%)	6 (6%)	25	9
All	All	635/636 (100%)	601 (95%)	34 (5%)	27	10

5 of 34 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	19	LEU
1	D	2	LEU
1	F	85	LYS
1	C	124	LEU
1	A	86	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	129/130 (99%)	-0.26	0	100	100	17, 21, 28, 36	0
1	B	130/130 (100%)	-0.25	1 (0%)	87	87	17, 21, 29, 39	0
1	C	129/130 (99%)	-0.22	1 (0%)	87	87	16, 21, 29, 36	0
1	D	129/130 (99%)	-0.20	3 (2%)	64	62	17, 22, 29, 35	0
1	E	129/130 (99%)	-0.14	2 (1%)	74	74	16, 21, 28, 35	0
1	F	129/130 (99%)	-0.15	3 (2%)	64	62	16, 21, 28, 37	0
All	All	775/780 (99%)	-0.20	10 (1%)	79	79	16, 21, 29, 39	0

The worst 5 of 10 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	E	78	GLU	3.8
1	C	48	GLY	3.1
1	B	130	GLY	3.0
1	F	28	VAL	2.7
1	F	129	VAL	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.