



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:19 AM GMT

PDB ID : 3AAE
Title : Crystal structure of Actin capping protein in complex with CARMIL fragment
Authors : Takeda, S.; Minakata, S.; Narita, A.; Kitazawa, M.; Yamakuni, T.; Maeda, Y.;
Nitanai, Y.
Deposited on : 2009-11-16
Resolution : 3.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2060 (3.40-3.20)
Clashscore	102246	1058 (3.38-3.22)
Ramachandran outliers	100387	1038 (3.38-3.22)
Sidechain outliers	100360	1037 (3.38-3.22)
RSRZ outliers	91569	2070 (3.40-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	<div> <div>3%</div> <div>78%</div> <div>15%</div> <div>6%</div> </div>
1	C	286	<div> <div>%</div> <div>76%</div> <div>16%</div> <div>6%</div> </div>
1	E	286	<div> <div>6%</div> <div>74%</div> <div>18%</div> <div>6%</div> </div>
1	G	286	<div> <div>4%</div> <div>75%</div> <div>17%</div> <div>6%</div> </div>
1	I	286	<div> <div>5%</div> <div>74%</div> <div>18%</div> <div>6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	277	<p>78% 16% • 5%</p>
2	D	277	<p>79% 14% • 5%</p>
2	F	277	<p>76% 13% • 10%</p>
2	H	277	<p>73% 15% • 10%</p>
2	J	277	<p>75% 14% • 10%</p>
3	V	37	<p>49% 8% 43%</p>
3	W	37	<p>51% 16% • 30%</p>
3	X	37	<p>59% 14% 5% 22%</p>
3	Y	37	<p>70% 5% 24%</p>
3	Z	37	<p>62% 11% • 24%</p>

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 22177 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called F-actin-capping protein subunit alpha-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	270	Total	C	N	O	S	0	0	0
			2196	1384	385	422	5			
1	C	270	Total	C	N	O	S	0	0	0
			2196	1384	385	422	5			
1	E	270	Total	C	N	O	S	0	0	0
			2196	1384	385	422	5			
1	G	270	Total	C	N	O	S	0	0	0
			2196	1384	385	422	5			
1	I	270	Total	C	N	O	S	0	0	0
			2196	1384	385	422	5			

- Molecule 2 is a protein called F-actin-capping protein subunit beta isoforms 1 and 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	264	Total	C	N	O	S	0	0	0
			2090	1305	363	412	10			
2	D	264	Total	C	N	O	S	0	0	0
			2090	1305	363	412	10			
2	F	250	Total	C	N	O	S	0	0	0
			1967	1227	340	390	10			
2	H	249	Total	C	N	O	S	0	0	0
			1959	1223	338	388	10			
2	J	249	Total	C	N	O	S	0	0	0
			1959	1223	338	388	10			

- Molecule 3 is a protein called 32mer peptide from Leucine-rich repeat-containing protein 16A.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	X	29	Total	C	N	O	0	0	0
			247	154	50	43			
3	Y	28	Total	C	N	O	0	0	0
			239	148	49	42			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
3	Z	28	Total	C	N	O	0	0	0
			239	148	49	42			
3	W	26	Total	C	N	O	0	0	0
			225	139	47	39			
3	V	21	Total	C	N	O	0	0	0
			182	113	39	30			


There are 25 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
X	966	GLY	-	EXPRESSION TAG	UNP Q6EDY6
X	967	PRO	-	EXPRESSION TAG	UNP Q6EDY6
X	968	LEU	-	EXPRESSION TAG	UNP Q6EDY6
X	969	GLY	-	EXPRESSION TAG	UNP Q6EDY6
X	970	SER	-	EXPRESSION TAG	UNP Q6EDY6
Y	966	GLY	-	EXPRESSION TAG	UNP Q6EDY6
Y	967	PRO	-	EXPRESSION TAG	UNP Q6EDY6
Y	968	LEU	-	EXPRESSION TAG	UNP Q6EDY6
Y	969	GLY	-	EXPRESSION TAG	UNP Q6EDY6
Y	970	SER	-	EXPRESSION TAG	UNP Q6EDY6
Z	966	GLY	-	EXPRESSION TAG	UNP Q6EDY6
Z	967	PRO	-	EXPRESSION TAG	UNP Q6EDY6
Z	968	LEU	-	EXPRESSION TAG	UNP Q6EDY6
Z	969	GLY	-	EXPRESSION TAG	UNP Q6EDY6
Z	970	SER	-	EXPRESSION TAG	UNP Q6EDY6
W	966	GLY	-	EXPRESSION TAG	UNP Q6EDY6
W	967	PRO	-	EXPRESSION TAG	UNP Q6EDY6
W	968	LEU	-	EXPRESSION TAG	UNP Q6EDY6
W	969	GLY	-	EXPRESSION TAG	UNP Q6EDY6
W	970	SER	-	EXPRESSION TAG	UNP Q6EDY6
V	966	GLY	-	EXPRESSION TAG	UNP Q6EDY6
V	967	PRO	-	EXPRESSION TAG	UNP Q6EDY6
V	968	LEU	-	EXPRESSION TAG	UNP Q6EDY6
V	969	GLY	-	EXPRESSION TAG	UNP Q6EDY6
V	970	SER	-	EXPRESSION TAG	UNP Q6EDY6


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: F-actin-capping protein subunit alpha-1




- Molecule 1: F-actin-capping protein subunit alpha-1




- Molecule 1: F-actin-capping protein subunit alpha-1




- Molecule 1: F-actin-capping protein subunit alpha-1









- Molecule 1: F-actin-capping protein subunit alpha-1





- Molecule 2: F-actin-capping protein subunit beta isoforms 1 and 2




- Molecule 2: F-actin-capping protein subunit beta isoforms 1 and 2




- Molecule 2: F-actin-capping protein subunit beta isoforms 1 and 2




- Molecule 2: F-actin-capping protein subunit beta isoforms 1 and 2




- Molecule 2: F-actin-capping protein subunit beta isoforms 1 and 2




- Molecule 3: 32mer peptide from Leucine-rich repeat-containing protein 16A




- Molecule 3: 32mer peptide from Leucine-rich repeat-containing protein 16A



- Molecule 3: 32mer peptide from Leucine-rich repeat-containing protein 16A



- Molecule 3: 32mer peptide from Leucine-rich repeat-containing protein 16A



- Molecule 3: 32mer peptide from Leucine-rich repeat-containing protein 16A

Chain V: 

GLY	PRO	LEU	GLY	SER	SER	GLY	LEU	ILE	SER	GLU	L978	E982	G983	R984	R985	L986	R998	LYS	GLN	GLN	PRO
-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	-----	------	------	------	------	------	------	------	-----	-----	-----	-----

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	231.15Å 104.36Å 186.63Å 90.00° 119.09° 90.00°	Depositor
Resolution (Å)	49.70 – 3.30 49.70 – 3.30	Depositor EDS
% Data completeness (in resolution range)	99.5 (49.70-3.30) 99.5 (49.70-3.30)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.65 (at 3.33Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.243 , 0.271 0.228 , 0.255	Depositor DCC
R_{free} test set	2955 reflections (5.33%)	DCC
Wilson B-factor (Å ²)	55.6	Xtriage
Anisotropy	0.049	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 16.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 58417 reflections	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	22177	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2247	0.90	8/3046 (0.3%)
1	C	0.47	0/2247	0.60	1/3046 (0.0%)
1	E	0.49	0/2247	0.63	1/3046 (0.0%)
1	G	0.48	0/2247	0.59	0/3046
1	I	0.49	0/2247	0.76	5/3046 (0.2%)
2	B	0.54	0/2125	0.63	4/2869 (0.1%)
2	D	0.51	1/2125 (0.0%)	0.77	5/2869 (0.2%)
2	F	0.48	0/2001	0.75	4/2704 (0.1%)
2	H	0.51	0/1993	0.62	3/2693 (0.1%)
2	J	0.52	0/1993	0.61	2/2693 (0.1%)
3	V	0.52	0/185	0.63	0/244
3	W	0.56	0/229	0.64	0/303
3	X	0.52	0/251	0.71	0/333
3	Y	0.49	0/243	0.62	0/322
3	Z	0.51	0/243	0.65	0/322
All	All	0.50	1/22623 (0.0%)	0.69	33/30582 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	8	CYS	CB-SG	-5.62	1.72	1.81

The worst 5 of 33 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	129	ARG	NE-CZ-NH1	-18.03	111.28	120.30
1	A	129	ARG	NE-CZ-NH2	17.82	129.21	120.30
1	A	91	ARG	NE-CZ-NH2	17.79	129.19	120.30
1	A	91	ARG	NE-CZ-NH1	-17.58	111.51	120.30
1	I	15	ARG	NE-CZ-NH2	-17.44	111.58	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2196	0	2117	36	0
1	C	2196	0	2117	31	0
1	E	2196	0	2117	33	0
1	G	2196	0	2117	35	0
1	I	2196	0	2117	35	0
2	B	2090	0	2076	37	0
2	D	2090	0	2076	24	0
2	F	1967	0	1943	24	0
2	H	1959	0	1937	29	0
2	J	1959	0	1937	27	0
3	V	182	0	192	1	0
3	W	225	0	234	3	0
3	X	247	0	261	14	0
3	Y	239	0	250	1	0
3	Z	239	0	250	3	0
All	All	22177	0	21741	295	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 295 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:X:975:ILE:HG23	3:X:976:SER:H	1.03	1.12
3:X:975:ILE:HG23	3:X:976:SER:N	1.64	1.06
2:H:174:MET:HG2	2:H:192:SER:HB3	1.42	1.01
2:B:174:MET:HG2	2:B:192:SER:HB3	1.44	0.99
2:D:174:MET:HG2	2:D:192:SER:HB3	1.42	0.98

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	268/286 (94%)	247 (92%)	17 (6%)	4 (2%)	13	49
1	C	268/286 (94%)	247 (92%)	16 (6%)	5 (2%)	10	45
1	E	268/286 (94%)	246 (92%)	17 (6%)	5 (2%)	10	45
1	G	268/286 (94%)	247 (92%)	16 (6%)	5 (2%)	10	45
1	I	268/286 (94%)	247 (92%)	16 (6%)	5 (2%)	10	45
2	B	262/277 (95%)	249 (95%)	11 (4%)	2 (1%)	24	62
2	D	262/277 (95%)	247 (94%)	13 (5%)	2 (1%)	24	62
2	F	248/277 (90%)	233 (94%)	14 (6%)	1 (0%)	39	76
2	H	247/277 (89%)	235 (95%)	9 (4%)	3 (1%)	16	54
2	J	247/277 (89%)	232 (94%)	14 (6%)	1 (0%)	39	76
3	V	19/37 (51%)	17 (90%)	2 (10%)	0	100	100
3	W	24/37 (65%)	19 (79%)	5 (21%)	0	100	100
3	X	27/37 (73%)	24 (89%)	2 (7%)	1 (4%)	4	27
3	Y	26/37 (70%)	23 (88%)	3 (12%)	0	100	100
3	Z	26/37 (70%)	26 (100%)	0	0	100	100
All	All	2728/3000 (91%)	2539 (93%)	155 (6%)	34 (1%)	16	54

5 of 34 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	115	SER
1	E	115	SER
1	G	84	ASN
1	I	115	SER
2	J	36	CYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	239/252 (95%)	233 (98%)	6 (2%)	55	82
1	C	239/252 (95%)	230 (96%)	9 (4%)	40	75
1	E	239/252 (95%)	230 (96%)	9 (4%)	40	75
1	G	239/252 (95%)	232 (97%)	7 (3%)	50	80
1	I	239/252 (95%)	232 (97%)	7 (3%)	50	80
2	B	235/248 (95%)	226 (96%)	9 (4%)	40	75
2	D	235/248 (95%)	225 (96%)	10 (4%)	35	72
2	F	221/248 (89%)	213 (96%)	8 (4%)	42	76
2	H	220/248 (89%)	212 (96%)	8 (4%)	42	76
2	J	220/248 (89%)	212 (96%)	8 (4%)	42	76
3	V	20/33 (61%)	19 (95%)	1 (5%)	30	68
3	W	25/33 (76%)	22 (88%)	3 (12%)	6	27
3	X	28/33 (85%)	26 (93%)	2 (7%)	18	55
3	Y	27/33 (82%)	26 (96%)	1 (4%)	41	76
3	Z	27/33 (82%)	24 (89%)	3 (11%)	8	31
All	All	2453/2665 (92%)	2362 (96%)	91 (4%)	41	76

5 of 91 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	129	ARG
2	F	227	THR
3	Z	981	GLU
1	E	191	VAL
2	F	108	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 67 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	34	ASN
2	F	4	GLN
1	I	197	HIS
1	E	58	ASN
1	E	84	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	270/286 (94%)	0.49	8 (2%) 54 47	31, 51, 85, 98	0
1	C	270/286 (94%)	0.45	2 (0%) 89 86	31, 51, 85, 98	0
1	E	270/286 (94%)	0.51	17 (6%) 23 19	31, 51, 85, 98	0
1	G	270/286 (94%)	0.45	12 (4%) 38 31	31, 51, 85, 98	0
1	I	270/286 (94%)	0.50	14 (5%) 31 25	31, 51, 85, 98	0
2	B	264/277 (95%)	0.38	3 (1%) 82 78	29, 44, 68, 81	0
2	D	264/277 (95%)	0.33	3 (1%) 82 78	29, 44, 69, 81	0
2	F	250/277 (90%)	0.11	3 (1%) 81 76	30, 43, 63, 72	0
2	H	249/277 (89%)	0.22	5 (2%) 68 62	30, 43, 62, 72	0
2	J	249/277 (89%)	0.22	7 (2%) 56 50	30, 43, 62, 72	0
3	V	21/37 (56%)	0.80	1 (4%) 34 28	56, 65, 79, 80	0
3	W	26/37 (70%)	0.87	3 (11%) 6 5	56, 70, 83, 85	0
3	X	29/37 (78%)	0.90	3 (10%) 9 7	56, 70, 79, 82	0
3	Y	28/37 (75%)	1.20	8 (28%) 1 1	56, 73, 82, 83	0
3	Z	28/37 (75%)	0.61	1 (3%) 46 39	56, 72, 81, 82	0
All	All	2758/3000 (91%)	0.39	90 (3%) 50 43	29, 47, 79, 98	0

The worst 5 of 90 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	X	1001	GLN	6.3
2	H	141	SER	5.8
2	J	141	SER	5.3
1	E	112	ASP	5.1
2	B	141	SER	5.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.