



# Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 11:06 PM GMT

PDB ID : 5AGP  
Title : Structure of rat neuronal nitric oxide synthase heme domain in complex with (S)-2-Amino-5-(2-mercaptoacetimidamido)pentanoic acid  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2015-02-02  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3939 (2.10-2.10)
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	VUR	A	800	-	-	-	X
5	ACT	A	860	-	-	-	X

## 2 Entry composition [\(i\)](#)


There are 7 unique types of molecules in this entry. The entry contains 7099 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, BRAIN.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	408	3320	2125	567	607	21	0	1	1
1	B	411	3360	2150	574	614	22	0	4	0

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	Fe	N	O		
2	A	1	43	34	1	4	4	0	0
2	B	1	43	34	1	4	4	0	0

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: C<sub>9</sub>H<sub>15</sub>N<sub>5</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 17 9 5 3	0	0
3	B	1	Total C N O 17 9 5 3	0	0

- Molecule 4 is (S)-2-AMINO-5-(2-MERCAPTOACETIMIDAMIDO)PENTANOIC ACID (three-letter code: VUR) (formula: C<sub>7</sub>H<sub>15</sub>N<sub>3</sub>O<sub>2</sub>S).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C N O S 13 7 3 2 1	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
4	B	1	13	7	3	2	1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	C	O		
5	A	1	4	2	2	0	0
5	B	1	4	2	2	0	0

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
6	B	1	1	1	0	0


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
7	A	123	123	123	0	0
7	B	141	141	141	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



- Molecule 1: NITRIC OXIDE SYNTHASE, BRAIN



## 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.87 Å   110.62 Å   164.21 Å 90.00°   90.00°   90.00°	Depositor
Resolution (Å)	91.91 – 2.10 46.96 – 2.10	Depositor EDS
% Data completeness (in resolution range)	98.2 (91.91-2.10) 98.4 (46.96-2.10)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) >$ <sup>1</sup>	2.68 (at 2.10 Å)	Xtriage
Refinement program	REFMAC 5.8.0049	Depositor
$R$ , $R_{free}$	0.196 , 0.251 0.196 , 0.250	Depositor DCC
$R_{free}$ test set	2740 reflections (5.22%)	DCC
Wilson B-factor (Å <sup>2</sup> )	26.9	Xtriage
Anisotropy	1.006	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 45.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$<  L  > = 0.48$ , $< L^2 > = 0.31$	Xtriage
Outliers	5 of 55188 reflections (0.009%)	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7099	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	39.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.80% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $< |L| >$ ,  $< L^2 >$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN, VUR, H4B, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	0.81	1/3416 (0.0%)	0.89	4/4635 (0.1%)
1	B	0.82	0/3465	0.86	5/4697 (0.1%)
All	All	0.81	1/6881 (0.0%)	0.87	9/9332 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	360	GLU	CD-OE1	7.38	1.33	1.25

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	547	ARG	NE-CZ-NH1	7.59	124.09	120.30
1	A	410	ARG	NE-CZ-NH2	-7.35	116.63	120.30
1	A	363	ASP	CB-CG-OD1	6.85	124.47	118.30
1	A	410	ARG	NE-CZ-NH1	6.78	123.69	120.30
1	B	547	ARG	NE-CZ-NH2	-5.90	117.35	120.30
1	B	471	ASP	CB-CG-OD1	5.54	123.29	118.30
1	A	709	ASP	CB-CG-OD1	5.52	123.27	118.30
1	B	414	ARG	NE-CZ-NH2	-5.33	117.63	120.30
1	B	489	ASP	CB-CG-OD1	5.21	122.99	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	299	ARG	Peptide

## 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3320	0	3227	43	0
1	B	3360	0	3280	36	0
2	A	43	0	30	0	0
2	B	43	0	30	5	0
3	A	17	0	15	0	0
3	B	17	0	15	1	0
4	A	13	0	12	1	0
4	B	13	0	12	1	0
5	A	4	0	3	0	0
5	B	4	0	3	1	0
6	B	1	0	0	0	0
7	A	123	0	0	5	0
7	B	141	0	0	0	0
All	All	7099	0	6627	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:608:GLU:HG2	7:A:2091:HOH:O	1.55	1.05
1:A:523:LEU:HD22	1:A:531:PRO:HB2	1.46	0.98
1:A:631:VAL:HG11	1:B:628:GLN:HG2	1.57	0.86
1:B:620:LYS:HE2	1:B:621:THR:H	1.43	0.81
1:A:300:PHE:HD2	1:A:315:THR:HG22	1.46	0.81
1:B:607:LEU:HD13	1:B:626:LYS:HG2	1.66	0.78
1:B:620:LYS:HE2	1:B:621:THR:N	2.05	0.70

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:620:LYS:HB2	1:A:620:LYS:NZ	2.12	0.64
1:A:316:LEU:HD12	1:A:319:LYS:HE2	1.80	0.64
1:B:420:GLN:HE21	5:B:860:ACT:H2	1.64	0.63
1:A:369:ILE:HG13	1:A:371:ARG:HG3	1.80	0.62
1:A:507:GLN:O	1:A:507:GLN:HG2	1.98	0.62
1:A:350:THR:N	1:A:353:GLN:NE2	2.47	0.61
1:A:631:VAL:CG1	1:B:628:GLN:HG2	2.28	0.61
1:B:351:LYS:HB3	1:B:351:LYS:NZ	2.17	0.59
2:B:750:HEM:HBC2	2:B:750:HEM:CMC	2.33	0.58
1:B:706:TYR:OH	2:B:750:HEM:O2D	2.17	0.57
1:A:300:PHE:CD2	1:A:315:THR:HG22	2.34	0.57
1:A:609:GLU:CB	7:A:2092:HOH:O	2.53	0.56
1:A:571:LEU:HD21	1:A:578:GLU:HB3	1.87	0.56
2:B:750:HEM:HBC2	2:B:750:HEM:HMC1	1.88	0.56
1:A:414:ARG:HD3	1:A:678:TRP:CD2	2.40	0.56
1:A:536:ILE:O	1:A:538:PRO:HD3	2.05	0.56
1:A:478:GLN:HB2	1:A:481:ARG:HG3	1.88	0.56
1:B:409:TRP:CE3	1:B:421:TRP:HA	2.41	0.55
1:A:306:TRP:CD2	1:B:336:MET:HE3	2.44	0.53
1:A:455:LEU:HD13	1:A:564:LEU:HD12	1.92	0.51
1:A:525:GLN:HG3	1:A:529:ASN:O	2.11	0.51
1:B:323:GLU:O	1:B:699:ARG:HD3	2.11	0.51
1:B:388:ILE:O	1:B:392:SER:HA	2.11	0.50
1:B:684:SER:HB3	1:B:687:ILE:HG12	1.93	0.50
1:B:595:VAL:O	1:B:599:CYS:HB2	2.12	0.50
1:A:380:ARG:HD3	7:A:2024:HOH:O	2.11	0.50
1:A:321:THR:HG23	1:A:322:LEU:HG	1.93	0.50
1:B:299:ARG:CZ	1:B:299:ARG:HB3	2.42	0.49
1:A:336:MET:HE3	1:B:306:TRP:CD2	2.47	0.49
1:A:508:GLN:NE2	1:A:716:TRP:CH2	2.81	0.49
1:A:620:LYS:HZ1	1:A:620:LYS:HB2	1.78	0.48
1:B:308:THR:O	1:B:309:ASP:HB2	2.13	0.48
1:A:686:SER:HA	1:A:691:PHE:CG	2.48	0.48
1:A:632:GLU:OE2	1:B:628:GLN:NE2	2.48	0.47
1:B:566:ALA:HA	1:B:584:PHE:O	2.14	0.47
1:A:485:TYR:CE2	1:A:512:ALA:HB1	2.49	0.47
1:B:470:HIS:O	1:B:527:ASN:HA	2.16	0.46
7:A:2114:HOH:O	1:B:337:LEU:HD12	2.15	0.46
1:A:609:GLU:HB3	7:A:2092:HOH:O	2.15	0.46
1:B:391:THR:O	1:B:392:SER:CB	2.64	0.46
1:B:322:LEU:HD13	1:B:699:ARG:HH21	1.82	0.45

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:465:ARG:HD2	1:A:578:GLU:OE2	2.16	0.45
1:B:445:HIS:C	1:B:445:HIS:CD2	2.90	0.45
1:B:610:VAL:HG21	1:B:633:ILE:HD11	1.98	0.44
1:B:462:PHE:HB2	1:B:581:ALA:HB3	1.98	0.44
1:B:474:VAL:HG11	1:B:568:SER:HB2	1.99	0.44
2:B:750:HEM:HHC	2:B:750:HEM:HBB2	1.99	0.44
1:B:620:LYS:HA	1:B:620:LYS:CE	2.47	0.44
1:A:550:LYS:HB2	1:A:550:LYS:HZ3	1.82	0.43
1:A:350:THR:N	1:A:353:GLN:HE22	2.16	0.43
1:A:415:CYS:HB3	1:A:418:ARG:HG3	2.00	0.43
1:A:475:TRP:HB2	1:A:523:LEU:HB3	2.00	0.43
1:A:336:MET:HE2	1:B:306:TRP:CD1	2.55	0.42
1:B:607:LEU:HD13	1:B:626:LYS:CG	2.42	0.42
2:B:750:HEM:CBC	2:B:750:HEM:HMC1	2.49	0.42
1:B:678:TRP:HA	3:B:760:H4B:N1	2.34	0.42
1:A:420:GLN:OE1	1:A:423:LYS:HE2	2.20	0.42
1:A:353:GLN:HB2	1:A:353:GLN:HE21	1.65	0.42
1:B:388:ILE:O	1:B:392:SER:N	2.52	0.42
1:A:355:PHE:CE1	1:A:385:ASN:HB2	2.54	0.42
1:A:525:GLN:HB2	1:A:531:PRO:HB3	2.02	0.42
1:A:475:TRP:CE3	1:A:523:LEU:HD13	2.56	0.41
1:A:626:LYS:HB3	1:B:687:ILE:HD12	2.03	0.41
4:A:800:VUR:HNE	4:A:800:VUR:HBA	1.72	0.41
1:B:299:ARG:O	1:B:318:LEU:HD11	2.21	0.41
4:B:800:VUR:CD	4:B:800:VUR:S3	3.09	0.40
1:B:701:THR:HA	1:B:702:PRO:C	2.42	0.40
1:A:460:THR:O	1:A:583:PRO:HD2	2.21	0.40
1:A:409:TRP:CE3	1:A:421:TRP:HA	2.56	0.40
1:A:694:GLU:HB3	1:B:335:ILE:HD13	2.03	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles

#### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/422 (96%)	386 (95%)	19 (5%)	0	100	100
1	B	411/422 (97%)	398 (97%)	13 (3%)	0	100	100
All	All	816/844 (97%)	784 (96%)	32 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	364/377 (97%)	347 (95%)	17 (5%)	32	30
1	B	370/377 (98%)	353 (95%)	17 (5%)	33	31
All	All	734/754 (97%)	700 (95%)	34 (5%)	32	31

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	360	GLU
1	A	364	GLN
1	A	371	ARG
1	A	381	LEU
1	A	390	SER
1	A	507	GLN
1	A	523	LEU
1	A	547	ARG
1	A	550	LYS
1	A	552	ASP
1	A	569	ASN
1	A	578	GLU
1	A	601	ASN
1	A	608	GLU
1	A	612	LYS
1	A	645	LYS
1	A	663	GLU
1	B	302	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	330	ILE
1	B	337	LEU
1	B	351	LYS
1	B	353	GLN
1	B	360	GLU
1	B	390	SER
1	B	392	SER
1	B	540	LEU
1	B	547	ARG
1	B	550	LYS
1	B	573	GLU
1	B	608	GLU
1	B	612	LYS
1	B	620	LYS
1	B	626	LYS
1	B	717	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	353	GLN
1	A	364	GLN
1	A	425	GLN
1	A	454	ASN
1	A	507	GLN
1	A	569	ASN
1	A	605	ASN
1	A	642	GLN
1	A	697	ASN
1	B	364	GLN
1	B	420	GLN
1	B	454	ASN
1	B	507	GLN
1	B	601	ASN
1	B	605	ASN
1	B	697	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 9 ligands modelled in this entry, 1 is monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	HEM	A	750	1	30,50,50	2.42	8 (26%)	24,82,82	2.97	12 (50%)
3	H4B	A	760	-	13,18,18	1.14	1 (7%)	11,26,26	3.06	6 (54%)
4	VUR	A	800	-	7,12,12	2.94	1 (14%)	5,14,14	1.89	1 (20%)
5	ACT	A	860	-	1,3,3	1.98	0	0,3,3	0.00	-
2	HEM	B	750	1,4	30,50,50	1.98	6 (20%)	24,82,82	2.87	11 (45%)
3	H4B	B	760	-	13,18,18	1.47	3 (23%)	11,26,26	2.98	3 (27%)
4	VUR	B	800	2	7,12,12	3.15	2 (28%)	5,14,14	1.61	1 (20%)
5	ACT	B	860	-	1,3,3	2.48	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	750	1	-	0/10/54/54	0/0/8/8
3	H4B	A	760	-	-	0/8/17/17	0/2/2/2
4	VUR	A	800	-	-	1/7/13/13	0/0/0/0
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
2	HEM	B	750	1,4	-	0/10/54/54	0/0/8/8

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	H4B	B	760	-	-	0/8/17/17	0/2/2/2
4	VUR	B	800	2	-	0/7/13/13	0/0/0/0
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	800	VUR	CD-NE	-7.84	1.27	1.46
4	A	800	VUR	CD-NE	-7.62	1.28	1.46
2	A	750	HEM	C2D-C3D	-6.74	1.34	1.54
2	B	750	HEM	C2D-C3D	-6.31	1.35	1.54
2	A	750	HEM	C3D-C4D	-5.65	1.44	1.51
2	B	750	HEM	C2C-C1C	-5.41	1.42	1.52
2	A	750	HEM	C2C-C1C	-5.29	1.42	1.52
2	A	750	HEM	C2B-C1B	-3.31	1.41	1.51
2	B	750	HEM	C2B-C1B	-3.07	1.41	1.51
2	A	750	HEM	C3B-C4B	-2.96	1.49	1.51
2	B	750	HEM	C3D-C4D	-2.34	1.48	1.51
2	B	750	HEM	C3C-CAC	-2.21	1.47	1.51
2	A	750	HEM	C3C-CAC	-2.12	1.47	1.51
3	B	760	H4B	C6-N5	2.06	1.50	1.45
3	B	760	H4B	C7-N8	2.32	1.49	1.46
5	B	860	ACT	CH3-C	2.48	1.52	1.48
4	B	800	VUR	C2-C1	2.50	1.54	1.49
3	A	760	H4B	C2-N2	2.59	1.39	1.34
2	B	750	HEM	FE-NC	2.63	2.06	1.95
3	B	760	H4B	C2-N1	2.75	1.40	1.35
2	A	750	HEM	CHC-C1C	2.83	1.43	1.36
2	A	750	HEM	FE-NC	3.90	2.11	1.95

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	750	HEM	CBA-CAA-C2A	-6.88	100.21	112.53
2	B	750	HEM	CBA-CAA-C2A	-6.70	100.52	112.53
3	A	760	H4B	N3-C2-N1	-5.11	117.16	125.53
2	A	750	HEM	C3C-CAC-CBC	-3.72	118.74	124.46
2	A	750	HEM	C1D-CHD-C4C	-3.67	119.68	125.82
2	B	750	HEM	C1D-CHD-C4C	-3.67	119.69	125.82
2	B	750	HEM	C3C-CAC-CBC	-3.48	119.12	124.46
2	A	750	HEM	CBD-CAD-C3D	-2.90	105.11	113.55
2	A	750	HEM	C3B-C4B-NB	-2.71	106.46	111.63

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	750	HEM	C3B-C4B-NB	-2.64	106.58	111.63
2	B	750	HEM	CAA-C2A-C1A	-2.50	124.29	127.01
2	B	750	HEM	C2D-C3D-C4D	2.13	105.10	101.50
3	B	760	H4B	C4-N3-C2	2.24	119.05	115.94
2	A	750	HEM	CMD-C2D-C3D	2.54	125.59	114.35
2	A	750	HEM	C3B-C4B-CHC	2.62	126.86	123.16
3	A	760	H4B	C4A-C8A-N8	2.84	121.78	118.43
2	A	750	HEM	CAD-C3D-C4D	3.17	123.66	112.47
3	B	760	H4B	C4A-C8A-N8	3.18	122.18	118.43
2	B	750	HEM	CMD-C2D-C3D	3.25	128.72	114.35
2	A	750	HEM	C2D-C3D-C4D	3.36	107.20	101.50
3	A	760	H4B	N2-C2-N3	3.42	122.86	117.20
4	B	800	VUR	CG-CD-NE	3.45	122.30	112.19
2	A	750	HEM	CMB-C2B-C3B	3.56	125.41	116.53
3	A	760	H4B	C4-N3-C2	3.61	120.95	115.94
2	B	750	HEM	CAD-C3D-C4D	3.79	125.84	112.47
4	A	800	VUR	CG-CD-NE	3.84	123.43	112.19
2	B	750	HEM	CMC-C2C-C3C	3.94	126.37	116.53
3	A	760	H4B	C2-N1-C8A	4.21	124.00	114.54
3	A	760	H4B	C4-C4A-C8A	4.41	118.55	114.56
2	B	750	HEM	CMB-C2B-C3B	4.62	128.05	116.53
2	A	750	HEM	CMC-C2C-C3C	5.09	129.24	116.53
2	B	750	HEM	CAD-C3D-C2D	5.51	129.05	113.22
2	A	750	HEM	CAD-C3D-C2D	5.52	129.09	113.22
3	B	760	H4B	C4-C4A-C8A	8.70	122.44	114.56

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	800	VUR	NH-C1-NE-CD

There are no ring outliers.

5 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	800	VUR	1	0
2	B	750	HEM	5	0
3	B	760	H4B	1	0
4	B	800	VUR	1	0
5	B	860	ACT	1	0

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	408/422 (96%)	0.10	12 (2%) 55 63	22, 39, 70, 106	0
1	B	411/422 (97%)	-0.02	11 (2%) 58 65	21, 33, 60, 95	0
All	All	819/844 (97%)	0.04	23 (2%) 56 64	21, 36, 68, 106	0

All (23) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	299	ARG	5.8
1	B	300	PHE	5.4
1	B	352	ASP	5.0
1	B	350	THR	4.9
1	B	348	VAL	4.4
1	A	352	ASP	3.9
1	B	718	GLY	3.9
1	A	715	VAL	3.5
1	A	321	THR	3.5
1	A	488	PRO	3.1
1	A	300	PHE	3.0
1	B	389	GLU	2.7
1	B	299	ARG	2.5
1	A	717	LYS	2.5
1	B	351	LYS	2.5
1	A	551	PHE	2.4
1	A	469	LYS	2.3
1	B	349	ARG	2.3
1	A	318	LEU	2.2
1	B	321	THR	2.1
1	B	619	ARG	2.0
1	A	470	HIS	2.0
1	A	373	GLY	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	VUR	A	800	13/13	0.88	0.18	8.16	40,48,54,55	0
5	ACT	A	860	4/4	0.93	0.16	7.36	65,69,70,71	0
4	VUR	B	800	13/13	0.88	0.14	1.89	36,40,47,49	0
5	ACT	B	860	4/4	0.94	0.13	0.77	50,52,53,54	0
2	HEM	A	750	43/43	0.97	0.12	0.73	26,29,35,40	0
6	ZN	B	900	1/1	0.99	0.12	0.56	31,31,31,31	0
2	HEM	B	750	43/43	0.98	0.12	0.46	22,25,37,37	0
3	H4B	A	760	17/17	0.94	0.12	-0.07	26,27,29,29	0
3	H4B	B	760	17/17	0.94	0.12	-0.20	22,23,29,29	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.