



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:24 AM GMT

PDB ID : 3AIB
Title : Crystal Structure of Glucansucrase
Authors : Ito, K.; Ito, S.; Shimamura, T.; Iwata, S.
Deposited on : 2010-05-12
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1114 (3.14-3.06)
Clashscore	102246	1222 (3.14-3.06)
Ramachandran outliers	100387	1174 (3.14-3.06)
Sidechain outliers	100360	1174 (3.14-3.06)
RSRZ outliers	91569	1119 (3.14-3.06)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	844	<div> <div>2%</div> <div>69%</div> <div>27%</div> <div>.</div> </div>
1	B	844	<div> <div>4%</div> <div>52%</div> <div>32%</div> <div>5%</div> <div>11%</div> </div>
1	C	844	<div> <div>73%</div> <div>24%</div> <div>.</div> </div>
1	D	844	<div> <div>2%</div> <div>69%</div> <div>27%</div> <div>.</div> </div>
1	E	844	<div> <div>3%</div> <div>69%</div> <div>27%</div> <div>.</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	844	
1	G	844	
1	H	844	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	MES	D	5001	-	-	-	X
3	MES	E	5001	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 52615 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glucosyltransferase-SI.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	842	Total	C	N	O	S	0	0	0
			6643	4184	1141	1302	16			
1	B	749	Total	C	N	O	S	0	0	0
			5868	3692	1008	1154	14			
1	C	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	D	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	G	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	E	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	F	844	Total	C	N	O	S	0	0	0
			6660	4196	1143	1305	16			
1	H	808	Total	C	N	O	S	0	0	0
			6378	4022	1094	1246	16			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	597	ASP	ASN	SEE REMARK 999	UNP P13470
A	600	LYS	ARG	SEE REMARK 999	UNP P13470
A	727	ILE	THR	SEE REMARK 999	UNP P13470
A	734	VAL	ALA	SEE REMARK 999	UNP P13470
B	597	ASP	ASN	SEE REMARK 999	UNP P13470
B	600	LYS	ARG	SEE REMARK 999	UNP P13470
B	727	ILE	THR	SEE REMARK 999	UNP P13470
B	734	VAL	ALA	SEE REMARK 999	UNP P13470
C	597	ASP	ASN	SEE REMARK 999	UNP P13470
C	600	LYS	ARG	SEE REMARK 999	UNP P13470
C	727	ILE	THR	SEE REMARK 999	UNP P13470
C	734	VAL	ALA	SEE REMARK 999	UNP P13470
D	597	ASP	ASN	SEE REMARK 999	UNP P13470

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	600	LYS	ARG	SEE REMARK 999	UNP P13470
D	727	ILE	THR	SEE REMARK 999	UNP P13470
D	734	VAL	ALA	SEE REMARK 999	UNP P13470
G	597	ASP	ASN	SEE REMARK 999	UNP P13470
G	600	LYS	ARG	SEE REMARK 999	UNP P13470
G	727	ILE	THR	SEE REMARK 999	UNP P13470
G	734	VAL	ALA	SEE REMARK 999	UNP P13470
E	597	ASP	ASN	SEE REMARK 999	UNP P13470
E	600	LYS	ARG	SEE REMARK 999	UNP P13470
E	727	ILE	THR	SEE REMARK 999	UNP P13470
E	734	VAL	ALA	SEE REMARK 999	UNP P13470
F	597	ASP	ASN	SEE REMARK 999	UNP P13470
F	600	LYS	ARG	SEE REMARK 999	UNP P13470
F	727	ILE	THR	SEE REMARK 999	UNP P13470
F	734	VAL	ALA	SEE REMARK 999	UNP P13470
H	597	ASP	ASN	SEE REMARK 999	UNP P13470
H	600	LYS	ARG	SEE REMARK 999	UNP P13470
H	727	ILE	THR	SEE REMARK 999	UNP P13470
H	734	VAL	ALA	SEE REMARK 999	UNP P13470

- Molecule 2 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	G	1	Total Ca 1 1	0	0
2	D	1	Total Ca 1 1	0	0
2	E	1	Total Ca 1 1	0	0
2	H	1	Total Ca 1 1	0	0
2	B	1	Total Ca 1 1	0	0
2	C	1	Total Ca 1 1	0	0
2	A	1	Total Ca 1 1	0	0
2	F	1	Total Ca 1 1	0	0

- Molecule 3 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula: C₆H₁₃NO₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	C	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	D	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	G	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	E	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	F	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
3	H	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 4 is SUGAR (MALTOSE) (three-letter code: MAL) (formula: C₁₂H₂₂O₁₁).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	G	1	Total	C	O	0	0
			23	12	11		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	35	Total	O	0	0
			35	35		
5	B	20	Total	O	0	0
			20	20		
5	C	59	Total	O	0	0
			59	59		
5	D	50	Total	O	0	0
			50	50		
5	G	46	Total	O	0	0
			46	46		
5	E	38	Total	O	0	0
			38	38		
5	F	23	Total	O	0	0
			23	23		
5	H	28	Total	O	0	0
			28	28		


3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Glucosyltransferase-SI




• Molecule 1: Glucosyltransferase-SI






• Molecule 1: Glucosyltransferase-SI


Chain C: 73% 24%




• Molecule 1: Glucosyltransferase-SI




• Molecule 1: Glucosyltransferase-SI




• Molecule 1: Glucosyltransferase-SI




• Molecule 1: Glucosyltransferase-SI





• Molecule 1: Glucosyltransferase-SI



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	295.42Å 213.94Å 220.98Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.67 – 3.09 50.67 – 3.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (50.67-3.09) 97.8 (50.67-3.09)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.58 (at 3.07Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, R_{free}	0.211 , 0.241 0.207 , 0.238	Depositor DCC
R_{free} test set	12657 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	60.4	Xtriage
Anisotropy	0.010	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 29.8	EDS
Estimated twinning fraction	0.000 for -h,l,k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Outliers	0 of 249322 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	52615	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.12 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 9.8896e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: CA, MAL, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.07	9/6784 (0.1%)	0.97	5/9213 (0.1%)
1	B	0.95	6/5990 (0.1%)	0.96	6/8132 (0.1%)
1	C	1.10	11/6802 (0.2%)	0.98	18/9237 (0.2%)
1	D	1.06	9/6802 (0.1%)	0.97	10/9237 (0.1%)
1	E	1.07	6/6802 (0.1%)	0.96	10/9237 (0.1%)
1	F	0.87	2/6802 (0.0%)	0.88	6/9237 (0.1%)
1	G	1.10	15/6802 (0.2%)	0.99	15/9237 (0.2%)
1	H	0.96	3/6516 (0.0%)	0.95	19/8845 (0.2%)
All	All	1.03	61/53300 (0.1%)	0.96	89/72375 (0.1%)

The worst 5 of 61 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	244	SER	CB-OG	8.60	1.53	1.42
1	H	314	ASN	CG-ND2	-8.57	1.11	1.32
1	E	632	GLU	CG-CD	8.44	1.64	1.51
1	G	608	VAL	CB-CG1	7.45	1.68	1.52
1	E	574	GLU	CD-OE1	7.36	1.33	1.25

The worst 5 of 89 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	245	PHE	CB-CA-C	-10.38	89.64	110.40
1	F	1002	ASP	CB-CG-OD1	-8.53	110.63	118.30
1	B	1002	ASP	CB-CG-OD1	8.37	125.84	118.30
1	H	725	ASP	CB-CG-OD1	8.19	125.67	118.30
1	D	425	ARG	NE-CZ-NH1	-7.86	116.37	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6643	0	6475	243	0
1	B	5868	0	5704	333	0
1	C	6660	0	6489	181	0
1	D	6660	0	6489	261	0
1	E	6660	0	6489	314	0
1	F	6660	0	6489	349	0
1	G	6660	0	6489	209	0
1	H	6378	0	6201	272	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	12	0	12	4	0
3	B	12	0	12	1	0
3	C	12	0	12	1	0
3	D	12	0	12	1	0
3	E	12	0	12	1	0
3	F	12	0	12	1	0
3	G	12	0	12	2	0
3	H	12	0	12	1	0
4	G	23	0	22	0	0
5	A	35	0	0	1	0
5	B	20	0	0	0	0
5	C	59	0	0	5	0
5	D	50	0	0	2	0
5	E	38	0	0	0	0
5	F	23	0	0	2	0
5	G	46	0	0	1	0
5	H	28	0	0	0	0
All	All	52615	0	50943	2162	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 2162 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1087:PRO:CG	1:B:1087:PRO:CB	1.77	1.54
1:E:271:TRP:CZ2	1:E:357:ARG:HD3	1.51	1.43
1:C:475:ARG:HB2	1:C:953:MET:CE	1.53	1.39
1:E:277:ILE:CD1	1:E:291:ASP:HB3	1.50	1.38
1:H:278:LEU:HD11	1:H:282:LYS:N	1.41	1.34

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	840/844 (100%)	791 (94%)	45 (5%)	4 (0%)	34	72
1	B	745/844 (88%)	693 (93%)	41 (6%)	11 (2%)	13	46
1	C	842/844 (100%)	809 (96%)	29 (3%)	4 (0%)	34	72
1	D	842/844 (100%)	789 (94%)	46 (6%)	7 (1%)	24	63
1	E	842/844 (100%)	791 (94%)	41 (5%)	10 (1%)	16	52
1	F	842/844 (100%)	783 (93%)	53 (6%)	6 (1%)	26	65
1	G	842/844 (100%)	796 (94%)	43 (5%)	3 (0%)	39	75
1	H	804/844 (95%)	743 (92%)	53 (7%)	8 (1%)	19	58
All	All	6599/6752 (98%)	6195 (94%)	351 (5%)	53 (1%)	24	63

5 of 53 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	713	TYR
1	B	761	ASN
1	D	351	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	281	GLY
1	G	351	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	713/715 (100%)	671 (94%)	42 (6%)	24	60
1	B	628/715 (88%)	575 (92%)	53 (8%)	14	46
1	C	715/715 (100%)	678 (95%)	37 (5%)	29	65
1	D	715/715 (100%)	660 (92%)	55 (8%)	16	50
1	E	715/715 (100%)	665 (93%)	50 (7%)	19	54
1	F	715/715 (100%)	666 (93%)	49 (7%)	19	55
1	G	715/715 (100%)	675 (94%)	40 (6%)	26	62
1	H	683/715 (96%)	645 (94%)	38 (6%)	26	62
All	All	5599/5720 (98%)	5235 (94%)	364 (6%)	21	57

5 of 364 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	740	ASN
1	G	606	ASN
1	H	368	SER
1	D	854	ARG
1	D	1076	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 187 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	864	GLN
1	G	782	GLN
1	H	470	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	1052	ASN
1	G	367	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 17 ligands modelled in this entry, 8 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	MES	A	5001	-	11,12,12	4.03	4 (36%)	14,16,16	12.15	8 (57%)
3	MES	B	5001	-	11,12,12	2.38	3 (27%)	14,16,16	10.36	10 (71%)
3	MES	C	5001	-	11,12,12	3.47	4 (36%)	14,16,16	7.56	5 (35%)
3	MES	D	5001	-	11,12,12	2.80	4 (36%)	14,16,16	7.65	10 (71%)
3	MES	E	5001	-	11,12,12	3.02	3 (27%)	14,16,16	9.05	9 (64%)
3	MES	F	5001	-	11,12,12	3.38	4 (36%)	14,16,16	8.68	9 (64%)
3	MES	G	5001	-	11,12,12	2.92	3 (27%)	14,16,16	9.45	8 (57%)
4	MAL	G	5044	-	24,24,24	2.04	5 (20%)	35,35,35	3.24	15 (42%)
3	MES	H	5001	-	11,12,12	2.81	5 (45%)	14,16,16	7.93	9 (64%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MES	A	5001	-	-	0/6/14/14	0/1/1/1
3	MES	B	5001	-	-	0/6/14/14	0/1/1/1
3	MES	C	5001	-	-	0/6/14/14	0/1/1/1
3	MES	D	5001	-	-	0/6/14/14	0/1/1/1
3	MES	E	5001	-	-	0/6/14/14	0/1/1/1
3	MES	F	5001	-	-	0/6/14/14	0/1/1/1
3	MES	G	5001	-	-	0/6/14/14	0/1/1/1
4	MAL	G	5044	-	-	0/8/48/48	0/2/2/2
3	MES	H	5001	-	-	0/6/14/14	0/1/1/1

The worst 5 of 35 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	H	5001	MES	C7-N4	-2.23	1.42	1.47
3	F	5001	MES	C7-C8	2.01	1.59	1.52
3	H	5001	MES	C5-C6	2.07	1.58	1.50
3	C	5001	MES	C5-N4	2.22	1.53	1.46
3	A	5001	MES	C5-N4	2.23	1.53	1.46

The worst 5 of 83 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	5001	MES	O1S-S-C8	-33.66	78.18	106.91
3	B	5001	MES	O2S-S-C8	-30.49	80.89	106.91
3	G	5001	MES	O1S-S-C8	-29.18	82.01	106.91
3	A	5001	MES	O2S-S-C8	-28.21	82.84	106.91
3	E	5001	MES	O1S-S-C8	-26.71	84.11	106.91

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	5001	MES	4	0
3	B	5001	MES	1	0
3	C	5001	MES	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	5001	MES	1	0
3	E	5001	MES	1	0
3	F	5001	MES	1	0
3	G	5001	MES	2	0
3	H	5001	MES	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	842/844 (99%)	-0.21	18 (2%) 67 44	23, 36, 90, 112	0
1	B	749/844 (88%)	-0.01	34 (4%) 37 17	30, 51, 98, 148	0
1	C	844/844 (100%)	-0.33	4 (0%) 91 83	25, 35, 61, 71	0
1	D	844/844 (100%)	-0.26	15 (1%) 71 50	25, 37, 89, 110	0
1	E	844/844 (100%)	-0.17	25 (2%) 54 29	22, 36, 111, 136	0
1	F	844/844 (100%)	-0.14	6 (0%) 89 78	39, 55, 78, 91	0
1	G	844/844 (100%)	-0.34	4 (0%) 91 83	24, 36, 72, 85	0
1	H	808/844 (95%)	-0.17	26 (3%) 51 27	25, 43, 101, 144	0
All	All	6619/6752 (98%)	-0.21	132 (1%) 68 46	22, 40, 88, 148	0

The worst 5 of 132 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1087	PRO	5.9
1	E	1081	PRO	5.5
1	H	1087	PRO	5.1
1	H	277	ILE	4.9
1	A	1087	PRO	4.8

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	MES	D	5001	12/12	0.86	0.27	4.86	32,34,36,36	0
3	MES	E	5001	12/12	0.88	0.24	3.68	32,35,37,37	0
3	MES	H	5001	12/12	0.91	0.21	1.49	30,32,34,35	0
3	MES	B	5001	12/12	0.92	0.23	1.39	39,41,43,44	0
3	MES	C	5001	12/12	0.90	0.20	1.33	30,31,32,33	0
3	MES	G	5001	12/12	0.91	0.19	1.17	29,31,32,33	0
3	MES	F	5001	12/12	0.90	0.22	1.08	38,38,39,39	0
4	MAL	G	5044	23/23	0.94	0.18	0.94	30,34,38,40	0
2	CA	B	4001	1/1	0.96	0.19	0.67	32,32,32,32	0
2	CA	G	4001	1/1	0.96	0.17	-0.09	28,28,28,28	0
2	CA	E	4001	1/1	0.91	0.17	-0.58	32,32,32,32	0
2	CA	F	4001	1/1	0.93	0.17	-0.65	40,40,40,40	0
2	CA	H	4001	1/1	0.97	0.16	-0.68	32,32,32,32	0
2	CA	C	4001	1/1	0.89	0.17	-0.88	29,29,29,29	0
2	CA	D	4001	1/1	0.96	0.15	-1.21	30,30,30,30	0
2	CA	A	4001	1/1	0.96	0.15	-1.42	29,29,29,29	0
3	MES	A	5001	12/12	0.85	0.26	-	31,33,33,34	0

6.5 Other polymers

There are no such residues in this entry.