



# Full wwPDB X-ray Structure Validation Report ⓘ

May 5, 2016 – 07:42 PM EDT

PDB ID : 5AID  
Title : Crystal structure of the Mep2 mutant delta442 from *Candida albicans*  
Authors : van den Berg, B.; Chembath, A.; Rutherford, J.  
Deposited on : 2015-02-12  
Resolution : 3.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027457  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027457


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 3.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1476 (3.50-3.30)
Clashscore	102246	1611 (3.50-3.30)
Ramachandran outliers	100387	1571 (3.50-3.30)
Sidechain outliers	100360	1571 (3.50-3.30)
RSRZ outliers	91569	1485 (3.50-3.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	448	<div> <div>12%</div> <div>57%</div> <div>29%</div> <div>•</div> <div>11%</div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3041 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MEP2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	398	3041	2008	488	517	28	7	3	0


There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	4	GLN	ASN	CONFLICT	UNP Q59UP8
A	443	HIS	-	EXPRESSION TAG	UNP Q59UP8
A	444	HIS	-	EXPRESSION TAG	UNP Q59UP8
A	445	HIS	-	EXPRESSION TAG	UNP Q59UP8
A	446	HIS	-	EXPRESSION TAG	UNP Q59UP8
A	447	HIS	-	EXPRESSION TAG	UNP Q59UP8
A	448	HIS	-	EXPRESSION TAG	UNP Q59UP8

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: MEP2



## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	114.73Å 114.73Å 289.96Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	96.66 – 3.40 96.65 – 3.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (96.66-3.40) 92.3 (96.65-3.40)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.33 (at 3.41Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.236 , 0.289 0.233 , 0.282	Depositor DCC
$R_{free}$ test set	983 reflections (10.18%)	DCC
Wilson B-factor (Å <sup>2</sup> )	127.7	Xtriage
Anisotropy	0.107	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 86.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for $-1/3^*h+1/3^*k+1/3^*l,-k,8/3^*h+4/3^*k+1/3^*l$ 0.025 for $-2/3^*h-1/3^*k-1/3^*l,-1/3^*h-2/3^*k+1/3^*l,-4/3^*h+4/3^*k+1/3^*l$ 0.015 for $-h,1/3^*h-1/3^*k-1/3^*l,-4/3^*h-8/3^*k+1/3^*l$	Xtriage
$F_o, F_c$ correlation	0.75	EDS
Total number of atoms	3041	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	131.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.71% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.45	0/3144	0.63	2/4287 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	260	LEU	CA-CB-CG	8.77	135.48	115.30
1	A	61	LEU	CA-CB-CG	5.98	129.06	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3041	0	2992	84	0
All	All	3041	0	2992	84	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:13:ASP:HB3	1:A:16:LYS:HB3	1.67	0.77

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:THR:HG21	1:A:310:VAL:HG13	1.74	0.69
1:A:232:LEU:O	1:A:236:TRP:HD1	1.74	0.69
1:A:143:ARG:HD3	1:A:413:ARG:HB2	1.75	0.69
1:A:4:GLN:HA	1:A:17:VAL:HG11	1.77	0.66
1:A:34:ALA:HA	1:A:37:LEU:HD12	1.79	0.65
1:A:57:HIS:HE2	1:A:139[A]:CYS:HG	1.25	0.64
1:A:380:GLN:HA	1:A:383:TYR:HD2	1.62	0.63
1:A:356:PHE:HA	1:A:374:MET:HG2	1.80	0.63
1:A:167:TRP:HZ2	1:A:183:GLY:HA3	1.64	0.62
1:A:89:GLY:HA2	1:A:95:THR:HG23	1.84	0.59
1:A:27:ASP:OD1	1:A:247:SER:OG	2.17	0.58
1:A:295:VAL:HG11	1:A:342:HIS:HA	1.87	0.57
1:A:276:ARG:NH1	1:A:325:ASP:OD2	2.38	0.56
1:A:396:PHE:O	1:A:400:SER:OG	2.14	0.56
1:A:104:VAL:HG13	1:A:116:ASP:HB3	1.88	0.56
1:A:141:ARG:HH21	1:A:416:LEU:HD12	1.69	0.56
1:A:219:PRO:O	1:A:220:HIS:ND1	2.37	0.55
1:A:201:TRP:HB3	1:A:404:LEU:HD21	1.89	0.55
1:A:407:ASP:HB2	1:A:414:ILE:HD11	1.87	0.55
1:A:61:LEU:HA	1:A:64:ALA:HB3	1.88	0.54
1:A:399:THR:O	1:A:403:LEU:HG	2.09	0.53
1:A:236:TRP:CD1	1:A:294:LEU:HD13	2.44	0.52
1:A:89:GLY:HA3	1:A:94:GLY:HA2	1.92	0.51
1:A:57:HIS:NE2	1:A:139[A]:CYS:SG	2.65	0.51
1:A:288:MET:HE2	1:A:341:LEU:HD21	1.91	0.51
1:A:351:PHE:CE1	1:A:381:LEU:HD11	2.46	0.50
1:A:199:SER:O	1:A:332:ILE:HD13	2.11	0.50
1:A:50:SER:N	1:A:61:LEU:HD11	2.27	0.49
1:A:277:THR:O	1:A:279:GLY:N	2.45	0.49
1:A:262:ALA:HB2	1:A:294:LEU:HD21	1.94	0.49
1:A:357:ALA:O	1:A:372:GLY:N	2.46	0.49
1:A:380:GLN:NE2	1:A:384:GLN:OE1	2.45	0.48
1:A:48:LEU:HD23	1:A:49:TYR:CE1	2.49	0.48
1:A:223:SER:O	1:A:227:MET:N	2.41	0.48
1:A:240:ASN:ND2	1:A:258:THR:OG1	2.46	0.48
1:A:225:ILE:O	1:A:229:THR:N	2.41	0.47
1:A:362:ALA:O	1:A:367:THR:HB	2.15	0.47
1:A:52:ILE:HA	1:A:225:ILE:HD11	1.97	0.47
1:A:305:PRO:HB2	1:A:308:THR:HG23	1.96	0.47
1:A:262:ALA:HB2	1:A:294:LEU:CD2	2.45	0.46
1:A:398:VAL:HA	1:A:401:ILE:HG22	1.98	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:90:SER:O	1:A:170:ASN:HB2	2.15	0.46
1:A:266:GLY:HA2	1:A:289:GLY:HA3	1.96	0.46
1:A:407:ASP:CB	1:A:414:ILE:HD11	2.46	0.46
1:A:377:HIS:ND1	1:A:380:GLN:HB2	2.31	0.45
1:A:398:VAL:O	1:A:402:ILE:HG12	2.15	0.45
1:A:236:TRP:HZ3	1:A:298:THR:HG1	1.62	0.45
1:A:165:TRP:O	1:A:173:LEU:HB2	2.16	0.45
1:A:88:ASN:ND2	1:A:95:THR:HG21	2.31	0.45
1:A:188:HIS:O	1:A:192:GLY:N	2.51	0.44
1:A:76:TRP:HA	1:A:80:GLY:HA3	1.99	0.43
1:A:141:ARG:NH2	1:A:416:LEU:HD12	2.34	0.43
1:A:119:PHE:O	1:A:123:GLN:HB2	2.18	0.43
1:A:4:GLN:HG2	1:A:16:LYS:NZ	2.33	0.43
1:A:358:ALA:HB1	1:A:360:TYR:CE2	2.54	0.43
1:A:371:GLY:HA3	1:A:375:ASN:HD21	1.84	0.43
1:A:79:TRP:CZ2	1:A:152:PHE:HE1	2.36	0.43
1:A:91:VAL:HA	1:A:170:ASN:O	2.19	0.43
1:A:232:LEU:O	1:A:236:TRP:CD1	2.64	0.42
1:A:154:TRP:HD1	1:A:402:ILE:HD11	1.84	0.42
1:A:167:TRP:CZ2	1:A:183:GLY:HA3	2.50	0.42
1:A:320:CYS:HA	1:A:323:ALA:HB3	2.02	0.42
1:A:264:THR:OG1	1:A:313:GLY:HA3	2.20	0.42
1:A:31:ILE:HB	1:A:118:LEU:HD21	2.02	0.42
1:A:47:LEU:HD13	1:A:227:MET:HG3	2.02	0.42
1:A:393:ALA:O	1:A:397:THR:HG23	2.19	0.42
1:A:25:ARG:HD3	1:A:28:MET:CE	2.50	0.41
1:A:78:TRP:CD1	1:A:79:TRP:CE3	3.08	0.41
1:A:162:ILE:HG23	1:A:391:VAL:HG22	2.02	0.41
1:A:145:GLY:O	1:A:149:VAL:HG23	2.21	0.41
1:A:277:THR:O	1:A:280:LYS:N	2.52	0.41
1:A:291:ILE:O	1:A:295:VAL:HG23	2.21	0.41
1:A:414:ILE:CD1	1:A:415:ARG:H	2.34	0.41
1:A:48:LEU:O	1:A:52:ILE:HB	2.21	0.41
1:A:50:SER:H	1:A:61:LEU:HD11	1.85	0.41
1:A:18:ASP:O	1:A:21:GLU:HG3	2.21	0.41
1:A:111:VAL:HG12	1:A:113:THR:HG22	2.02	0.40
1:A:225:ILE:HD13	1:A:284:VAL:HG13	2.04	0.40
1:A:330:LEU:O	1:A:331:GLN:HB3	2.22	0.40
1:A:375:ASN:OD1	1:A:375:ASN:N	2.53	0.40
1:A:96:LEU:HA	1:A:96:LEU:HD23	1.78	0.40
1:A:269:TRP:CH2	1:A:288:MET:HE1	2.57	0.40

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:402:ILE:O	1:A:406:MET:HB2	2.22	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	395/448 (88%)	334 (85%)	49 (12%)	12 (3%)	5	39

All (12) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	SER
1	A	278	GLY
1	A	413	ARG
1	A	220	HIS
1	A	414	ILE
1	A	78	TRP
1	A	334	ASP
1	A	141	ARG
1	A	60	SER
1	A	330	LEU
1	A	410	PRO
1	A	314	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/346 (89%)	287 (94%)	20 (6%)	21	61

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	40	ILE
1	A	57	HIS
1	A	59	LEU
1	A	61	LEU
1	A	79	TRP
1	A	110	ILE
1	A	166	THR
1	A	174	VAL
1	A	191	SER
1	A	218	LYS
1	A	260	LEU
1	A	270	MET
1	A	276	ARG
1	A	328	ASP
1	A	329	LEU
1	A	379	LYS
1	A	401	ILE
1	A	403	LEU
1	A	412	LEU
1	A	414	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	398/448 (88%)	0.67	55 (13%) 4 4	81, 126, 187, 228	2 (0%)

All (55) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	4	GLN	9.9
1	A	333	ASP	9.6
1	A	7	GLY	7.9
1	A	54	ARG	6.6
1	A	6	THR	6.2
1	A	55	LYS	6.2
1	A	11	GLY	5.6
1	A	8	THR	5.5
1	A	5	PHE	5.5
1	A	103	ASP	5.3
1	A	231	PHE	5.1
1	A	40	ILE	4.7
1	A	12	GLY	4.5
1	A	10	THR	4.3
1	A	332	ILE	4.3
1	A	234	PHE	4.2
1	A	9	GLY	3.9
1	A	104	VAL	3.6
1	A	201	TRP	3.6
1	A	237	TYR	3.6
1	A	230	ILE	3.5
1	A	66	LEU	3.4
1	A	404	LEU	3.3
1	A	36	VAL	3.2
1	A	37	LEU	3.1
1	A	43	PRO	3.0
1	A	306	VAL	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	182	ALA	2.9
1	A	102	LYS	2.9
1	A	67	MET	2.8
1	A	92	PHE	2.8
1	A	110	ILE	2.8
1	A	105	LEU	2.7
1	A	17	VAL	2.7
1	A	334	ASP	2.6
1	A	360	TYR	2.6
1	A	278	GLY	2.6
1	A	16	LYS	2.6
1	A	240	ASN	2.5
1	A	41	MET	2.5
1	A	39	TRP	2.5
1	A	62[A]	MET	2.5
1	A	14	VAL	2.5
1	A	65	ALA	2.4
1	A	227	MET	2.4
1	A	101	LEU	2.4
1	A	107	ALA	2.2
1	A	309[A]	SER	2.1
1	A	279	GLY	2.1
1	A	19	LEU	2.1
1	A	244	THR	2.1
1	A	112	LYS	2.1
1	A	123	GLN	2.1
1	A	126	PHE	2.0
1	A	68	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.