



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 19, 2017 – 11:15 AM EST

PDB ID : 5AOH
Title : Crystal Structure of CarF
Authors : Tichy, E.M.; Hardwick, S.W.; Luisi, B.F.; C Salmond, G.P.
Deposited on : 2015-09-10
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4533 (1.80-1.80)
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	273	
1	B	273	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	K	B	1267	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4290 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CARF.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	0	1	0
			2001	1266	340	386	9			
1	B	261	Total	C	N	O	S	0	0	0
			1982	1253	338	382	9			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	269	HIS	-	EXPRESSION TAG	UNP V3T1Y6
A	270	HIS	-	EXPRESSION TAG	UNP V3T1Y6
A	271	HIS	-	EXPRESSION TAG	UNP V3T1Y6
A	272	HIS	-	EXPRESSION TAG	UNP V3T1Y6
A	273	HIS	-	EXPRESSION TAG	UNP V3T1Y6
B	269	HIS	-	EXPRESSION TAG	UNP V3T1Y6
B	270	HIS	-	EXPRESSION TAG	UNP V3T1Y6
B	271	HIS	-	EXPRESSION TAG	UNP V3T1Y6
B	272	HIS	-	EXPRESSION TAG	UNP V3T1Y6
B	273	HIS	-	EXPRESSION TAG	UNP V3T1Y6

- Molecule 2 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	K	0	0
			1	1		
2	A	1	Total	K	0	0
			1	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	165	Total	O	0	0
			165	165		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	140	Total	O	0	0
			140	140		

- Molecule 1: CARF



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	49.47Å 55.45Å 176.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	88.10 – 1.80 36.13 – 1.80	Depositor EDS
% Data completeness (in resolution range)	97.2 (88.10-1.80) 97.2 (36.13-1.80)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.44 (at 1.81Å)	Xtriage
Refinement program	REFMAC 5.8.0131	Depositor
R, R_{free}	0.137 , 0.208 0.147 , 0.213	Depositor DCC
R_{free} test set	2185 reflections (5.15%)	DCC
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.969	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 50.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	4290	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section:
K

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	1/2060 (0.0%)	0.97	4/2809 (0.1%)
1	B	1.09	2/2037 (0.1%)	0.98	10/2777 (0.4%)
All	All	1.08	3/4097 (0.1%)	0.97	14/5586 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67	CYS	CB-SG	-12.98	1.60	1.82
1	B	67	CYS	CB-SG	-9.02	1.67	1.82
1	B	7	GLU	CD-OE2	-5.15	1.20	1.25

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	CYS	CB-CA-C	-6.56	97.28	110.40
1	B	246	ASP	CB-CG-OD1	6.48	124.14	118.30
1	A	67	CYS	CA-CB-SG	-6.23	102.79	114.00
1	B	254	MET	CG-SD-CE	6.04	109.87	100.20
1	B	142	ARG	NE-CZ-NH2	-6.04	117.28	120.30
1	B	64	SER	N-CA-C	5.97	127.13	111.00
1	B	67	CYS	CA-CB-SG	-5.85	103.47	114.00
1	B	67	CYS	CB-CA-C	-5.72	98.96	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	246	ASP	CB-CG-OD2	-5.69	113.18	118.30
1	B	74	ASP	CB-CG-OD1	5.62	123.36	118.30
1	A	74	ASP	CB-CG-OD2	-5.39	113.45	118.30
1	B	84	ARG	NE-CZ-NH1	5.32	122.96	120.30
1	A	254	MET	CB-CG-SD	-5.09	97.14	112.40
1	B	64	SER	CB-CA-C	-5.06	100.48	110.10

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	224	GLY	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2001	0	1859	20	1
1	B	1982	0	1844	14	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	165	0	0	4	1
3	B	140	0	0	0	0
All	All	4290	0	3703	34	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:125:GLU:O	1:A:128:THR:HG22	1.72	0.90
1:B:25:ASP:OD2	1:B:28:ALA:HB2	1.79	0.82
1:B:25:ASP:OD2	1:B:28:ALA:CB	2.28	0.81
1:A:56:ALA:HB1	1:A:61:TYR:HB2	1.62	0.80

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2:ILE:O	1:B:2:ILE:HD12	1.89	0.73
1:A:118:GLN:HB3	3:A:2073:HOH:O	1.89	0.72
1:A:116:ASP:OD1	1:A:118:GLN:HG2	1.90	0.71
1:B:25:ASP:OD2	1:B:28:ALA:HB3	2.03	0.58
1:A:46:TYR:CE2	1:A:50:GLN:OE1	2.57	0.58
1:A:64:SER:HB2	3:A:2051:HOH:O	2.02	0.58
1:A:81:ASP:OD2	1:A:85:HIS:HE1	1.87	0.57
1:A:31:ASN:OD1	1:A:32:THR:HG23	2.05	0.56
1:A:124:PRO:HB2	1:A:128:THR:HG23	1.88	0.56
1:A:36:PRO:HB2	1:A:265:ILE:HG13	1.88	0.56
1:B:56:ALA:HB1	1:B:61:TYR:HB2	1.89	0.54
1:A:186:THR:HG22	1:A:188:THR:H	1.73	0.53
1:A:113:VAL:HB	1:A:118:GLN:O	2.09	0.52
1:B:81:ASP:OD2	1:B:85:HIS:HE1	1.92	0.52
1:A:68:ASN:ND2	1:A:85:HIS:HB2	2.26	0.50
1:B:125:GLU:O	1:B:126:GLU:CB	2.59	0.50
1:A:172:LYS:HE2	1:A:174:GLN:HE21	1.76	0.49
1:A:64:SER:CB	3:A:2051:HOH:O	2.59	0.49
1:B:85:HIS:HD2	1:B:191:THR:OG1	1.95	0.49
1:B:254:MET:CE	1:B:255:SER:H	2.26	0.48
1:A:85:HIS:HD2	1:A:191:THR:OG1	1.97	0.47
1:B:252:PHE:CE2	1:B:254:MET:HG2	2.50	0.46
1:B:215:LEU:HD11	1:B:231:CYS:SG	2.55	0.46
1:A:215:LEU:HD11	1:A:231:CYS:SG	2.56	0.45
1:B:31:ASN:OD1	1:B:32:THR:HG23	2.17	0.45
1:A:168:ARG:HD3	3:A:2102:HOH:O	2.17	0.44
1:A:105:GLN:HB3	1:A:106:HIS:HD2	1.83	0.43
1:B:114:THR:HG22	1:B:132:ILE:HD12	2.02	0.42
1:A:25:ASP:O	1:A:29:HIS:HD2	2.02	0.41
1:B:7:GLU:HB3	1:B:38:SER:HB2	2.03	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:2051:HOH:O	3:A:2080:HOH:O[4_455]	1.85	0.35
1:A:225:LYS:NZ	1:B:74:ASP:OD2[1_565]	2.18	0.02

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	264/273 (97%)	259 (98%)	4 (2%)	1 (0%)	39	23
1	B	257/273 (94%)	250 (97%)	6 (2%)	1 (0%)	39	23
All	All	521/546 (95%)	509 (98%)	10 (2%)	2 (0%)	39	23

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	210	ASN
1	B	210	ASN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	202/219 (92%)	199 (98%)	3 (2%)	72	62
1	B	202/219 (92%)	202 (100%)	0	100	100
All	All	404/438 (92%)	401 (99%)	3 (1%)	88	86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	A	183	VAL
1	A	265	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	HIS
1	A	68	ASN
1	A	85	HIS
1	A	106	HIS
1	A	174	GLN
1	A	216	ASN
1	B	85	HIS
1	B	105	GLN
1	B	187	GLN
1	B	216	ASN
1	B	248	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	265/273 (97%)	-0.29	6 (2%) 64 59	11, 24, 46, 75	0
1	B	261/273 (95%)	-0.35	3 (1%) 82 80	13, 23, 42, 61	0
All	All	526/546 (96%)	-0.32	9 (1%) 73 69	11, 24, 46, 75	0

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	224	GLY	5.3
1	B	224	GLY	5.0
1	B	223	GLY	4.2
1	B	131	GLY	3.3
1	A	223	GLY	3.2
1	A	2	ILE	2.9
1	A	107	ASN	2.2
1	A	3	LEU	2.1
1	A	128	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron

density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	K	B	1267	1/1	0.99	0.13	2.03	18,18,18,18	0
2	K	A	1267	1/1	1.00	0.12	1.40	16,16,16,16	0

6.5 Other polymers [i](#)

There are no such residues in this entry.