wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 01:55 PM BST

PDB ID	:	1ARQ
Title	:	RELAXATION MATRIX REFINEMENT OF THE SOLUTION STRUCTURE OF THE ARC REPRESSOR
Authors	:	Bonvin, A.M.J.J.; Vis, H.; Burgering, M.J.M.; Breg, J.N.; Boelens, R.; Kaptein, R.
Deposited on	:	1993-08-24

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	53	36% 28% 9% 26%
1	B	53	40% 25% 9% 26%

2 Ensemble composition and analysis (i)

This entry contains 16 models. Model 2 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:8-A:46, B:8-B:46 (78)	0.33	2

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 3 single-model clusters were found.

Cluster number	Models
1	2, 4, 7, 8, 11, 13, 15
2	5, 12, 14, 16
3	1, 6
Single-model clusters	3; 9; 10

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 870 atoms, of which 0 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called ARC REPRESSOR.

Mol	Chain	Residues	Atoms					Trace
			Total	C	N	O	S	
1	A	53	435	271	82	78	4	0
1	B	53	435	271	82	78	4	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: ARC REPRESSOR

Chain A: 36% 28% 9% 26%

M1 K2 G3 M4 S5 K6 M7 P8 Q9 F10 N11 L12 R13 - R16 E17 V18 L19 D20 L21 V22 R23 K24 V25 - R31 S32 - S35 E36 I37 Y38 Q39 R40 - F43 S44 F45 K46 K47 E48 G49 R50 I51 G52 A53

- Molecule 1: ARC REPRESSOR

Chain B: 40% 25% 9% 26%

M1 K2 G3 M4 S5 K6 M7 P8 Q9 F10 - R13 - R16 E17 V18 L19 - V22 R23 K24 V25 - N29 G30 R31 S32 V33 - E36 I37 Y38 Q39 R40 - F45 K46 K47 E48 G49 R50 I51 G52 A53

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 2. Colouring as in section 4.1 above.

- Molecule 1: ARC REPRESSOR

Chain A: 36% 21% 15% • 26%

M1 K2 G3 M4 S5 K6 M7 P8 Q9 F10 N11 L12 R13 W14 P15 R16 - L19 D20 L21 V22 R23 K24 V25 - R31 S32 - S35 E36 I37 Y38 Q39 R40 V41 - F45 K46 K47 E48 G49 R50 I51 G52 A53

- Molecule 1: ARC REPRESSOR

Chain B: 47% 19% 8% 26%

M1 K2 G3 M4 S5 K6 M7 P8 Q9 F10 N11 - R16 - L19 - R23 K24 V25 - R31 - I37 Y38 Q39 R40 - E43 S44 F45 K46 K47 E48 G49 R50 I51 G52 A53

5 Refinement protocol and experimental data overview (i)

Of the ? calculated structures, 16 were deposited, based on the following criterion: *?*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
IRMA	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.01±0.03	0±0/339 (0.0±0.1%)	2.26±0.16	17±3/456 (3.7±0.6%)
1	B	0.88±0.02	0±0/339 (0.0±0.0%)	1.68±0.07	7±2/456 (1.6±0.3%)
All	All	0.95	1/10848 (0.0%)	2.00	383/14592 (2.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.8±0.4	6.1±0.9
1	B	0.0±0.0	5.2±1.2
All	All	13	182

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	A	10	PHE	CE2-CZ	5.03	1.47	1.37	16	1

5 of 82 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	38	TYR	CB-CG-CD1	-30.32	102.81	121.00	8	16
1	A	10	PHE	CB-CG-CD2	-28.79	100.65	120.80	15	13
1	A	45	PHE	CB-CG-CD2	-15.81	109.73	120.80	14	11
1	B	38	TYR	CB-CG-CD2	-15.65	111.61	121.00	12	2
1	B	45	PHE	CB-CG-CD2	-15.64	109.85	120.80	8	11

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	16	ARG	CA	13

5 of 30 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	38	TYR	Sidechain	16
1	A	45	PHE	Sidechain	15
1	A	23	ARG	Mainchain	15
1	B	45	PHE	Sidechain	15
1	B	38	TYR	Sidechain	14

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	332	0	331	5±2
1	B	332	0	331	6±2
All	All	10624	0	10592	132

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

5 of 76 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:39:GLN:HE22	1:B:40:ARG:CZ	0.69	2.00	1	9
1:A:11:ASN:HD21	1:B:9:GLN:HB2	0.67	1.49	2	3
1:A:39:GLN:HE22	1:A:40:ARG:CZ	0.66	2.04	4	6
1:A:31:ARG:NH1	1:A:39:GLN:HE22	0.61	1.93	12	4
1:B:39:GLN:HE21	1:B:40:ARG:NH2	0.58	1.96	7	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	39/53 (74%)	36±1 (92±3%)	2±1 (6±3%)	1±1 (2±2%)	12 49
1	B	39/53 (74%)	36±1 (93±3%)	2±1 (4±2%)	1±1 (3±2%)	9 42
All	All	1248/1696 (74%)	1152 (92%)	62 (5%)	34 (3%)	10 45

5 of 9 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	B	32	SER	7
1	B	8	PRO	6
1	B	33	VAL	5
1	A	46	LYS	4
1	A	33	VAL	4

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	37/47 (79%)	34±1 (91±3%)	4±1 (9±3%)	15 60
1	B	37/47 (79%)	33±1 (90±4%)	4±1 (10±4%)	14 59
All	All	1184/1504 (79%)	1070 (90%)	114 (10%)	15 60

5 of 27 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	13	ARG	15
1	B	13	ARG	13
1	B	36	GLU	9
1	B	39	GLN	8
1	A	43	GLU	7

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided