



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 02:52 PM GMT

PDB ID : 4ATN  
Title : Crystal structure of C2498 2'-O-ribose methyltransferase RlmM from Escherichia coli  
Authors : Punekar, A.S.; Shepherd, T.R.; Liljeruhm, J.; Forster, A.C.; Selmer, M.  
Deposited on : 2012-05-08  
Resolution : 1.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	375	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1359	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	1362	-	-	-	X
3	EDO	A	1365	-	-	-	X
3	EDO	A	1366	-	-	-	X
3	EDO	A	1367	-	-	-	X
3	EDO	A	1372	-	-	-	X
3	EDO	A	1373	-	-	-	X
3	EDO	A	1377	-	-	-	X
3	EDO	A	1378	-	-	-	X

## 2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3245 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called RIBOSOMAL RNA LARGE SUBUNIT METHYLTRANSFERASE M.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	357	Total	C	N	O	S	0	7	0
			2916	1872	502	519	23			

There are 9 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	367	SER	-	EXPRESSION TAG	UNP P0ADR6
A	368	LYS	-	EXPRESSION TAG	UNP P0ADR6
A	369	GLY	-	EXPRESSION TAG	UNP P0ADR6
A	370	HIS	-	EXPRESSION TAG	UNP P0ADR6
A	371	HIS	-	EXPRESSION TAG	UNP P0ADR6
A	372	HIS	-	EXPRESSION TAG	UNP P0ADR6
A	373	HIS	-	EXPRESSION TAG	UNP P0ADR6
A	374	HIS	-	EXPRESSION TAG	UNP P0ADR6
A	375	HIS	-	EXPRESSION TAG	UNP P0ADR6

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		
2	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula:  $C_2H_6O_2$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0
3	A	1	Total C O 4 2 2	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	195	Total	O	0	0
			195	195		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RIBOSOMAL RNA LARGE SUBUNIT METHYLTRANSFERASE M





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	92.09 Å 92.09 Å 83.09 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.77 – 1.95 28.77 – 1.95	Depositor EDS
% Data completeness (in resolution range)	99.8 (28.77-1.95) 99.8 (28.77-1.95)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.30 (at 1.95 Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, $R_{free}$	0.169 , 0.206 0.162 , 0.200	Depositor DCC
$R_{free}$ test set	1501 reflections (5.00%)	DCC
Wilson B-factor (Å <sup>2</sup> )	33.0	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.1	EDS
Estimated twinning fraction	0.034 for -h,-k,l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 30025 reflections	Xtriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	3245	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.67% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, SO4, CSS, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.40	0/3002	0.52	0/4068

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2916	0	2893	53	0
2	A	30	0	40	4	0
3	A	84	0	126	11	0
4	A	20	0	0	1	0
5	A	195	0	0	5	0
All	All	3245	0	3059	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:260:ASP:H	3:A:1371:EDO:H22	1.37	0.85
1:A:309:MET:HB2	3:A:1368:EDO:H12	1.56	0.85
1:A:61:PHE:H	1:A:167:ASN:HD21	1.34	0.74
4:A:1387:SO4:O4	5:A:2122:HOH:O	2.07	0.72
1:A:166:ASN:H	1:A:166:ASN:HD22	1.37	0.72
1:A:335:GLN:HE22	1:A:353:ARG:HH21	1.38	0.68
1:A:69:GLN:HE22	1:A:169:SER:H	1.44	0.67
1:A:61:PHE:H	1:A:167:ASN:ND2	1.97	0.63
1:A:241:ASN:O	1:A:258:ARG:NH1	2.33	0.61
1:A:69:GLN:HE21	1:A:70:TRP:H	1.48	0.61
1:A:270:ASN:H	1:A:270:ASN:HD22	1.47	0.60
1:A:117[A]:LEU:CD1	1:A:121:ARG:NH2	2.65	0.59
1:A:268:ARG:HH11	1:A:270:ASN:HD21	1.51	0.58
1:A:60:PRO:HB3	3:A:1364:EDO:H12	1.86	0.58
1:A:69:GLN:NE2	1:A:169:SER:H	2.00	0.57
1:A:107:VAL:HG22	1:A:117[A]:LEU:HD23	1.86	0.57
1:A:337:GLN:HE21	1:A:351:HIS:HD2	1.51	0.57
3:A:1375:EDO:O1	5:A:2195:HOH:O	2.17	0.56
1:A:106:GLU:OE1	1:A:149:HIS:HD2	1.90	0.55
1:A:337:GLN:HG3	5:A:2185:HOH:O	2.09	0.53
1:A:59:LEU:HD12	1:A:60:PRO:HD2	1.91	0.53
1:A:328:ASP:OD2	5:A:2183:HOH:O	2.19	0.52
1:A:15:GLU:OE2	1:A:312:ARG:HD2	2.11	0.51
1:A:50:ASP:HB2	3:A:1372:EDO:H21	1.93	0.50
1:A:69:GLN:HE22	1:A:168:ASN:HB2	1.77	0.50
1:A:72:VAL:O	1:A:161:GLY:HA3	2.12	0.49
1:A:77:LEU:HB2	1:A:92[B]:MET:SD	2.53	0.49
1:A:176:PRO:HG2	1:A:194:GLU:HG3	1.94	0.49
1:A:97:VAL:HA	2:A:1360:GOL:H11	1.95	0.48
1:A:58:GLU:OE1	5:A:2030:HOH:O	2.20	0.48
1:A:51:ASP:N	3:A:1372:EDO:H12	2.29	0.48
1:A:312:ARG:NH2	1:A:346[A]:GLU:OE2	2.47	0.48
1:A:320:LEU:HD13	2:A:1361:GOL:H11	1.96	0.48
1:A:149:HIS:HE1	1:A:163:SER:OG	1.99	0.46
1:A:171:PHE:CE1	2:A:1362:GOL:H2	2.51	0.46
1:A:166:ASN:HD21	3:A:1376:EDO:H12	1.80	0.46
1:A:88:PRO:O	1:A:92[B]:MET:HG3	2.16	0.46
1:A:211:ASN:HD22	1:A:211:ASN:C	2.20	0.45
1:A:117[A]:LEU:HD11	1:A:121:ARG:NH2	2.31	0.45
1:A:117[A]:LEU:HD11	1:A:121:ARG:HH21	1.81	0.45
1:A:168:ASN:HD22	1:A:168:ASN:C	2.21	0.44
1:A:332:ILE:HA	3:A:1381:EDO:H21	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:294:VAL:HA	1:A:332:ILE:HD13	2.00	0.43
1:A:117[A]:LEU:HD12	1:A:121:ARG:NH2	2.32	0.43
1:A:117[A]:LEU:CD1	1:A:121:ARG:CZ	2.97	0.43
1:A:270:ASN:N	1:A:270:ASN:HD22	2.11	0.43
1:A:117[A]:LEU:CD1	1:A:121:ARG:HH21	2.32	0.42
1:A:168:ASN:ND2	1:A:168:ASN:C	2.73	0.42
1:A:260:ASP:N	3:A:1371:EDO:H22	2.18	0.42
1:A:308:PRO:HB2	3:A:1368:EDO:H21	2.02	0.42
1:A:259:GLU:HB3	3:A:1371:EDO:O2	2.20	0.42
1:A:146:PRO:HG2	2:A:1360:GOL:H2	2.01	0.42
1:A:240:ASP:O	1:A:258:ARG:HA	2.19	0.41
1:A:181:PRO:HG2	1:A:184:ALA:HB2	2.01	0.41
1:A:335:GLN:NE2	1:A:353:ARG:HE	2.19	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	361/375 (96%)	353 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/318 (98%)	295 (95%)	16 (5%)	29	13

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	69	GLN
1	A	70	TRP
1	A	94	GLN
1	A	142	THR
1	A	166	ASN
1	A	168	ASN
1	A	211	ASN
1	A	265	ARG
1	A	270	ASN
1	A	280	GLU
1	A	304	ASN
1	A	305	LEU
1	A	315	GLU
1	A	329	GLU
1	A	354	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (17) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2	ASN
1	A	26	GLN
1	A	69	GLN
1	A	149	HIS
1	A	166	ASN
1	A	167	ASN
1	A	168	ASN
1	A	211	ASN
1	A	228	GLN
1	A	233	ASN
1	A	246	GLN
1	A	270	ASN
1	A	295	ASN
1	A	304	ASN
1	A	335	GLN
1	A	340	GLN
1	A	351	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	CSS	A	157	1	4,6,7	0.98	0	3,6,8	1.70	1 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSS	A	157	1	-	0/1/5/7	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	157	CSS	O-C-CA	-2.23	119.69	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

30 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	A	1358	-	5,5,5	0.27	0	5,5,5	0.38	0
2	GOL	A	1359	-	5,5,5	0.40	0	5,5,5	0.24	0
2	GOL	A	1360	-	5,5,5	0.25	0	5,5,5	0.50	0
2	GOL	A	1361	-	5,5,5	0.37	0	5,5,5	0.18	0
2	GOL	A	1362	-	5,5,5	0.32	0	5,5,5	0.30	0
3	EDO	A	1363	-	3,3,3	0.47	0	2,2,2	0.42	0
3	EDO	A	1364	-	3,3,3	0.46	0	2,2,2	0.50	0
3	EDO	A	1365	-	3,3,3	0.48	0	2,2,2	0.30	0
3	EDO	A	1366	-	3,3,3	0.51	0	2,2,2	0.50	0
3	EDO	A	1367	-	3,3,3	0.58	0	2,2,2	0.15	0
3	EDO	A	1368	-	3,3,3	0.49	0	2,2,2	0.27	0
3	EDO	A	1369	-	3,3,3	0.42	0	2,2,2	0.44	0
3	EDO	A	1370	-	3,3,3	0.47	0	2,2,2	0.56	0
3	EDO	A	1371	-	3,3,3	0.52	0	2,2,2	0.36	0
3	EDO	A	1372	-	3,3,3	0.48	0	2,2,2	0.34	0
3	EDO	A	1373	-	3,3,3	0.53	0	2,2,2	0.35	0
3	EDO	A	1374	-	3,3,3	0.47	0	2,2,2	0.47	0
3	EDO	A	1375	-	3,3,3	0.44	0	2,2,2	0.48	0
3	EDO	A	1376	-	3,3,3	0.34	0	2,2,2	0.55	0
3	EDO	A	1377	-	3,3,3	0.51	0	2,2,2	0.38	0
3	EDO	A	1378	-	3,3,3	0.47	0	2,2,2	0.35	0
3	EDO	A	1379	-	3,3,3	0.54	0	2,2,2	0.29	0
3	EDO	A	1380	-	3,3,3	0.54	0	2,2,2	0.24	0
3	EDO	A	1381	-	3,3,3	0.46	0	2,2,2	0.47	0
3	EDO	A	1382	-	3,3,3	0.47	0	2,2,2	0.51	0
3	EDO	A	1383	-	3,3,3	0.49	0	2,2,2	0.43	0
4	SO4	A	1384	-	4,4,4	1.22	0	6,6,6	0.16	0
4	SO4	A	1385	-	4,4,4	1.19	0	6,6,6	0.08	0
4	SO4	A	1386	-	4,4,4	1.20	0	6,6,6	0.08	0
4	SO4	A	1387	-	4,4,4	1.33	0	6,6,6	0.07	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	1358	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1359	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1360	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1361	-	-	0/4/4/4	0/0/0/0
2	GOL	A	1362	-	-	0/4/4/4	0/0/0/0
3	EDO	A	1363	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1364	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1365	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1366	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1367	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1368	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1369	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1370	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1371	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1372	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1373	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1374	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1375	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1376	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1377	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1378	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1379	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1380	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1381	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1382	-	-	0/1/1/1	0/0/0/0
3	EDO	A	1383	-	-	0/1/1/1	0/0/0/0
4	SO4	A	1384	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1385	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1386	-	-	0/0/0/0	0/0/0/0
4	SO4	A	1387	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

11 monomers are involved in 16 short contacts:



Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1360	GOL	2	0
2	A	1361	GOL	1	0
2	A	1362	GOL	1	0
3	A	1364	EDO	1	0
3	A	1368	EDO	2	0
3	A	1371	EDO	3	0
3	A	1372	EDO	2	0
3	A	1375	EDO	1	0
3	A	1376	EDO	1	0
3	A	1381	EDO	1	0
4	A	1387	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	356/375 (94%)	0.11	17 (4%)	34 45	21, 31, 49, 60	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	357	ALA	5.1
1	A	140	TYR	4.5
1	A	356	TRP	4.2
1	A	79	HIS	4.1
1	A	267	THR	3.6
1	A	182	ALA	3.3
1	A	350	VAL	3.3
1	A	180	PHE	3.2
1	A	302	ILE	3.1
1	A	258	ARG	2.7
1	A	275	VAL	2.6
1	A	143	PRO	2.6
1	A	269	SER	2.6
1	A	303	PHE	2.5
1	A	179	LYS	2.4
1	A	301	THR	2.3
1	A	192	LEU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
1	CSS	A	157	7/8	0.95	0.07	-	22,27,49,62	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	EDO	A	1367	4/4	0.85	0.23	13.56	32,47,49,49	0
3	EDO	A	1377	4/4	0.84	0.31	10.40	51,52,54,55	0
3	EDO	A	1365	4/4	0.86	0.41	6.68	48,51,52,58	0
3	EDO	A	1372	4/4	0.89	0.24	4.97	40,41,51,52	0
3	EDO	A	1366	4/4	0.93	0.34	4.33	43,50,51,63	0
3	EDO	A	1378	4/4	0.87	0.37	3.28	45,56,57,60	0
3	EDO	A	1373	4/4	0.90	0.15	2.82	51,51,54,54	0
2	GOL	A	1359	6/6	0.91	0.13	2.47	39,41,47,47	0
2	GOL	A	1362	6/6	0.85	0.25	2.08	40,48,56,57	0
2	GOL	A	1360	6/6	0.83	0.14	1.93	35,39,44,45	0
3	EDO	A	1381	4/4	0.88	0.28	1.20	42,53,54,67	0
3	EDO	A	1379	4/4	0.88	0.14	0.97	47,50,52,53	0
3	EDO	A	1380	4/4	0.92	0.19	0.66	41,47,62,63	0
3	EDO	A	1371	4/4	0.84	0.15	0.27	47,54,56,58	0
2	GOL	A	1358	6/6	0.95	0.09	0.09	28,32,34,36	0
3	EDO	A	1369	4/4	0.95	0.12	-0.17	39,48,54,55	0
2	GOL	A	1361	6/6	0.96	0.10	-0.32	32,36,41,42	0
3	EDO	A	1363	4/4	0.90	0.10	-0.56	48,53,58,64	0
3	EDO	A	1370	4/4	0.87	0.20	-	42,44,48,58	0
3	EDO	A	1382	4/4	0.88	0.14	-	52,60,62,65	0
4	SO4	A	1386	5/5	0.93	0.30	-	67,72,78,82	0
3	EDO	A	1383	4/4	0.84	0.33	-	57,61,63,66	0
4	SO4	A	1385	5/5	0.83	0.35	-	74,80,92,98	0
3	EDO	A	1364	4/4	0.91	0.23	-	40,45,50,55	0
3	EDO	A	1375	4/4	0.92	0.23	-	35,37,41,51	0
3	EDO	A	1374	4/4	0.93	0.19	-	52,53,55,59	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
4	SO4	A	1387	5/5	0.89	0.38	-	68,70,83,89	0
4	SO4	A	1384	5/5	0.99	0.06	-	31,35,37,37	5
3	EDO	A	1368	4/4	0.83	0.16	-	46,48,50,52	0
3	EDO	A	1376	4/4	0.92	0.10	-	43,47,54,57	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.