



Full wwPDB X-ray Structure Validation Report i

Jan 31, 2016 – 06:29 PM GMT

PDB ID : 1AZS
Title : COMPLEX OF GS-ALPHA WITH THE CATALYTIC DOMAINS OF MAMMALIAN ADENYLYL CYCLASE
Authors : Tesmer, J.J.G.; Sprang, S.R.
Deposited on : 1997-11-20
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org

A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	FKP	A	1	-	-	-	X

2 Entry composition i

There are 7 unique types of molecules in this entry. The entry contains 5881 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VC1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	190	1484	933	260	274	17	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	362	HIS	VAL	CONFLICT	UNP P30803
A	476	MET	VAL	ENGINEERED	UNP P30803

- Molecule 2 is a protein called IIC2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	189	1467	936	242	279	10	0	0	0


- Molecule 3 is a protein called GS-ALPHA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	339	2789	1769	487	520	13	0	0	0

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total Mg 1 1	0	0

- Molecule 5 is 5'-GUANOSINE-DIPHOSPHATE-MONOTHIOPHOSPHATE (three-letter code: GSP) (formula: C₁₀H₁₆N₅O₁₃P₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O	P	S		
5	C	1	32	10	5	13	3	1	0	0

- Molecule 6 is METHYLPIPERAZINOFORSKOLIN (three-letter code: FKP) (formula: C₃₀H₅₀N₂O₇).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
			Total	C	N	O				
6	A	1	39	30	2	7			0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	13	Total O 13 13	0	0
7	B	23	Total O 23 23	0	0
7	C	33	Total O 33 33	0	0

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	119.00 Å 134.05 Å 70.65 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	15.00 – 2.30 14.98 – 2.32	Depositor EDS
% Data completeness (in resolution range)	73.2 (15.00-2.30) 74.7 (14.98-2.32)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$< I/\sigma(I) >$ ¹	3.27 (at 2.32 Å)	Xtriage
Refinement program	X-PLOR 3.851	Depositor
R , R_{free}	0.219 , 0.282 0.233 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	27.5	Xtriage
Anisotropy	1.359	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 59.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.50$, $< L^2 > = 0.33$	Xtriage
Outliers	0 of 37183 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5881	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [\(i\)](#)

5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FKP, GSP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.37	0/1512	0.56	0/2038
2	B	0.41	0/1492	0.56	0/2014
3	C	0.41	0/2848	0.55	0/3853
All	All	0.40	0/5852	0.56	0/7905

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1484	0	1454	26	0
2	B	1467	0	1470	28	0
3	C	2789	0	2734	32	0
4	C	1	0	0	0	0
5	C	32	0	12	1	0
6	A	39	0	50	10	0
7	A	13	0	0	1	0
7	B	23	0	0	1	0
7	C	33	0	0	1	0
All	All	5881	0	5720	92	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (92) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:1:FKP:H201	6:A:1:FKP:H173	1.43	0.97
3:C:88:LYS:HE3	3:C:197:LEU:HD11	1.48	0.96
6:A:1:FKP:H202	6:A:1:FKP:H193	1.51	0.90
2:B:1009:VAL:HG11	2:B:1015:PRO:HB3	1.71	0.72
1:A:403:LEU:HD13	1:A:479:VAL:HG11	1.71	0.72
3:C:151:LYS:O	3:C:155:GLU:HG2	1.93	0.68
6:A:1:FKP:C17	6:A:1:FKP:H201	2.22	0.67
3:C:87:GLU:O	3:C:90:THR:HG22	1.97	0.65
1:A:411:GLU:HA	1:A:414:MET:HE2	1.79	0.65
2:B:879:HIS:HB2	2:B:1007:ALA:O	1.97	0.65
3:C:384:GLN:O	3:C:388:LEU:HG	1.97	0.65
2:B:995:LYS:HB2	2:B:1036:LEU:HD22	1.80	0.64
1:A:386:LYS:HE3	1:A:491:ARG:HD3	1.79	0.63
6:A:1:FKP:H202	6:A:1:FKP:C19	2.28	0.61
3:C:195:GLN:O	3:C:199:ARG:HG2	2.02	0.60
1:A:441:CYS:SG	6:A:1:FKP:H302	2.42	0.59
2:B:953:ILE:H	2:B:953:ILE:HD12	1.67	0.59
2:B:1062:ILE:HD12	2:B:1072:THR:HG21	1.84	0.58
2:B:1013:GLN:O	2:B:1015:PRO:HD3	2.03	0.58
2:B:1009:VAL:CG1	2:B:1015:PRO:HB3	2.35	0.57
3:C:283:ARG:HG3	3:C:283:ARG:HH11	1.70	0.56
3:C:127:ARG:NH1	7:C:427:HOH:O	2.39	0.56
2:B:1002:HIS:CE1	2:B:1048:ILE:HD12	2.42	0.54
6:A:1:FKP:O3	6:A:1:FKP:H231	2.07	0.54
1:A:401:THR:OG1	2:B:1065:LYS:HE2	2.08	0.54
1:A:413:VAL:O	1:A:417:ASN:HB2	2.07	0.53
2:B:1043:GLU:O	2:B:1046:SER:HB3	2.07	0.53
1:A:460:VAL:O	1:A:464:MET:HG2	2.08	0.53
1:A:411:GLU:HA	1:A:414:MET:CE	2.38	0.52
1:A:505:ASP:HB3	1:A:507:TRP:CZ2	2.45	0.52
2:B:1030:MET:HE3	2:B:1040:GLN:HG2	1.92	0.51
1:A:393:LEU:HD23	1:A:394:PHE:N	2.25	0.51
2:B:891:SER:HB2	7:B:31:HOH:O	2.10	0.51
3:C:326:PRO:HG3	3:C:335:THR:HG21	1.94	0.50
3:C:207:ILE:HG12	3:C:224:VAL:HG12	1.92	0.50
2:B:917:GLU:O	2:B:921:ASP:HB2	2.11	0.50
3:C:321:PRO:HG2	3:C:324:ALA:HB2	1.93	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:286:SER:HB2	3:C:386:MET:SD	2.51	0.50
2:B:1064:VAL:HG21	2:B:1070:LEU:HD21	1.94	0.49
1:A:382:ILE:O	1:A:382:ILE:HD12	2.12	0.49
1:A:507:TRP:O	1:A:508:SER:HB3	2.13	0.48
3:C:172:ILE:HD12	3:C:174:CYS:SG	2.53	0.48
3:C:38:ARG:HB3	3:C:391:TYR:HE2	1.79	0.48
2:B:969:ILE:HG22	2:B:1049:LEU:HD21	1.95	0.48
7:A:55:HOH:O	2:B:938:LYS:HB2	2.14	0.47
3:C:166:SER:HA	3:C:169:TYR:CE2	2.49	0.47
3:C:297:LEU:O	3:C:301:VAL:HG23	2.15	0.47
6:A:1:FKP:O3	6:A:1:FKP:H173	2.15	0.46
3:C:151:LYS:HE2	3:C:155:GLU:OE2	2.15	0.46
6:A:1:FKP:H242	6:A:1:FKP:H6	1.97	0.46
1:A:403:LEU:CD1	1:A:477:THR:HG21	2.45	0.46
1:A:463:GLY:O	1:A:467:ILE:HG13	2.15	0.46
1:A:452:ARG:HD3	1:A:454:ASP:HB2	1.97	0.46
2:B:1022:ASN:HA	2:B:1025:ASN:HD22	1.81	0.46
3:C:386:MET:HG2	3:C:389:ARG:NH1	2.30	0.45
3:C:298:ALA:O	3:C:302:LEU:HG	2.17	0.45
1:A:426:LEU:HD13	1:A:465:ASP:HB3	1.99	0.45
3:C:160:ARG:O	3:C:164:GLU:HG2	2.17	0.45
2:B:975:PHE:O	2:B:978:ALA:HB3	2.17	0.45
1:A:514:ALA:O	1:A:517:MET:HB2	2.17	0.44
3:C:353:GLY:O	3:C:356:ARG:HB2	2.17	0.44
3:C:254:ASN:OD1	3:C:255:MET:HG3	2.18	0.43
1:A:477:THR:HB	1:A:479:VAL:HG12	2.00	0.43
2:B:953:ILE:HD12	2:B:953:ILE:N	2.32	0.43
2:B:1025:ASN:O	2:B:1029:ARG:HG2	2.18	0.43
3:C:277:TRP:NE1	3:C:349:SER:HA	2.32	0.43
2:B:925:LEU:HD12	2:B:979:LEU:HD23	2.00	0.43
1:A:434:ARG:HA	1:A:444:CYS:HB3	2.01	0.43
3:C:293:LYS:HG2	5:C:404:GSP:C6	2.54	0.43
3:C:239:ASN:O	3:C:240:ASP:HB2	2.19	0.42
1:A:474:ARG:HG3	1:A:479:VAL:O	2.19	0.42
2:B:1044:GLU:H	2:B:1044:GLU:CD	2.21	0.42
1:A:441:CYS:SG	6:A:1:FKP:C30	3.07	0.42
2:B:1030:MET:CE	2:B:1040:GLN:HG2	2.48	0.42
1:A:504:PHE:HB3	2:B:1014:LYS:NZ	2.34	0.42
3:C:100:LYS:O	3:C:104:GLU:HG2	2.19	0.42
6:A:1:FKP:C17	6:A:1:FKP:C20	2.95	0.42
2:B:1026:VAL:O	2:B:1030:MET:HG2	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:452:ARG:HG2	1:A:454:ASP:N	2.34	0.42
1:A:470:ILE:HG21	1:A:483:MET:HG2	2.01	0.42
2:B:995:LYS:HD2	2:B:1037:ASP:OD1	2.20	0.41
2:B:888:MET:HB2	2:B:1000:ILE:HG12	2.01	0.41
3:C:121:ASN:HB2	3:C:124:ASN:ND2	2.36	0.41
3:C:187:GLN:HB2	3:C:190:TYR:HB2	2.02	0.41
1:A:403:LEU:HD22	1:A:481:VAL:HG11	2.01	0.41
1:A:456:ALA:O	1:A:460:VAL:HG23	2.20	0.41
3:C:99:LEU:CD1	3:C:179:LEU:HD23	2.50	0.41
3:C:104:GLU:HB2	3:C:135:MET:CE	2.51	0.41
3:C:128:VAL:O	3:C:132:LEU:HG	2.21	0.41
3:C:213:GLN:OE1	3:C:216:LYS:HA	2.20	0.41
3:C:382:ILE:H	3:C:382:ILE:HG13	1.66	0.41
2:B:879:HIS:CB	2:B:1008:GLY:HA3	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	188/220 (86%)	178 (95%)	9 (5%)	1 (0%)	34 41
2	B	185/212 (87%)	176 (95%)	8 (4%)	1 (0%)	34 41
3	C	335/402 (83%)	321 (96%)	12 (4%)	2 (1%)	30 36
All	All	708/834 (85%)	675 (95%)	29 (4%)	4 (1%)	30 36

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
3	C	138	PRO
2	B	1013	GLN
1	A	508	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	C	175	ALA

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	159/186 (86%)	156 (98%)	3 (2%)	65 81
2	B	162/184 (88%)	158 (98%)	4 (2%)	55 73
3	C	306/357 (86%)	301 (98%)	5 (2%)	70 84
All	All	627/727 (86%)	615 (98%)	12 (2%)	65 81

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	415	THR
1	A	530	LYS
1	A	550	ARG
2	B	921	ASP
2	B	938	LYS
2	B	1037	ASP
2	B	1059	ARG
3	C	141	ASP
3	C	203	LEU
3	C	229	ASP
3	C	357	HIS
3	C	378	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	385	GLN
1	A	537	ASN
2	B	879	HIS
2	B	1001	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	1002	HIS
2	B	1025	ASN
2	B	1076	ASN
3	C	97	ASN
3	C	220	HIS
3	C	362	HIS
3	C	371	ASN
3	C	377	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 3 ligands modelled in this entry, 1 is monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	FKP	A	1	-	37,42,42	1.29	4 (10%)	49,68,68	1.29	5 (10%)
5	GSP	C	404	4	25,34,34	1.20	2 (8%)	31,54,54	2.29	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	FKP	A	1	-	-	0/14/97/97	0/4/4/4
5	GSP	C	404	4	-	0/15/38/38	0/3/3/3

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	404	GSP	C8-N7	-2.96	1.28	1.34
6	A	1	FKP	C17-C8	2.08	1.55	1.51
6	A	1	FKP	C2-C1	2.35	1.55	1.52
5	C	404	GSP	C6-N1	2.45	1.37	1.33
6	A	1	FKP	C12-C11	2.93	1.55	1.50
6	A	1	FKP	C10-C5	3.02	1.62	1.56

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	404	GSP	C5-C6-N1	-9.04	111.23	123.59
6	A	1	FKP	C19-C4-C3	-2.42	103.90	109.00
5	C	404	GSP	PA-O3A-PB	-2.41	125.95	132.73
6	A	1	FKP	C30-N2-C27	-2.08	107.38	110.63
6	A	1	FKP	O1-C13-C12	2.09	113.70	111.23
6	A	1	FKP	C19-C4-C5	2.51	120.86	111.85
6	A	1	FKP	C9-C10-C1	3.00	113.70	110.90
5	C	404	GSP	C6-N1-C2	6.30	124.69	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1	FKP	10	0
5	C	404	GSP	1	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	190/220 (86%)	0.36	14 (7%) 17 25	29, 63, 93, 97	0
2	B	189/212 (89%)	-0.10	8 (4%) 40 49	23, 42, 88, 96	0
3	C	339/402 (84%)	-0.05	14 (4%) 41 50	25, 46, 83, 97	0
All	All	718/834 (86%)	0.04	36 (5%) 32 41	23, 48, 88, 97	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	C	37	TYR	8.3
2	B	953	ILE	8.1
2	B	952	ALA	7.1
1	A	376	ASP	6.9
3	C	391	TYR	6.3
3	C	393	LEU	5.6
1	A	476	MET	5.4
1	A	480	ASN	5.1
2	B	964	ARG	5.1
3	C	392	GLU	5.0
3	C	38	ARG	4.6
3	C	66	ASN	4.5
1	A	475	GLU	4.3
2	B	1077	THR	4.1
3	C	354	ASP	3.8
3	C	36	VAL	3.5
1	A	377	MET	3.4
3	C	86	GLY	3.2
1	A	472	LEU	3.2
3	C	87	GLU	3.1
1	A	478	GLY	3.0
1	A	474	ARG	2.8
3	C	139	ASP	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	1065	LYS	2.6
1	A	482	ASN	2.5
1	A	398	GLU	2.4
1	A	468	GLU	2.3
1	A	471	SER	2.3
3	C	322	GLU	2.3
1	A	479	VAL	2.2
2	B	966	TYR	2.2
3	C	381	ASP	2.1
1	A	406	GLN	2.1
2	B	965	GLN	2.1
2	B	951	SER	2.1
3	C	157	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
6	FKP	A	1	39/39	0.89	0.18	2.37	29,42,86,88	0
5	GSP	C	404	32/32	0.99	0.10	-0.67	23,32,41,48	0
4	MG	C	403	1/1	0.95	0.05	-1.78	28,28,28,28	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.