



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 07:32 AM GMT

PDB ID : 3B4V
Title : X-Ray structure of Activin in complex with FSTL3
Authors : Thompson, T.B.
Deposited on : 2007-10-24
Resolution : 2.48 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.48 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4309 (2.50-2.46)
Clashscore	102246	5050 (2.50-2.46)
Ramachandran outliers	100387	4961 (2.50-2.46)
Sidechain outliers	100360	4963 (2.50-2.46)
RSRZ outliers	91569	4319 (2.50-2.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	116	<div> <div>8%</div> <div>85%</div> <div>9%</div> <div>6%</div> </div>
1	B	116	<div> <div>6%</div> <div>87%</div> <div>11%</div> <div>•</div> </div>
1	E	116	<div> <div>7%</div> <div>76%</div> <div>16%</div> <div>•</div> <div>8%</div> </div>
1	F	116	<div> <div>3%</div> <div>85%</div> <div>12%</div> <div>••</div> </div>
2	C	237	<div> <div>8%</div> <div>79%</div> <div>8%</div> <div>13%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	237	
2	G	237	
2	H	237	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	C	240	-	-	-	X

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9726 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inhibin beta A chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	109	Total	C	N	O	S	0	0	0
			855	540	147	155	13			
1	B	114	Total	C	N	O	S	0	0	0
			886	558	153	162	13			
1	E	107	Total	C	N	O	S	0	0	0
			845	534	145	153	13			
1	F	114	Total	C	N	O	S	0	0	0
			880	553	152	162	13			

- Molecule 2 is a protein called follistatin-like 3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	206	Total	C	N	O	S	0	0	0
			1494	896	287	282	29			
2	D	213	Total	C	N	O	S	0	0	0
			1533	919	296	289	29			
2	G	206	Total	C	N	O	S	0	0	0
			1490	894	286	281	29			
2	H	211	Total	C	N	O	S	0	0	0
			1521	913	293	286	29			

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	G	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is SULFATE ION (three-letter code: SO₄) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is SUGAR (N-ACETYL-D-GLUCOSAMINE) (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	G	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	19	Total	O	0	0
			19	19		
6	B	19	Total	O	0	0
			19	19		
6	C	16	Total	O	0	0
			16	16		
6	D	36	Total	O	0	0
			36	36		
6	E	15	Total	O	0	0
			15	15		
6	F	16	Total	O	0	0
			16	16		
6	G	22	Total	O	0	0
			22	22		
6	H	34	Total	O	0	0
			34	34		


● Molecule 2: follistatin-like 3

Chain D: 





● Molecule 2: follistatin-like 3

Chain G: 



● Molecule 2: follistatin-like 3

Chain H: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	63.61Å 71.38Å 100.20Å 98.55° 90.64° 90.11°	Depositor
Resolution (Å)	31.78 – 2.48 31.01 – 2.48	Depositor EDS
% Data completeness (in resolution range)	98.6 (31.78-2.48) 91.3 (31.01-2.48)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.68 (at 2.48Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.227 , 0.279 0.244 , 0.270	Depositor DCC
R_{free} test set	3141 reflections (5.41%)	DCC
Wilson B-factor (Å ²)	52.9	Xtriage
Anisotropy	0.023	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.27 , 2.2	EDS
Estimated twinning fraction	0.407 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 61223 reflections	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	9726	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, EDO, NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.51	0/876	0.58	0/1178
1	B	0.45	0/907	0.52	0/1218
1	E	0.49	0/866	0.57	0/1164
1	F	0.83	3/901 (0.3%)	0.54	0/1211
2	C	0.42	0/1521	0.51	0/2060
2	D	0.53	2/1563 (0.1%)	0.55	0/2121
2	G	0.56	2/1517 (0.1%)	0.56	1/2055 (0.0%)
2	H	0.72	3/1550 (0.2%)	0.54	0/2103
All	All	0.58	10/9701 (0.1%)	0.55	1/13110 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	6	GLY	N-CA	17.18	1.71	1.46
2	H	6	PRO	CA-CB	14.27	1.82	1.53
2	H	6	PRO	N-CA	14.20	1.71	1.47
2	G	74	ASP	CG-OD2	11.19	1.51	1.25
2	G	74	ASP	CG-OD1	10.22	1.48	1.25
1	F	6	GLY	C-O	10.10	1.39	1.23
2	H	6	PRO	CG-CD	8.52	1.78	1.50
2	D	33	ALA	C-N	8.41	1.53	1.34
2	D	33	ALA	C-O	6.07	1.34	1.23
1	F	6	GLY	CA-C	5.33	1.60	1.51

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	G	74	ASP	CB-CG-OD1	-8.71	110.46	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	33	ALA	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	855	0	801	9	0
1	B	886	0	843	10	0
1	E	845	0	797	15	0
1	F	880	0	825	17	0
2	C	1494	0	1407	8	0
2	D	1533	0	1438	11	0
2	G	1490	0	1401	13	0
2	H	1521	0	1425	15	0
3	A	4	0	6	0	0
3	C	4	0	6	0	0
3	G	4	0	6	0	0
4	C	5	0	0	0	0
5	C	14	0	13	1	0
5	G	14	0	13	0	0
6	A	19	0	0	0	0
6	B	19	0	0	0	0
6	C	16	0	0	0	0
6	D	36	0	0	0	0
6	E	15	0	0	0	0
6	F	16	0	0	0	0
6	G	22	0	0	2	0
6	H	34	0	0	0	0
All	All	9726	0	8981	77	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (77) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:6:PRO:CA	2:H:6:PRO:CB	1.82	1.57
2:H:6:PRO:CD	2:H:6:PRO:CG	1.78	1.53
2:H:6:PRO:CA	2:H:6:PRO:N	1.71	1.53
1:F:6:GLY:CA	1:F:6:GLY:N	1.71	1.50
2:H:6:PRO:CD	2:H:6:PRO:CA	2.63	0.76
2:H:6:PRO:CA	2:H:6:PRO:CG	2.69	0.70
2:H:6:PRO:CD	2:H:6:PRO:CB	2.71	0.66
1:F:45:PRO:HG2	1:F:56:LEU:HD21	1.84	0.60
2:H:6:PRO:C	2:H:6:PRO:CB	2.68	0.60
2:H:6:PRO:N	2:H:6:PRO:CG	2.62	0.58
2:C:159:ASP:OD1	2:C:161:THR:HB	2.03	0.57
2:D:106:LEU:O	2:D:118:ASP:HA	2.04	0.57
1:B:86:LEU:HD23	1:B:110:VAL:HA	1.87	0.56
1:E:110:VAL:HG23	1:F:59:HIS:NE2	2.21	0.56
1:E:45:PRO:HG2	1:E:56:LEU:HD21	1.88	0.56
1:F:6:GLY:C	1:F:6:GLY:N	2.55	0.55
1:A:18:VAL:HG21	1:B:62:VAL:CG1	2.37	0.55
2:C:117:ARG:HB2	2:C:121:GLU:HG3	1.89	0.55
1:A:82:VAL:HG21	1:B:82:VAL:HG11	1.90	0.54
1:A:110:VAL:HG23	1:B:59:HIS:NE2	2.23	0.53
1:F:55:SER:HB3	2:G:24:LEU:HG	1.90	0.53
2:C:189:ASN:OD1	2:C:212:ARG:NH1	2.42	0.52
2:H:11:VAL:HG21	2:H:55:ILE:HG12	1.91	0.52
2:C:98:ASP:OD1	2:C:100:SER:HB3	2.11	0.51
1:F:65:HIS:HD2	6:G:248:HOH:O	1.93	0.49
2:G:159:ASP:OD1	2:G:161:THR:HB	2.12	0.49
1:A:62:VAL:CG1	1:B:18:VAL:HG21	2.42	0.49
2:C:71:ASP:OD2	2:C:71:ASP:N	2.44	0.49
2:H:43:THR:HG23	2:H:69:CYS:SG	2.53	0.48
2:G:31:THR:HG23	2:G:34:GLU:H	1.79	0.48
1:F:30:ILE:CG2	2:G:167:VAL:HG11	2.43	0.48
1:E:38:ASN:HB2	1:E:110:VAL:HB	1.96	0.48
2:D:117:ARG:NH2	2:D:121:GLU:OE1	2.48	0.47
1:E:59:HIS:NE2	1:F:110:VAL:HG23	2.30	0.47
2:H:119:GLU:OE2	2:H:123:ARG:NH1	2.48	0.47
2:D:11:VAL:HG21	2:D:55:ILE:HG12	1.97	0.47
1:F:97:GLY:O	2:G:210:GLY:HA2	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:55:SER:CB	2:G:24:LEU:HG	2.46	0.46
1:E:18:VAL:HG13	1:E:23:ILE:HD11	1.98	0.46
1:F:100:ILE:HG22	2:G:159:ASP:HA	1.98	0.46
2:D:156:CYS:HA	2:D:165:HIS:O	2.15	0.46
1:E:18:VAL:HG22	1:F:66:TYR:CE1	2.50	0.45
2:C:126:ARG:HD3	2:C:133:LEU:O	2.15	0.45
2:D:43:THR:HG23	2:D:69:CYS:SG	2.56	0.45
1:E:82:VAL:HG11	1:F:82:VAL:HG11	1.97	0.45
1:E:87:ARG:HB2	1:E:88:PRO:HD2	1.97	0.45
1:F:31:ALA:HB2	2:G:167:VAL:HG22	1.98	0.45
1:B:9:ASN:OD1	1:B:9:ASN:N	2.44	0.44
1:E:18:VAL:HG21	1:F:62:VAL:CG1	2.48	0.44
1:A:55:SER:HB3	2:D:25:VAL:O	2.18	0.43
1:F:65:HIS:CD2	6:G:248:HOH:O	2.71	0.43
2:H:106:LEU:O	2:H:118:ASP:HA	2.18	0.43
2:C:187:ASN:O	5:C:239:NAG:H82	2.18	0.43
2:G:43:THR:HG23	2:G:69:CYS:SG	2.59	0.43
1:B:25:TRP:CG	2:D:57:LEU:HD22	2.53	0.43
2:G:176:VAL:HA	2:G:177:PRO:HD3	1.86	0.43
2:H:12:CYS:HB2	2:H:30:VAL:HG21	2.00	0.43
1:E:2:LEU:HD23	1:E:10:ILE:HD13	1.99	0.43
2:G:102:LEU:HD13	2:G:119:GLU:HG2	2.01	0.43
1:A:72:SER:HA	1:A:73:PRO:HA	1.91	0.42
2:C:73:CYS:O	2:C:76:VAL:HG22	2.19	0.42
1:A:59:HIS:NE2	1:B:110:VAL:HG23	2.35	0.42
2:D:85:ARG:NE	2:D:94:GLU:OE2	2.52	0.42
2:D:50:HIS:HB3	2:D:54:LYS:HB3	2.02	0.42
1:E:103:LYS:HB3	1:E:103:LYS:HE2	1.70	0.42
2:D:15:GLN:OE1	2:D:20:ALA:HB1	2.19	0.41
2:H:7:ALA:HA	2:H:8:PRO:HD2	1.86	0.41
1:A:18:VAL:HG22	1:B:66:TYR:CE1	2.56	0.41
1:E:2:LEU:HD12	1:E:2:LEU:HA	1.94	0.41
2:G:156:CYS:HA	2:G:165:HIS:O	2.20	0.41
1:E:25:TRP:HB3	1:E:29:ILE:HD12	2.02	0.41
2:H:150:CYS:HA	2:H:151:PRO:HD2	1.90	0.41
1:B:45:PRO:HG2	1:B:56:LEU:HD21	2.01	0.41
1:A:58:PHE:HA	2:D:60:PHE:CE2	2.55	0.41
1:E:41:GLU:HG3	1:F:77:LEU:HD11	2.02	0.40
2:G:14:LEU:HB2	2:G:26:LEU:HD13	2.02	0.40
1:E:72:SER:HA	1:E:73:PRO:HA	1.86	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	103/116 (89%)	99 (96%)	3 (3%)	1 (1%)	19	32
1	B	108/116 (93%)	102 (94%)	6 (6%)	0	100	100
1	E	101/116 (87%)	97 (96%)	3 (3%)	1 (1%)	19	32
1	F	109/116 (94%)	103 (94%)	6 (6%)	0	100	100
2	C	202/237 (85%)	195 (96%)	7 (4%)	0	100	100
2	D	211/237 (89%)	201 (95%)	10 (5%)	0	100	100
2	G	202/237 (85%)	198 (98%)	4 (2%)	0	100	100
2	H	207/237 (87%)	200 (97%)	7 (3%)	0	100	100
All	All	1243/1412 (88%)	1195 (96%)	46 (4%)	2 (0%)	52	73

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	38	ASN
1	E	38	ASN

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	95/102 (93%)	95 (100%)	0	100	100
1	B	99/102 (97%)	97 (98%)	2 (2%)	63	85
1	E	95/102 (93%)	94 (99%)	1 (1%)	80	93

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	97/102 (95%)	97 (100%)	0	100	100
2	C	166/191 (87%)	162 (98%)	4 (2%)	57	81
2	D	168/191 (88%)	162 (96%)	6 (4%)	42	68
2	G	165/191 (86%)	160 (97%)	5 (3%)	48	74
2	H	167/191 (87%)	158 (95%)	9 (5%)	27	47
All	All	1052/1172 (90%)	1025 (97%)	27 (3%)	54	79

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	21	LYS
1	B	54	SER
2	C	21	THR
2	C	66	CYS
2	C	67	LEU
2	C	152	ARG
2	D	14	LEU
2	D	66	CYS
2	D	67	LEU
2	D	161	THR
2	D	167	VAL
2	D	189	ASN
1	E	85	LYS
2	G	66	CYS
2	G	67	LEU
2	G	168	VAL
2	G	195	SER
2	G	216	SER
2	H	14	LEU
2	H	21	THR
2	H	24	LEU
2	H	26	LEU
2	H	66	CYS
2	H	67	LEU
2	H	76	VAL
2	H	115	THR
2	H	161	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	F	65	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	117	-	3,3,3	0.51	0	2,2,2	0.29	0
4	SO4	C	238	-	4,4,4	0.17	0	6,6,6	0.14	0
5	NAG	C	239	2	14,14,15	0.81	1 (7%)	15,19,21	1.25	2 (13%)
3	EDO	C	240	-	3,3,3	0.52	0	2,2,2	0.27	0
5	NAG	G	238	2	14,14,15	0.53	0	15,19,21	1.64	2 (13%)
3	EDO	G	239	-	3,3,3	0.52	0	2,2,2	0.25	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	117	-	-	0/1/1/1	0/0/0/0
4	SO4	C	238	-	-	0/0/0/0	0/0/0/0
5	NAG	C	239	2	-	0/6/23/26	0/1/1/1
3	EDO	C	240	-	-	0/1/1/1	0/0/0/0
5	NAG	G	238	2	-	0/6/23/26	0/1/1/1
3	EDO	G	239	-	-	0/1/1/1	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	239	NAG	C8-C7	2.06	1.54	1.50

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	C	239	NAG	C3-C4-C5	-2.13	106.48	110.20
5	G	238	NAG	O5-C5-C6	2.39	112.53	107.35
5	C	239	NAG	O5-C5-C6	2.65	113.08	107.35
5	G	238	NAG	C2-N2-C7	4.48	128.79	123.04

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	C	239	NAG	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	109/116 (93%)	0.54	9 (8%) 14 15	17, 29, 65, 69	0
1	B	114/116 (98%)	0.44	7 (6%) 25 27	14, 32, 54, 59	0
1	E	107/116 (92%)	0.39	8 (7%) 17 18	18, 28, 66, 71	0
1	F	114/116 (98%)	0.27	3 (2%) 59 63	16, 32, 52, 56	0
2	C	206/237 (86%)	0.57	19 (9%) 11 12	23, 46, 73, 83	0
2	D	213/237 (89%)	0.26	7 (3%) 50 54	16, 40, 65, 70	0
2	G	206/237 (86%)	0.49	15 (7%) 18 19	26, 45, 70, 80	0
2	H	211/237 (89%)	0.32	11 (5%) 31 35	20, 38, 66, 72	0
All	All	1280/1412 (90%)	0.41	79 (6%) 24 26	14, 37, 68, 83	0

All (79) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	48	ILE	5.6
1	B	49	ALA	5.4
1	B	7	LYS	5.2
1	B	8	VAL	5.0
1	A	1	GLY	4.9
2	C	218	ALA	4.6
2	G	92	ARG	4.6
2	C	92	ARG	4.5
1	F	1	GLY	4.5
2	C	76	VAL	4.3
1	A	47	HIS	4.3
2	H	8	PRO	4.2
2	H	48	LEU	4.2
2	G	117	ARG	4.1
1	B	48	ILE	4.0
2	C	13	TRP	4.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	E	1	GLY	3.8
1	A	2	LEU	3.8
2	H	7	ALA	3.7
1	F	49	ALA	3.6
2	G	86	MET	3.6
2	C	9	GLY	3.5
2	C	37	ALA	3.5
1	E	73	PRO	3.4
1	B	2	LEU	3.4
1	E	76	ASN	3.3
1	F	48	ILE	3.3
2	G	26	LEU	3.2
2	G	100	SER	3.2
2	D	8	PRO	3.2
2	G	85	ARG	3.2
2	G	72	SER	3.2
2	C	20	ALA	3.1
2	C	212	ARG	3.1
2	C	65	HIS	3.0
1	E	2	LEU	2.9
2	D	33	ALA	2.9
2	C	66	CYS	2.9
1	E	47	HIS	2.8
2	C	67	LEU	2.8
2	H	67	LEU	2.8
2	C	33	ALA	2.8
2	G	20	ALA	2.7
1	A	75	ALA	2.7
1	A	78	LYS	2.7
2	G	19	GLU	2.7
2	H	87	LEU	2.6
2	C	91	PRO	2.6
2	C	77	GLU	2.6
2	C	18	GLN	2.6
2	G	116	TYR	2.6
2	G	49	THR	2.6
2	H	218	ALA	2.5
2	C	100	SER	2.5
2	D	89	GLY	2.5
1	A	74	PHE	2.5
2	D	75	GLY	2.4
1	E	5	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	H	92	ARG	2.4
2	G	18	GLN	2.4
2	H	6	PRO	2.3
1	E	48	ILE	2.3
2	D	9	GLY	2.3
2	H	65	HIS	2.3
2	G	65	HIS	2.3
2	D	91	PRO	2.3
2	C	193	ILE	2.2
1	A	46	SER	2.2
1	B	1	GLY	2.2
1	B	6	GLY	2.2
2	G	91	PRO	2.2
1	E	78	LYS	2.1
2	H	75	GLY	2.1
2	D	92	ARG	2.1
2	H	24	LEU	2.1
2	C	22	CYS	2.1
2	G	28	THR	2.1
2	C	180	PRO	2.0
1	A	69	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	EDO	C	240	4/4	0.86	0.25	6.37	62,65,66,67	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	SO4	C	238	5/5	0.93	0.17	-0.08	78,79,80,80	0
5	NAG	C	239	14/15	0.78	0.16	-0.42	73,79,81,81	0
5	NAG	G	238	14/15	0.69	0.17	-	76,82,83,83	0
3	EDO	G	239	4/4	0.89	0.29	-	72,73,73,74	0
3	EDO	A	117	4/4	0.88	0.28	-	60,60,61,62	0

6.5 Other polymers [i](#)

There are no such residues in this entry.