



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 12:38 AM GMT

PDB ID : 2B8W
Title : Crystal-structure of the N-terminal Large GTPase Domain of human Guanylate Binding protein 1 (hGBP1) in complex with GMP/AlF₄
Authors : Ghosh, A.; Praefcke, G.J.K.; Renault, L.; Wittinghofer, A.; Herrmann, C.
Deposited on : 2005-10-10
Resolution : 2.22 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.22 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4405 (2.24-2.20)
Clashscore	102246	5146 (2.24-2.20)
Ramachandran outliers	100387	5065 (2.24-2.20)
Sidechain outliers	100360	5066 (2.24-2.20)
RSRZ outliers	91569	4414 (2.24-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	328	 9% 71% 14% • 12%
1	B	328	 9% 69% 15% 5% • 12%

2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4821 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Interferon-induced guanylate-binding protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	288	2290	1470	385	421	14	0	2	0
1	B	290	2304	1479	387	424	14	0	2	0


There are 22 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	CLONING ARTIFACT	UNP P32455
A	-9	HIS	-	EXPRESSION TAG	UNP P32455
A	-8	HIS	-	EXPRESSION TAG	UNP P32455
A	-7	HIS	-	EXPRESSION TAG	UNP P32455
A	-6	HIS	-	EXPRESSION TAG	UNP P32455
A	-5	HIS	-	EXPRESSION TAG	UNP P32455
A	-4	HIS	-	EXPRESSION TAG	UNP P32455
A	-3	MET	-	CLONING ARTIFACT	UNP P32455
A	-2	ARG	-	CLONING ARTIFACT	UNP P32455
A	-1	GLY	-	CLONING ARTIFACT	UNP P32455
A	0	SER	-	CLONING ARTIFACT	UNP P32455
B	-10	MET	-	CLONING ARTIFACT	UNP P32455
B	-9	HIS	-	EXPRESSION TAG	UNP P32455
B	-8	HIS	-	EXPRESSION TAG	UNP P32455
B	-7	HIS	-	EXPRESSION TAG	UNP P32455
B	-6	HIS	-	EXPRESSION TAG	UNP P32455
B	-5	HIS	-	EXPRESSION TAG	UNP P32455
B	-4	HIS	-	EXPRESSION TAG	UNP P32455
B	-3	MET	-	CLONING ARTIFACT	UNP P32455
B	-2	ARG	-	CLONING ARTIFACT	UNP P32455
B	-1	GLY	-	CLONING ARTIFACT	UNP P32455
B	0	SER	-	CLONING ARTIFACT	UNP P32455

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Mg 1 1	0	0
2	A	1	Total Mg 1 1	0	0

- Molecule 3 is TETRAFLUOROALUMINATE ION (three-letter code: ALF) (formula: AlF_4).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Al F 5 1 4	0	0
3	B	1	Total Al F 5 1 4	0	0

- Molecule 4 is GUANOSINE-5'-MONOPHOSPHATE (three-letter code: 5GP) (formula: $\text{C}_{10}\text{H}_{14}\text{N}_5\text{O}_8\text{P}$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	0
			24	10	5	8	1		
4	B	1	Total	C	N	O	P	0	0
			24	10	5	8	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	82	Total	O	0	0
			82	82		
5	B	85	Total	O	0	0
			85	85		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	149.62Å 101.93Å 54.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.75 – 2.22 28.78 – 2.22	Depositor EDS
% Data completeness (in resolution range)	99.6 (29.75-2.22) 99.6 (28.78-2.22)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.70 (at 2.22Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.190 , 0.233 0.201 , 0.239	Depositor DCC
R_{free} test set	2758 reflections (7.11%)	DCC
Wilson B-factor (Å ²)	40.6	Xtrriage
Anisotropy	0.340	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 38.2	EDS
Estimated twinning fraction	No twinning to report.	Xtrriage
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.32$	Xtrriage
Outliers	0 of 41572 reflections	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4821	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 38.64 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 3.5758e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ALF, MG, 5GP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	2/2350 (0.1%)	0.43	0/3176
1	B	0.35	0/2364	0.43	0/3195
All	All	0.37	2/4714 (0.0%)	0.43	0/6371

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
1	B	0	6
All	All	0	10

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	210	LYS	CE-NZ	6.48	1.65	1.49
1	A	210	LYS	CG-CD	6.05	1.73	1.52

There are no bond angle outliers.

There are no chirality outliers.

All (10) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	153	ARG	Sidechain
1	A	195	PRO	Mainchain
1	A	210	LYS	Mainchain
1	A	240	ARG	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	155	LYS	Peptide
1	B	212	THR	Mainchain
1	B	240	ARG	Sidechain
1	B	308	ASP	Peptide
1	B	47	TYR	Sidechain
1	B	8	THR	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2290	0	2318	46	0
1	B	2304	0	2334	57	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	5	0	0	0	0
3	B	5	0	0	0	0
4	A	24	0	12	0	0
4	B	24	0	12	0	0
5	A	82	0	0	2	0
5	B	85	0	0	4	0
All	All	4821	0	4676	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:THR:HG23	1:A:129:ASN:ND2	1.68	1.08
1:A:149:THR:CG2	1:A:153:ARG:HH22	1.70	1.05
1:A:149:THR:CG2	1:A:153:ARG:NH2	2.20	1.04
1:A:49:THR:CG2	1:A:129:ASN:HD22	1.76	0.98
1:A:209:LYS:H	1:A:220:ASN:HD21	1.09	0.97
1:B:70:THR:HG22	1:B:72:GLN:H	1.32	0.92
1:B:62:LYS:O	1:B:63:LYS:HB3	1.67	0.91

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:209:LYS:H	1:B:220:ASN:HD21	1.20	0.89
1:A:149:THR:HG22	1:A:153:ARG:NH2	1.86	0.89
1:A:245:ARG:HD2	1:B:239:ASP:OD2	1.74	0.86
1:B:212:THR:O	1:B:213:SER:O	1.93	0.86
1:A:49:THR:HG23	1:A:129:ASN:HD22	1.30	0.84
1:A:225:CYS:HB3	5:A:677:HOH:O	1.77	0.84
1:B:128:TYR:CE1	5:B:641:HOH:O	2.32	0.82
1:A:16:ASN:HD21	1:A:111:ASN:HD21	1.31	0.79
1:B:70:THR:CG2	1:B:72:GLN:H	1.96	0.77
1:B:156:SER:HA	1:B:157:SER:HB3	1.68	0.75
1:B:210:LYS:HE2	1:B:210:LYS:H	1.51	0.75
1:A:172:VAL:O	1:A:172:VAL:HG22	1.85	0.74
1:B:62:LYS:O	1:B:63:LYS:CB	2.36	0.73
1:A:49:THR:CG2	1:A:129:ASN:ND2	2.40	0.72
1:B:149:THR:HG23	1:B:153:ARG:NH1	2.05	0.72
1:B:221:LEU:HD23	1:B:222:PRO:HD3	1.73	0.71
1:A:149:THR:HG23	1:A:153:ARG:NH2	2.07	0.70
1:A:149:THR:HG22	1:A:153:ARG:HH22	1.48	0.69
1:B:209:LYS:N	1:B:220:ASN:HD21	1.90	0.69
1:B:149:THR:CG2	1:B:153:ARG:HH12	2.04	0.68
1:A:245:ARG:CD	1:B:239:ASP:OD2	2.41	0.68
1:B:16:ASN:HD21	1:B:111:ASN:HD21	1.42	0.67
1:A:149:THR:HG23	1:A:153:ARG:CZ	2.27	0.65
1:A:16:ASN:ND2	1:A:111:ASN:HD21	1.94	0.65
1:A:259:ASP:CG	1:B:245:ARG:HH12	2.01	0.64
1:A:199:ASP:O	1:A:203:THR:CG2	2.46	0.64
1:B:211:GLY:HA3	1:B:216:ASP:OD1	1.98	0.63
1:A:199:ASP:O	1:A:203:THR:HG22	1.98	0.63
1:A:221:LEU:N	1:A:222:PRO:HD2	2.14	0.63
1:B:214:GLN:O	1:B:217:GLU:HB2	1.99	0.62
1:B:44:VAL:HG13	1:B:126:PHE:CE1	2.35	0.62
1:B:146:THR:HG22	5:B:664:HOH:O	1.99	0.62
1:A:149:THR:HG21	1:A:153:ARG:HH22	1.61	0.61
1:A:155:LYS:O	1:A:156:SER:CB	2.48	0.61
1:B:149:THR:CG2	1:B:153:ARG:NH1	2.64	0.60
1:B:70:THR:HG21	5:B:650:HOH:O	2.01	0.60
1:A:155:LYS:O	1:A:156:SER:HB3	2.03	0.59
1:B:221:LEU:N	1:B:222:PRO:HD2	2.18	0.58
1:A:49:THR:HG23	1:A:129:ASN:HD21	1.63	0.58
1:B:146:THR:CG2	5:B:664:HOH:O	2.52	0.58
1:B:31:LEU:O	1:B:290:GLY:HA3	2.04	0.57

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:209:LYS:N	1:A:220:ASN:HD21	1.91	0.56
1:A:172:VAL:O	1:A:172:VAL:CG2	2.52	0.56
1:A:11:MET:SD	1:A:30:ILE:HD12	2.46	0.55
1:B:227:ARG:HA	1:B:233:LYS:HE2	1.88	0.55
1:B:70:THR:HG22	1:B:72:GLN:N	2.11	0.54
1:A:142:LEU:O	1:A:146:THR:HG23	2.07	0.54
1:A:49:THR:HG22	1:A:51:LYS:HG3	1.90	0.54
1:A:221:LEU:N	1:A:222:PRO:CD	2.71	0.54
1:B:213:SER:OG	1:B:216:ASP:CG	2.47	0.53
1:A:221:LEU:HB3	1:A:222:PRO:HD3	1.91	0.53
1:B:199:ASP:O	1:B:203:THR:HG23	2.08	0.53
1:B:213:SER:O	1:B:216:ASP:HB2	2.08	0.53
1:A:155:LYS:HD3	1:A:155:LYS:N	2.23	0.52
1:B:67:LEU:CD1	1:B:251:GLU:HG3	2.40	0.52
1:B:212:THR:O	1:B:213:SER:C	2.48	0.52
1:A:49:THR:O	1:A:49:THR:HG23	2.10	0.52
1:B:16:ASN:ND2	1:B:111:ASN:HD21	2.08	0.52
1:A:16:ASN:HD21	1:A:111:ASN:ND2	2.04	0.51
1:A:31:LEU:O	1:A:290:GLY:HA3	2.10	0.51
1:B:156:SER:HA	1:B:157:SER:CB	2.35	0.51
1:B:209:LYS:H	1:B:220:ASN:ND2	2.00	0.50
1:A:28:LEU:HD22	1:A:294:GLU:HG3	1.94	0.49
1:A:44:VAL:CG1	1:A:45:GLY:N	2.76	0.49
1:B:70:THR:HG23	1:B:71:VAL:N	2.29	0.48
1:A:106:LYS:NZ	1:A:111:ASN:HD22	2.12	0.47
1:A:259:ASP:OD1	1:B:245:ARG:NH1	2.45	0.47
1:B:174:PHE:HB2	1:B:281:THR:HG22	1.97	0.47
1:B:106:LYS:HZ2	1:B:111:ASN:HD22	1.62	0.47
1:B:28:LEU:HD22	1:B:294:GLU:HG3	1.97	0.46
1:B:194:GLN:HE21	1:B:194:GLN:HB2	1.48	0.46
1:A:44:VAL:HG12	1:A:45:GLY:N	2.29	0.46
1:B:70:THR:CG2	1:B:71:VAL:N	2.78	0.46
1:B:210:LYS:CE	1:B:210:LYS:H	2.26	0.46
1:B:106:LYS:NZ	1:B:111:ASN:HD22	2.12	0.46
1:A:286:ILE:HG13	5:A:666:HOH:O	2.17	0.45
1:B:63:LYS:HG3	1:B:63:LYS:O	2.16	0.44
1:B:215:LYS:C	1:B:217:GLU:N	2.71	0.44
1:B:174:PHE:CB	1:B:281:THR:HG22	2.47	0.43
1:B:156:SER:CA	1:B:157:SER:HB3	2.44	0.43
1:A:200:GLU:HA	1:A:203:THR:HG23	1.99	0.43
1:B:114:TRP:HZ3	1:B:148:LEU:HD11	1.82	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:220:ASN:HA	1:B:220:ASN:HD22	1.65	0.42
1:B:259:ASP:HA	1:B:260:PRO:HD3	1.78	0.42
1:A:49:THR:HG21	1:A:129:ASN:HD22	1.76	0.41
1:B:87:LYS:HB3	1:B:87:LYS:HE2	1.92	0.41
1:A:259:ASP:HA	1:A:260:PRO:HD3	1.88	0.41
1:A:116:PHE:CD2	1:A:145:VAL:HG11	2.56	0.41
1:B:221:LEU:H	1:B:222:PRO:HD2	1.83	0.41
1:B:20:ARG:HA	1:B:20:ARG:HD3	1.67	0.41
1:B:44:VAL:CG1	1:B:126:PHE:CE1	3.04	0.40
1:B:114:TRP:CZ3	1:B:148:LEU:HD11	2.57	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	286/328 (87%)	276 (96%)	6 (2%)	4 (1%)	14	10
1	B	288/328 (88%)	275 (96%)	6 (2%)	7 (2%)	7	4
All	All	574/656 (88%)	551 (96%)	12 (2%)	11 (2%)	10	6

All (11) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	63	LYS
1	B	213	SER
1	B	244	ARG
1	A	155	LYS
1	A	210	LYS
1	A	244	ARG
1	B	216	ASP
1	B	306	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	214	GLN
1	A	307	GLY
1	B	307	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/293 (88%)	239 (92%)	20 (8%)	16	15
1	B	261/293 (89%)	238 (91%)	23 (9%)	12	11
All	All	520/586 (89%)	477 (92%)	43 (8%)	15	12

All (43) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ARG
1	A	47	TYR
1	A	48	ARG
1	A	49	THR
1	A	63	LYS
1	A	69	SER
1	A	116	PHE
1	A	149	THR
1	A	203	THR
1	A	212	THR
1	A	217	GLU
1	A	233	LYS
1	A	240	ARG
1	A	244	ARG
1	A	246	LYS
1	A	249	GLN
1	A	254	GLN
1	A	256	GLU
1	A	278	LYS
1	A	281	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	8	THR
1	B	20	ARG
1	B	22[A]	MET
1	B	22[B]	MET
1	B	26	GLU
1	B	47	TYR
1	B	48	ARG
1	B	61	LYS
1	B	63	LYS
1	B	70	THR
1	B	116	PHE
1	B	135	ASN
1	B	149	THR
1	B	194	GLN
1	B	203	THR
1	B	210	LYS
1	B	212	THR
1	B	221	LEU
1	B	240	ARG
1	B	245	ARG
1	B	249	GLN
1	B	254	GLN
1	B	281	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	111	ASN
1	A	220	ASN
1	A	264	GLN
1	B	18	ASN
1	B	111	ASN
1	B	150	HIS
1	B	194	GLN
1	B	220	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	5GP	A	593	3,2	21,26,26	1.57	5 (23%)	25,40,40	1.80	7 (28%)
3	ALF	A	594	2,5,4	0,4,4	0.00	-	0,6,6	0.00	-
4	5GP	B	593	3,2	21,26,26	1.60	4 (19%)	25,40,40	1.83	7 (28%)
3	ALF	B	594	2,5,4	0,4,4	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	5GP	A	593	3,2	-	0/6/26/26	0/3/3/3
3	ALF	A	594	2,5,4	-	0/0/0/0	0/0/0/0
4	5GP	B	593	3,2	-	0/6/26/26	0/3/3/3
3	ALF	B	594	2,5,4	-	0/0/0/0	0/0/0/0

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	593	5GP	P-O2P	2.12	1.62	1.54
4	A	593	5GP	O4'-C1'	2.42	1.44	1.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	B	593	5GP	C4-N3	2.65	1.39	1.35
4	A	593	5GP	C2-N1	2.90	1.40	1.35
4	A	593	5GP	C4-N3	2.93	1.40	1.35
4	B	593	5GP	C2-N1	3.04	1.40	1.35
4	B	593	5GP	O4'-C1'	3.10	1.45	1.41
4	A	593	5GP	C6-N1	4.16	1.40	1.33
4	B	593	5GP	C6-N1	4.21	1.40	1.33

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	593	5GP	N3-C2-N1	-5.26	119.44	127.44
4	B	593	5GP	N3-C2-N1	-5.06	119.74	127.44
4	B	593	5GP	C4-C5-N7	-3.16	106.57	109.48
4	B	593	5GP	C5-C6-N1	-3.04	119.42	123.59
4	A	593	5GP	C4-C5-N7	-3.00	106.72	109.48
4	A	593	5GP	C4'-O4'-C1'	-2.89	106.55	109.72
4	A	593	5GP	C5-C6-N1	-2.78	119.78	123.59
4	B	593	5GP	C4'-O4'-C1'	-2.51	106.96	109.72
4	B	593	5GP	C2'-C1'-N9	-2.42	110.59	114.29
4	A	593	5GP	C1'-N9-C4	-2.12	123.74	126.94
4	A	593	5GP	N2-C2-N1	2.00	120.52	117.20
4	B	593	5GP	N2-C2-N1	2.11	120.69	117.20
4	A	593	5GP	C6-N1-C2	3.03	120.14	115.94
4	B	593	5GP	C6-N1-C2	3.14	120.30	115.94

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	288/328 (87%)	0.30	28 (9%) 10 9	19, 27, 52, 69	1 (0%)
1	B	290/328 (88%)	0.33	31 (10%) 8 7	19, 27, 56, 69	0
All	All	578/656 (88%)	0.32	59 (10%) 9 8	19, 27, 55, 69	1 (0%)

All (59) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	212	THR	6.9
1	A	249	GLN	5.8
1	B	249	GLN	5.3
1	B	172	VAL	4.6
1	A	214	GLN	4.3
1	A	172	VAL	4.1
1	A	218	THR	4.0
1	B	43	ILE	3.9
1	A	44	VAL	3.9
1	A	127	VAL	3.8
1	A	43	ILE	3.7
1	A	217	GLU	3.6
1	A	215	LYS	3.5
1	A	42	ALA	3.5
1	B	127	VAL	3.3
1	A	221	LEU	3.3
1	B	210	LYS	3.3
1	A	41	VAL	3.3
1	B	246	LYS	3.3
1	B	216	ASP	3.3
1	B	251	GLU	3.3
1	B	211	GLY	3.3
1	B	41	VAL	3.2
1	A	306	SER	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	44	VAL	3.1
1	B	307	GLY	3.1
1	A	210	LYS	3.0
1	A	212	THR	2.9
1	B	180	TRP	2.9
1	A	126	PHE	2.9
1	A	191	ALA	2.8
1	A	180	TRP	2.8
1	A	40	VAL	2.7
1	B	217	GLU	2.7
1	B	306	SER	2.6
1	B	245	ARG	2.6
1	B	179	VAL	2.5
1	B	309	LEU	2.5
1	B	119	ALA	2.5
1	A	96	LEU	2.5
1	B	213	SER	2.4
1	B	130	SER	2.4
1	B	96	LEU	2.4
1	B	128	TYR	2.4
1	A	45	GLY	2.3
1	B	129	ASN	2.3
1	A	179	VAL	2.3
1	B	125	THR	2.2
1	A	120	VAL	2.2
1	A	46	LEU	2.2
1	A	128	TYR	2.2
1	B	45	GLY	2.1
1	A	49	THR	2.1
1	B	95	LEU	2.1
1	B	215	LYS	2.1
1	B	126	PHE	2.1
1	A	130	SER	2.0
1	A	156	SER	2.0
1	B	157	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	MG	B	595	1/1	0.95	0.19	0.59	24,24,24,24	0
2	MG	A	595	1/1	0.99	0.16	0.27	24,24,24,24	0
4	5GP	B	593	24/24	0.97	0.14	-0.23	19,24,27,28	0
3	ALF	A	594	5/5	0.99	0.12	-0.30	23,24,26,28	0
4	5GP	A	593	24/24	0.98	0.12	-0.41	17,24,26,28	0
3	ALF	B	594	5/5	0.98	0.11	-0.85	23,23,26,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.