



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:03 PM GMT

PDB ID : 4BCQ
Title : Structure of CDK2 in complex with cyclin A and a 2-amino-4-heteroaryl-pyrimidine inhibitor
Authors : Hole, A.J.; Baumli, S.; Wang, S.; Endicott, J.A.; Noble, M.E.M.
Deposited on : 2012-10-02
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 92%, green 5%, grey 4%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 92% 5% • </div> </div>
1	C	301	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 2%, orange 1%, yellow 89%, green 8%, grey 2%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> 2% 89% 8% • </div> </div>
2	B	262	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 1%, orange 1%, yellow 93%, green 5%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> % 93% 5% • </div> </div>
3	D	262	<div> <div style="width: 100%; height: 10px; background: linear-gradient(to right, red 3%, orange 1%, yellow 92%, green 5%, grey 1%);"></div> <div style="display: flex; justify-content: space-between; padding: 0 5px;"> 3% 92% 5% • </div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 9271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CYCLIN-DEPENDENT KINASE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	P	S	0	2	0
			2370	1538	404	419	1	8			
1	C	293	Total	C	N	O	P	S	0	1	0
			2366	1537	400	420	1	8			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	PRO	-	EXPRESSION TAG	UNP P24941
A	-1	GLY	-	EXPRESSION TAG	UNP P24941
A	0	SER	-	EXPRESSION TAG	UNP P24941
C	-2	PRO	-	EXPRESSION TAG	UNP P24941
C	-1	GLY	-	EXPRESSION TAG	UNP P24941
C	0	SER	-	EXPRESSION TAG	UNP P24941

- Molecule 2 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	257	Total	C	N	O	S	0	2	0
			2088	1352	340	385	11			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	432	LEU	-	EXPRESSION TAG	UNP P30274
B	303	THR	ALA	CONFLICT	UNP P30274
B	311	VAL	ILE	CONFLICT	UNP P30274
B	377	ILE	VAL	CONFLICT	UNP P30274
B	378	ARG	GLN	CONFLICT	UNP P30274
B	387	SER	LEU	CONFLICT	UNP P30274
B	391	MET	LEU	CONFLICT	UNP P30274
B	400	LYS	ARG	CONFLICT	UNP P30274


- Molecule 3 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	254	Total	C	N	O	S	0	0	0
			2054	1332	334	378	10			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	303	THR	ALA	CONFLICT	UNP P30274
D	311	VAL	ILE	CONFLICT	UNP P30274
D	377	ILE	VAL	CONFLICT	UNP P30274
D	378	ARG	GLN	CONFLICT	UNP P30274
D	386	SER	THR	CONFLICT	UNP P30274
D	390	MET	CYS	CONFLICT	UNP P30274
D	400	LYS	ARG	CONFLICT	UNP P30274

- Molecule 4 is 4-[4-METHYL-2-(METHYLAMINO)-1,3-THIAZOL-5-YL]-2-{{3-(MORPHOLIN-4-YLCARBONYL)PHENYL}AMINO}PYRIMIDINE-5-CARBONITRILE (three-letter code: TJF) (formula: C₂₁H₂₁N₇O₂S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			31	21	7	2	1		
4	C	1	Total	C	N	O	S	0	0
			31	21	7	2	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	127	Total 127	O 127	0	0
5	B	98	Total 98	O 98	0	0
5	C	57	Total 57	O 57	0	0
5	D	49	Total 49	O 49	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: CYCLIN-DEPENDENT KINASE 2




- Molecule 1: CYCLIN-DEPENDENT KINASE 2



- Molecule 2: CYCLIN-A2



- Molecule 3: CYCLIN-A2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	77.08Å 141.16Å 155.52Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.66 – 2.40 49.36 – 2.40	Depositor EDS
% Data completeness (in resolution range)	97.5 (48.66-2.40) 97.5 (49.36-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.81 (at 2.39Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.215 , 0.252 0.211 , 0.246	Depositor DCC
R_{free} test set	3321 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	43.8	Xtriage
Anisotropy	0.167	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 72.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 65384 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9271	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.65% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, TJF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2423	0.45	0/3286
1	C	0.24	0/2418	0.42	0/3279
2	B	0.27	0/2145	0.43	0/2912
3	D	0.24	0/2102	0.40	0/2851
All	All	0.26	0/9088	0.43	0/12328

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2370	0	2412	6	0
1	C	2366	0	2408	11	0
2	B	2088	0	2106	6	0
3	D	2054	0	2076	7	0
4	A	31	0	21	1	0
4	C	31	0	21	0	0
5	A	127	0	0	2	0
5	B	98	0	0	0	0
5	C	57	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	D	49	0	0	0	0
All	All	9271	0	9044	30	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:41:THR:HG22	1:C:42:GLU:HG2	1.67	0.76
1:C:72:THR:HG22	1:C:75:LYS:H	1.64	0.61
1:C:217:ARG:NH1	5:C:2055:HOH:O	2.37	0.58
1:A:40:GLU:HG3	1:A:42:GLU:H	1.69	0.57
1:C:74:ASN:ND2	5:C:2013:HOH:O	2.33	0.56
1:C:222:PRO:HG3	1:C:269:TYR:CZ	2.46	0.50
1:A:246:GLN:O	5:A:2088:HOH:O	2.20	0.49
1:A:223:ASP:H	1:A:226:VAL:HG12	1.82	0.45
1:C:181:SER:OG	1:C:182:THR:N	2.48	0.45
3:D:373:PRO:HD2	3:D:376:LEU:HD12	1.98	0.45
1:A:217:ARG:NH2	5:A:2086:HOH:O	2.38	0.45
4:A:1296:TJF:H27	4:A:1296:TJF:N09	2.32	0.44
1:C:154:VAL:HA	1:C:155:PRO:HA	1.85	0.44
3:D:375:SER:HA	3:D:378:ARG:HH11	1.81	0.44
1:C:237:LYS:HA	1:C:238:PRO:HD3	1.82	0.44
1:C:221:THR:HA	1:C:222:PRO:HD3	1.79	0.43
1:A:71:HIS:CD2	2:B:300:LYS:HE2	2.53	0.43
2:B:308:ALA:HA	2:B:309:PRO:HD3	1.91	0.43
2:B:176:PRO:HA	2:B:179[B]:HIS:CG	2.53	0.43
3:D:196:LYS:HE2	3:D:196:LYS:HB3	1.78	0.42
1:A:40:GLU:HG2	1:A:43:GLY:O	2.19	0.42
2:B:289:LYS:HE2	2:B:293:ARG:HE	1.84	0.42
2:B:194:LYS:HA	2:B:195:PRO:HD3	1.90	0.41
3:D:203:GLN:HA	3:D:204:PRO:HD3	1.88	0.41
3:D:272:PRO:HA	3:D:273:PRO:HD3	1.91	0.41
3:D:346:PRO:O	3:D:349:LYS:HG2	2.21	0.41
2:B:323:GLN:HA	2:B:324:PRO:HA	1.76	0.41
3:D:308:ALA:HA	3:D:309:PRO:HD3	1.87	0.41
1:C:288:ASP:OD1	1:C:288:ASP:N	2.54	0.40
1:C:256:ASP:HB3	1:C:257:GLU:H	1.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	290/301 (96%)	276 (95%)	13 (4%)	1 (0%)	46	63
1	C	289/301 (96%)	278 (96%)	10 (4%)	1 (0%)	46	63
2	B	257/262 (98%)	254 (99%)	3 (1%)	0	100	100
3	D	250/262 (95%)	245 (98%)	5 (2%)	0	100	100
All	All	1086/1126 (96%)	1053 (97%)	31 (3%)	2 (0%)	52	69

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	164	VAL
1	C	164	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	260/264 (98%)	254 (98%)	6 (2%)	58	78
1	C	259/264 (98%)	253 (98%)	6 (2%)	58	78
2	B	233/236 (99%)	230 (99%)	3 (1%)	76	89
3	D	228/236 (97%)	227 (100%)	1 (0%)	93	98
All	All	980/1000 (98%)	964 (98%)	16 (2%)	70	86

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	ILE
1	A	72	THR
1	A	122	ARG
1	A	248	PHE
1	A	293	VAL
1	A	295	HIS
2	B	177	ASP
2	B	334	MET
2	B	386	THR
1	C	72	THR
1	C	88	LYS
1	C	122	ARG
1	C	206	ASP
1	C	226	VAL
1	C	252	VAL
3	D	431	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	160	1	8,10,11	1.17	0	7,14,16	1.13	0
1	TPO	C	160	1	8,10,11	1.09	0	7,14,16	1.01	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical

component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/8/11/13	0/0/0/0
1	TPO	C	160	1	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	TJF	A	1296	-	29,34,34	2.28	7 (24%)	32,47,47	2.33	9 (28%)
4	TJF	C	1295	-	29,34,34	2.27	7 (24%)	32,47,47	2.37	10 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	TJF	A	1296	-	-	0/13/28/28	0/4/4/4
4	TJF	C	1295	-	-	0/13/28/28	0/4/4/4

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	1296	TJF	C20-N21	-6.24	1.36	1.47
4	C	1295	TJF	C20-N21	-6.07	1.36	1.47
4	C	1295	TJF	C25-N21	-6.05	1.36	1.47
4	A	1296	TJF	C25-N21	-6.05	1.36	1.47
4	A	1296	TJF	C07-C05	2.20	1.43	1.37
4	C	1295	TJF	C07-C05	2.21	1.44	1.37
4	A	1296	TJF	C30-C31	3.35	1.49	1.44
4	C	1295	TJF	C30-C31	3.37	1.49	1.44
4	A	1296	TJF	C22-N21	3.38	1.42	1.34
4	C	1295	TJF	C22-N21	3.76	1.43	1.34
4	C	1295	TJF	C10-N11	4.08	1.43	1.36
4	A	1296	TJF	C10-N11	4.12	1.43	1.36
4	C	1295	TJF	C03-N02	5.13	1.43	1.35
4	A	1296	TJF	C03-N02	5.23	1.43	1.35

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1296	TJF	N28-C10-N09	-5.94	120.34	126.67
4	C	1295	TJF	N28-C10-N09	-5.76	120.53	126.67
4	A	1296	TJF	C30-C29-N28	-3.39	119.22	124.30
4	C	1295	TJF	C30-C29-N28	-3.32	119.33	124.30
4	C	1295	TJF	O24-C22-N21	-2.69	118.15	122.42
4	C	1295	TJF	C01-N02-C03	-2.52	120.76	122.89
4	A	1296	TJF	O24-C22-N21	-2.51	118.43	122.42
4	C	1295	TJF	C25-N21-C22	-2.06	116.38	122.80
4	A	1296	TJF	C25-N21-C22	-2.04	116.44	122.80
4	C	1295	TJF	C29-N28-C10	2.85	121.40	115.95
4	A	1296	TJF	C29-N28-C10	2.86	121.41	115.95
4	A	1296	TJF	C07-C08-N09	3.01	119.05	114.97
4	C	1295	TJF	C07-C08-N09	3.02	119.05	114.97
4	A	1296	TJF	C11-C22-N21	3.08	122.86	118.76
4	C	1295	TJF	C11-C22-N21	3.83	123.87	118.76
4	C	1295	TJF	C08-N09-C10	5.70	121.47	116.66
4	A	1296	TJF	C08-N09-C10	5.93	121.67	116.66
4	C	1295	TJF	C25-N21-C20	6.03	123.74	112.56
4	A	1296	TJF	C25-N21-C20	6.40	124.43	112.56

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1296	TJF	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	292/301 (97%)	-0.25	4 (1%) 78 77	18, 31, 87, 129	0
1	C	292/301 (97%)	-0.12	7 (2%) 62 61	29, 52, 85, 115	0
2	B	257/262 (98%)	-0.21	2 (0%) 87 87	20, 33, 60, 99	0
3	D	254/262 (96%)	-0.08	9 (3%) 48 48	27, 53, 92, 120	0
All	All	1095/1126 (97%)	-0.17	22 (2%) 68 68	18, 42, 87, 129	0

All (22) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	39	THR	5.2
2	B	284	ASP	4.2
3	D	367	VAL	3.4
3	D	420	GLY	3.4
1	C	287	GLN	3.2
3	D	416	SER	3.2
1	C	282	ALA	3.1
3	D	284	ASP	2.7
3	D	421	VAL	2.7
1	C	258	ASP	2.7
1	C	288	ASP	2.7
1	C	250	LYS	2.6
3	D	430	LEU	2.5
1	A	73	GLU	2.4
3	D	415	ASN	2.4
1	A	38	ASP	2.3
2	B	323	GLN	2.2
3	D	423	LEU	2.1
3	D	369	GLY	2.1
1	C	257	GLU	2.1
1	A	72	THR	2.1
1	C	247	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	TPO	C	160	11/12	0.97	0.12	-	40,48,51,53	0
1	TPO	A	160	11/12	0.98	0.11	-	23,28,32,34	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	TJF	A	1296	31/31	0.85	0.18	1.36	56,63,89,90	0
4	TJF	C	1295	31/31	0.85	0.21	0.84	67,77,100,101	0

6.5 Other polymers [i](#)

There are no such residues in this entry.