wwPDB X-ray Structure Validation Summary Report (i)

Jan 31, 2016 – 06:34 PM GMT

PDB ID	:	1BH5
Title	:	HUMAN GLYOXALASE I Q33E, E172Q DOUBLE MUTANT
Authors	:	Cameron, A.D.; Jones, T.A.
Deposited on	:	1998-06-13
Resolution	:	2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	183	2%, 86%, 10%
1	B	183	4%, 85%, 11%
1	C	183	2%, 85%, 11%
1	D	183	8%, 80%, 17%

2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6317 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LACTOYLGLUTATHIONE LYASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	177	1406	895	230	271	10	0	0	0
1	B	182	1445	920	236	279	10	0	0	0
1	C	177	1406	895	230	271	10	0	0	0
1	D	182	1445	920	236	279	10	0	0	0

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	33	GLU	GLN	MUTATION	UNP Q04760
A	110	GLU	ALA	VARIANT	UNP Q04760
A	172	GLN	GLU	MUTATION	UNP Q04760
B	33	GLU	GLN	MUTATION	UNP Q04760
B	110	GLU	ALA	VARIANT	UNP Q04760
B	172	GLN	GLU	MUTATION	UNP Q04760
C	33	GLU	GLN	MUTATION	UNP Q04760
C	110	GLU	ALA	VARIANT	UNP Q04760
C	172	GLN	GLU	MUTATION	UNP Q04760
D	33	GLU	GLN	MUTATION	UNP Q04760
D	110	GLU	ALA	VARIANT	UNP Q04760
D	172	GLN	GLU	MUTATION	UNP Q04760

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	B	1	1	1	0	0
2	A	1	1	1	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Zn		
2	D	1	1	1	0	0
			Total	Zn		
2	C	1	1	1	0	0

- Molecule 3 is S-HEXYLGLUTATHIONE (three-letter code: GTX) (formula: $C_{16}H_{30}N_3O_6S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	26	16	3	6	1	0	0
			Total	C	N	O	S		
3	B	1	26	16	3	6	1	0	0
			Total	C	N	O	S		
3	C	1	26	16	3	6	1	0	0
			Total	C	N	O	S		
3	D	1	26	16	3	6	1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	141	141	141	0	0
			Total	O		
4	B	133	133	133	0	0
			Total	O		
4	C	130	130	130	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	D	103	103	103	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: LACTOYLGLUTATHIONE LYASE

Chain A:

- Molecule 1: LACTOYLGLUTATHIONE LYASE

Chain B:



- Molecule 1: LACTOYLGLUTATHIONE LYASE

Chain C:



- Molecule 1: LACTOYLGLUTATHIONE LYASE

Chain D:





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 41	Depositor
Cell constants a, b, c, α, β, γ	67.28Å 67.28Å 164.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.20 29.60 – 2.20	Depositor EDS
% Data completeness (in resolution range)	89.7 (30.00-2.20) 89.6 (29.60-2.20)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	5.51 (at 2.20Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.180 , 0.250 0.180 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å^2)	14.3	Xtriage
Anisotropy	0.751	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.29 , 49.2	EDS
Estimated twinning fraction	0.079 for h,-k,-l	Xtriage
L-test for twinning[2]	$< \|L\| > = 0.48$, $< L^2 > = 0.31$	Xtriage
Outliers	0 of 33195 reflections	Xtriage
F_o,F_c correlation	0.94	EDS
Total number of atoms	6317	wwPDB-VP
Average B, all atoms (Å^2)	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.84% of the height of the origin peak. No significant pseudotranslation is detected.*

[1] Intensities estimated from amplitudes.

[2] Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GTX, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\|>5$	RMSZ	$\#\|Z\|>5$
1	A	0.50	0/1438	1.21	8/1938 (0.4%)
1	B	0.51	0/1480	1.13	5/1998 (0.3%)
1	C	0.50	0/1438	1.19	5/1938 (0.3%)
1	D	0.52	0/1480	1.23	11/1998 (0.6%)
All	All	0.51	0/5836	1.19	29/7872 (0.4%)

There are no bond length outliers.

The worst 5 of 29 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
1	C	37	ARG	NE-CZ-NH2	-8.07	116.26	120.30
1	D	120	ASP	CB-CG-OD1	7.92	125.42	118.30
1	D	122	ARG	NE-CZ-NH1	7.77	124.18	120.30
1	C	140	ARG	NE-CZ-NH2	-7.35	116.62	120.30
1	A	76	ASP	CB-CG-OD1	7.14	124.73	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1406	0	1380	9	0
1	B	1445	0	1415	17	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1406	0	1380	10	0
1	D	1445	0	1415	18	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	26	0	28	1	0
3	B	26	0	28	2	0
3	C	26	0	28	3	0
3	D	26	0	28	0	0
4	A	141	0	0	0	0
4	B	133	0	0	4	1
4	C	130	0	0	0	0
4	D	103	0	0	2	0
All	All	6317	0	5702	47	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 47 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:200:GTX:HS62	1:B:92:LEU:HD11	1.62	0.81
1:B:2:GLU:HB3	1:B:3:PRO:HD3	1.70	0.72
1:C:179:MET:HE1	3:C:200:GTX:HS52	1.70	0.71
3:C:200:GTX:HS62	1:D:92:LEU:HD11	1.75	0.69
1:D:2:GLU:HB3	1:D:3:PRO:HD3	1.75	0.69

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:147:LYS:NZ	4:B:216:HOH:O[4_554]	2.01	0.19

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	175/183 (96%)	170 (97%)	5 (3%)	0	100	100
1	B	180/183 (98%)	176 (98%)	4 (2%)	0	100	100
1	C	175/183 (96%)	169 (97%)	5 (3%)	1 (1%)	30	29
1	D	180/183 (98%)	174 (97%)	5 (3%)	1 (1%)	30	29
All	All	710/732 (97%)	689 (97%)	19 (3%)	2 (0%)	46	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	156	LYS
1	C	182	LEU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	154/159 (97%)	150 (97%)	4 (3%)	54	66
1	B	159/159 (100%)	150 (94%)	9 (6%)	25	29
1	C	154/159 (97%)	150 (97%)	4 (3%)	54	66
1	D	159/159 (100%)	151 (95%)	8 (5%)	30	35
All	All	626/636 (98%)	601 (96%)	25 (4%)	38	47

5 of 25 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	179	MET
1	C	77	LYS
1	D	182	LEU
1	B	183	MET
1	C	83	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	58	GLN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

Of 8 ligands modelled in this entry, 4 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GTX	A	200	-	19,25,25	0.46	0	23,30,30	2.08	2 (8%)
3	GTX	B	200	-	19,25,25	0.69	0	23,30,30	2.77	5 (21%)
3	GTX	C	200	-	19,25,25	0.44	0	23,30,30	2.03	3 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\#\|Z\| > 2$	Counts	RMSZ	$\#\|Z\| > 2$
3	GTX	D	200	-	19,25,25	0.64	0	23,30,30	2.44	5 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GTX	A	200	-	-	0/24/30/30	0/0/0/0
3	GTX	B	200	-	-	0/24/30/30	0/0/0/0
3	GTX	C	200	-	-	0/24/30/30	0/0/0/0
3	GTX	D	200	-	-	0/24/30/30	0/0/0/0

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})
3	B	200	GTX	CA3-N3-C2	-10.60	107.77	122.34
3	D	200	GTX	CA3-N3-C2	-9.52	109.25	122.34
3	A	200	GTX	CA3-N3-C2	-8.20	111.06	122.34
3	C	200	GTX	CA3-N3-C2	-7.33	112.26	122.34
3	B	200	GTX	O2-C2-N3	-4.04	114.98	123.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	200	GTX	1	0
3	B	200	GTX	2	0
3	C	200	GTX	3	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	177/183 (96%)	-0.23	4 (2%)	64	63	7, 14, 24, 50	6 (3%)
1	B	182/183 (99%)	-0.21	8 (4%)	38	37	8, 14, 30, 48	7 (3%)
1	C	177/183 (96%)	-0.31	4 (2%)	64	63	8, 14, 24, 48	7 (3%)
1	D	182/183 (99%)	-0.05	14 (7%)	16	16	8, 14, 30, 47	6 (3%)
All	All	718/732 (98%)	-0.20	30 (4%)	40	39	7, 14, 28, 50	26 (3%)

The worst 5 of 30 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	7	SER	6.2
1	A	180	ALA	6.2
1	A	7	SER	4.9
1	B	183	MET	4.6
1	D	183	MET	4.3

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no carbohydrates in this entry.

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å^2)	Q<0.9
3	GTX	A	200	26/26	0.94	0.13	0.40	13,19,24,28	0
3	GTX	C	200	26/26	0.94	0.13	0.32	13,19,24,28	0
3	GTX	B	200	26/26	0.91	0.12	0.05	13,20,25,27	0
3	GTX	D	200	26/26	0.94	0.11	-0.30	14,20,25,27	0
2	ZN	B	201	1/1	0.99	0.04	-7.66	20,20,20,20	0
2	ZN	C	201	1/1	1.00	0.04	-	18,18,18,18	0
2	ZN	A	201	1/1	1.00	0.04	-	17,17,17,17	0
2	ZN	D	201	1/1	1.00	0.04	-	20,20,20,20	0

6.5 Other polymers (i)

There are no such residues in this entry.