



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:38 AM GMT

PDB ID : 3BJF  
Title : Pyruvate kinase M2 is a phosphotyrosine binding protein  
Authors : Wu, N.  
Deposited on : 2007-12-03  
Resolution : 2.03 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtrriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.03 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	9060 (2.04-2.00)
Ramachandran outliers	100387	8952 (2.04-2.00)
Sidechain outliers	100360	8951 (2.04-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	518	 77% 21% •
1	B	518	 78% 20% •
1	C	518	 74% 24% •
1	D	518	 73% 25% •

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 16573 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Pyruvate kinase isozymes M1/M2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	518	3963	2490	703	745	25	0	0	0
1	B	518	3963	2490	703	745	25	0	0	0
1	C	518	3963	2490	703	745	25	0	0	0
1	D	518	3963	2490	703	745	25	0	0	0

There are 4 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	379	ASN	HIS	CONFLICT	UNP P14618
B	379	ASN	HIS	CONFLICT	UNP P14618
C	379	ASN	HIS	CONFLICT	UNP P14618
D	379	ASN	HIS	CONFLICT	UNP P14618

- Molecule 2 is SUGAR (FRUCTOSE-1,6-DIPHOSPHATE) (three-letter code: FBP) (formula: C<sub>6</sub>H<sub>14</sub>O<sub>12</sub>P<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	O	P		
2	A	1	20	6	12	2	0	0
2	B	1	20	6	12	2	0	0
2	C	1	20	6	12	2	0	0
2	D	1	20	6	12	2	0	0

- Molecule 3 is OXALATE ION (three-letter code: OXL) (formula:  $C_2O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 2 4	0	0
3	B	1	Total C O 6 2 4	0	0
3	C	1	Total C O 6 2 4	0	0
3	D	1	Total C O 6 2 4	0	0

- Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total K 1 1	0	0
4	A	1	Total K 1 1	0	0
4	D	1	Total K 1 1	0	0
4	C	1	Total K 1 1	0	0

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	B	1	Total Mg 1 1	0	0
5	A	1	Total Mg 1 1	0	0
5	D	1	Total Mg 1 1	0	0
5	C	1	Total Mg 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	132	Total O 132 132	0	0
6	B	191	Total O 191 191	0	0
6	C	126	Total O 126 126	0	0

*Continued on next page...*

*Continued from previous page...*


<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	D	160	Total 160	O 160	0	0

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: Pyruvate kinase isozymes M1/M2




- Molecule 1: Pyruvate kinase isozymes M1/M2




- Molecule 1: Pyruvate kinase isozymes M1/M2





• Molecule 1: Pyruvate kinase isozymes M1/M2

Chain D: 73% 25%



## 4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	80.76Å 81.46Å 106.89Å 74.63° 69.91° 66.10°	Depositor
Resolution (Å)	44.29 – 2.03	Depositor
% Data completeness (in resolution range)	93.3 (44.29-2.03)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	CNS 1.2	Depositor
R, $R_{free}$	0.208 , 0.246	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	16573	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	29.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OXL, K, FBP, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.47	0/4026	0.69	2/5437 (0.0%)
1	B	0.54	1/4026 (0.0%)	0.74	2/5437 (0.0%)
1	C	0.48	0/4026	0.71	3/5437 (0.1%)
1	D	0.48	0/4026	0.70	1/5437 (0.0%)
All	All	0.49	1/16104 (0.0%)	0.71	8/21748 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	317	CYS	CB-SG	-6.49	1.71	1.82

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	294	ARG	NE-CZ-NH2	-9.13	115.73	120.30
1	C	467	ARG	NE-CZ-NH2	-8.56	116.02	120.30
1	A	467	ARG	NE-CZ-NH2	-6.84	116.88	120.30
1	D	294	ARG	NE-CZ-NH2	-6.75	116.92	120.30
1	C	294	ARG	NE-CZ-NH2	-6.53	117.04	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3963	0	4048	119	0
1	B	3963	0	4048	116	0
1	C	3963	0	4048	132	0
1	D	3963	0	4048	141	0
2	A	20	0	8	0	0
2	B	20	0	8	0	0
2	C	20	0	8	2	0
2	D	20	0	8	1	0
3	A	6	0	0	0	0
3	B	6	0	0	0	0
3	C	6	0	0	0	0
3	D	6	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	1	0	0	0	0
4	D	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
5	C	1	0	0	0	0
5	D	1	0	0	0	0
6	A	132	0	0	5	0
6	B	191	0	0	4	0
6	C	126	0	0	7	0
6	D	160	0	0	4	0
All	All	16573	0	16224	483	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 483 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:516:ARG:HH11	1:C:516:ARG:HB2	1.15	1.09
1:B:143:THR:HG22	1:B:145:ASP:H	1.13	1.09
1:A:64:MET:HE3	1:A:372:LEU:HD23	1.32	1.06
1:C:64:MET:HE3	1:C:372:LEU:HD23	1.40	1.02
1:D:255:ARG:HH11	1:D:255:ARG:HG3	1.31	0.96

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	516/518 (100%)	501 (97%)	13 (2%)	2 (0%)	39	32
1	B	516/518 (100%)	498 (96%)	14 (3%)	4 (1%)	24	15
1	C	516/518 (100%)	495 (96%)	16 (3%)	5 (1%)	19	10
1	D	516/518 (100%)	492 (95%)	16 (3%)	8 (2%)	12	4
All	All	2064/2072 (100%)	1986 (96%)	59 (3%)	19 (1%)	21	12

5 of 19 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	GLY
1	C	126	GLY
1	C	127	SER
1	B	128	GLY
1	C	130	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	426/426 (100%)	416 (98%)	10 (2%)	58	58
1	B	426/426 (100%)	412 (97%)	14 (3%)	45	42
1	C	426/426 (100%)	409 (96%)	17 (4%)	38	33
1	D	426/426 (100%)	416 (98%)	10 (2%)	58	58
All	All	1704/1704 (100%)	1653 (97%)	51 (3%)	48	46

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	525	MET
1	C	156	ILE
1	D	396	GLU
1	C	118	GLU
1	C	160	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 30 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	393	GLN
1	C	163	ASN
1	D	393	GLN
1	C	84	HIS
1	C	199	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	FBP	A	532	-	18,20,20	2.27	2 (11%)	21,32,32	2.32	8 (38%)
3	OXL	A	901	-	0,5,5	0.00	-	0,6,6	0.00	-
2	FBP	B	532	-	18,20,20	2.51	3 (16%)	21,32,32	3.81	13 (61%)
3	OXL	B	902	-	0,5,5	0.00	-	0,6,6	0.00	-
2	FBP	C	532	-	18,20,20	2.29	2 (11%)	21,32,32	2.32	7 (33%)
3	OXL	C	903	4	0,5,5	0.00	-	0,6,6	0.00	-
2	FBP	D	532	-	18,20,20	2.32	3 (16%)	21,32,32	2.39	7 (33%)
3	OXL	D	904	-	0,5,5	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FBP	A	532	-	-	0/13/32/32	0/1/1/1
3	OXL	A	901	-	-	0/0/4/4	0/0/0/0
2	FBP	B	532	-	-	0/13/32/32	0/1/1/1
3	OXL	B	902	-	-	0/0/4/4	0/0/0/0
2	FBP	C	532	-	-	0/13/32/32	0/1/1/1
3	OXL	C	903	4	-	0/0/4/4	0/0/0/0
2	FBP	D	532	-	-	0/13/32/32	0/1/1/1
3	OXL	D	904	-	-	0/0/4/4	0/0/0/0

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	532	FBP	O3-C3	-9.07	1.23	1.42
2	C	532	FBP	O3-C3	-8.37	1.25	1.42
2	D	532	FBP	O3-C3	-8.29	1.25	1.42
2	A	532	FBP	O3-C3	-8.27	1.25	1.42
2	B	532	FBP	P1-O1P	-2.34	1.43	1.51

The worst 5 of 35 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	532	FBP	O2P-P1-O1	-8.46	82.19	106.56
2	B	532	FBP	O1-P1-O1P	-7.71	87.51	107.14
2	B	532	FBP	O3P-P1-O1	-7.08	86.18	106.56
2	D	532	FBP	O6-C6-C5	-3.20	97.33	109.12
2	A	532	FBP	O6-C6-C5	-3.14	97.53	109.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	532	FBP	2	0
2	D	532	FBP	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers

EDS was not executed - this section will therefore be empty.