



wwPDB X-ray Structure Validation Summary Report ⓘ

Dec 15, 2016 – 10:41 AM EST

PDB ID : 2BKK
Title : Crystal structure of Aminoglycoside Phosphotransferase APH(3')-IIIa in complex with the designed ankyrin repeat inhibitor AR_3a
Authors : Kohl, A.; Amstutz, P.; Parizek, P.; Binz, H.K.; Briand, C.; Capitani, G.; Forrer, P.; Pluckthun, A.; Grutter, M.G.
Deposited on : 2005-02-16
Resolution : 2.15 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	unknown
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20028442
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20028442


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	264	<div> <div>5%</div> <div>69%</div> <div>20%</div> <div>6%</div> </div>
1	C	264	<div> <div>9%</div> <div>72%</div> <div>20%</div> <div>5%</div> <div>••</div> </div>
2	B	169	<div> <div>20%</div> <div>61%</div> <div>28%</div> <div>8%</div> <div>•</div> </div>
2	D	169	<div> <div>24%</div> <div>53%</div> <div>30%</div> <div>8%</div> <div>• 7%</div> </div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 7000 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called AMINOGLYCOSIDE 3'-PHOSPHOTRANSFERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	247	Total	C	N	O	S	0	0	0
			2038	1304	326	398	10			
1	C	262	Total	C	N	O	S	0	0	0
			2165	1382	343	430	10			


There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	19	SER	CYS	ENGINEERED MUTATION	UNP P00554
C	19	SER	CYS	ENGINEERED MUTATION	UNP P00554

- Molecule 2 is a protein called DESIGNED ANKYRIN REPEAT INHIBITOR AR_3A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	156	Total	C	N	O	S	0	0	1
			1194	756	201	235	2			
2	D	157	Total	C	N	O	S	0	0	1
			1198	758	202	236	2			

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	2	Total	Mg	0	0
			2	2		
4	C	2	Total	Mg	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	115	Total	O	0	0
			115	115		
5	B	53	Total	O	0	0
			53	53		
5	C	130	Total	O	0	0
			130	130		
5	D	49	Total	O	0	0
			49	49		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: AMINOGLYCOSIDE 3'-PHOSPHOTRANSFERASE




• Molecule 1: AMINOGLYCOSIDE 3'-PHOSPHOTRANSFERASE



• Molecule 2: DESIGNED ANKYRIN REPEAT INHIBITOR AR_3A



• Molecule 2: DESIGNED ANKYRIN REPEAT INHIBITOR AR_3A



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	59.67Å 98.08Å 81.30Å 90.00° 110.01° 90.00°	Depositor
Resolution (Å)	20.00 – 2.15 19.90 – 2.15	Depositor EDS
% Data completeness (in resolution range)	96.7 (20.00-2.15) 96.6 (19.90-2.15)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.49 (at 2.15Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.199 , 0.260 0.206 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	44.5	Xtriage
Anisotropy	0.130	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 57.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.032 for h,-k,-h-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7000	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.49% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.71	1/2081 (0.0%)	0.95	11/2800 (0.4%)
1	C	0.73	0/2213	0.95	9/2984 (0.3%)
2	B	0.63	0/1215	0.89	8/1652 (0.5%)
2	D	0.66	0/1219	0.98	13/1657 (0.8%)
All	All	0.70	1/6728 (0.0%)	0.94	41/9093 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	C	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	64	MET	SD-CE	-5.62	1.46	1.77

The worst 5 of 41 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	216	ASP	CB-CG-OD2	9.11	126.50	118.30
1	C	220	ASP	CB-CG-OD2	8.04	125.54	118.30
1	A	59	ARG	NE-CZ-NH2	-7.85	116.37	120.30
1	A	56	ASP	CB-CG-OD2	7.73	125.25	118.30
2	D	72	ASP	CB-CG-OD2	7.49	125.04	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	C	152	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2038	0	1992	39	0
1	C	2165	0	2091	39	0
2	B	1194	0	1162	35	0
2	D	1198	0	1165	45	0
3	A	27	0	12	1	0
3	C	27	0	12	0	0
4	A	2	0	0	0	0
4	C	2	0	0	0	0
5	A	115	0	0	12	1
5	B	53	0	0	11	0
5	C	130	0	0	8	1
5	D	49	0	0	13	0
All	All	7000	0	6434	154	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 154 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:146:LEU:HA	5:C:2081:HOH:O	1.34	1.23
2:D:82:THR:OG1	5:D:2030:HOH:O	1.74	1.04
2:D:116:PRO:HD3	5:D:2039:HOH:O	1.59	1.02
2:D:41:ASN:HD21	2:D:72:ASP:H	1.12	0.96
2:B:44:ASP:O	5:B:2025:HOH:O	1.84	0.93

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2109:HOH:O	5:C:2056:HOH:O[2_545]	1.89	0.31

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	243/264 (92%)	229 (94%)	12 (5%)	2 (1%)	24	15
1	C	260/264 (98%)	241 (93%)	10 (4%)	9 (4%)	4	1
2	B	154/169 (91%)	141 (92%)	11 (7%)	2 (1%)	15	8
2	D	155/169 (92%)	130 (84%)	22 (14%)	3 (2%)	10	3
All	All	812/866 (94%)	741 (91%)	55 (7%)	16 (2%)	9	3

5 of 16 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	25	GLY
1	C	4	MET
1	C	153	ASP
1	C	163	THR
1	C	193	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	226/242 (93%)	199 (88%)	27 (12%)	6 2

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	241/242 (100%)	207 (86%)	34 (14%)	4	1
2	B	123/135 (91%)	110 (89%)	13 (11%)	8	4
2	D	123/135 (91%)	99 (80%)	24 (20%)	2	0
All	All	713/754 (95%)	615 (86%)	98 (14%)	4	1

5 of 98 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	16	LYS
1	C	111	ILE
2	D	144	LYS
1	C	24	GLU
1	C	61	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	118	HIS
1	C	78	HIS
2	D	41	ASN
2	B	107	ASN
2	D	56	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	ADP	A	1265	4	24,29,29	1.41	3 (12%)	23,45,45	2.25	2 (8%)
3	ADP	C	1265	4	24,29,29	1.10	2 (8%)	23,45,45	3.50	3 (13%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1265	4	-	0/12/32/32	0/3/3/3
3	ADP	C	1265	4	-	0/12/32/32	0/3/3/3

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1265	ADP	O4'-C1'	2.35	1.44	1.41
3	C	1265	ADP	C2-N3	2.43	1.36	1.32
3	C	1265	ADP	C2-N1	2.68	1.39	1.33
3	A	1265	ADP	C2-N1	3.41	1.40	1.33
3	A	1265	ADP	C2-N3	3.72	1.38	1.32

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	1265	ADP	N3-C2-N1	-15.67	116.56	128.87
3	A	1265	ADP	N3-C2-N1	-9.66	121.28	128.87
3	C	1265	ADP	O4'-C1'-N9	2.38	112.59	108.11
3	A	1265	ADP	O2A-PA-O3A	2.51	116.02	105.27
3	C	1265	ADP	C2-N1-C6	3.46	124.94	118.77

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1265	ADP	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/264 (93%)	0.38	13 (5%) 30 40	29, 50, 87, 115	0
1	C	262/264 (99%)	0.48	25 (9%) 10 16	30, 47, 83, 112	1 (0%)
2	B	156/169 (92%)	1.09	34 (21%) 1 2	34, 59, 117, 123	0
2	D	157/169 (92%)	1.19	40 (25%) 1 1	31, 59, 117, 128	0
All	All	822/866 (94%)	0.70	112 (13%) 4 7	29, 52, 105, 128	1 (0%)

The worst 5 of 112 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	167	LYS	15.5
1	C	152	ALA	9.3
1	A	264	PHE	8.2
1	C	26	MET	6.8
2	B	145	PHE	6.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	ADP	C	1265	27/27	0.91	0.13	-0.50	26,42,71,74	0
4	MG	A	1267	1/1	0.92	0.12	-0.77	60,60,60,60	0
3	ADP	A	1265	27/27	0.95	0.08	-1.45	30,38,48,53	0
4	MG	C	1266	1/1	0.88	0.08	-	64,64,64,64	0
4	MG	C	1267	1/1	0.52	0.33	-	91,91,91,91	0
4	MG	A	1266	1/1	0.94	0.04	-	50,50,50,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.