



Full wwPDB X-ray Structure Validation Report ⓘ

Sep 26, 2016 – 10:09 PM EDT

PDB ID : 5BMM
Title : Src in complex with DNA-templated macrocyclic inhibitor MC25b
Authors : Georghiou, G.; Guja, K.E.; Aleem, S.; Kleiner, R.E.; Liu, D.R.; Miller, W.T.;
Garcia-Diaz, M.; Seeliger, M.A.
Deposited on : 2015-05-22
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3553 (2.50-2.50)
Clashscore	102246	4242 (2.50-2.50)
Ramachandran outliers	100387	4156 (2.50-2.50)
Sidechain outliers	100360	4158 (2.50-2.50)
RSRZ outliers	91569	3562 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	286	
1	B	286	
2	C	5	
2	D	5	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4534 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2147	1378	354	398	17			
1	B	275	Total	C	N	O	S	0	0	0
			2153	1381	361	394	17			

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523



- Molecule 2 is a protein called macrocyclic inhibitor MC25b.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	5	Total	C	F	N	O	0	0	0
			52	36	1	6	9			
2	D	5	Total	C	F	N	O	0	0	0
			52	36	1	6	9			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	58	Total	O	0	0
			58	58		
3	B	66	Total	O	0	0
			66	66		
3	C	6	Total	O	0	0
			6	6		

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	60.83Å 118.61Å 42.65Å 90.00° 90.08° 90.00°	Depositor
Resolution (Å)	42.65 – 2.50 42.65 – 2.50	Depositor EDS
% Data completeness (in resolution range)	83.7 (42.65-2.50) 83.6 (42.65-2.50)	Depositor EDS
R_{merge}	0.25	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.99 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.218 , 0.246 0.217 , 0.248	Depositor DCC
R_{free} test set	871 reflections (4.98%)	DCC
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.103	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 18.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.238 for -h,-k,l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	4534	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.07% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: ORN, ALC, FUM, PPN, PFF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.23	0/2199	0.43	0/2987
1	B	0.23	0/2205	0.43	0/2994
All	All	0.23	0/4404	0.43	0/5981

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2147	0	2084	21	0
1	B	2153	0	2105	27	0
2	C	52	0	38	3	0
2	D	52	0	39	4	0
3	A	58	0	0	1	0
3	B	66	0	0	0	0
3	C	6	0	0	1	0
All	All	4534	0	4266	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:264:ARG:HH22	1:A:333:PRO:HD2	1.44	0.82
1:B:264:ARG:NH1	1:B:333:PRO:O	2.19	0.77
1:A:320:GLU:O	1:A:401:LYS:HE2	1.87	0.74
1:A:264:ARG:HD3	1:A:329:VAL:HG11	1.70	0.71
1:B:377:VAL:HG13	1:B:382:TYR:HB3	1.72	0.70
1:B:310:GLU:OE1	1:B:409:ARG:NH2	2.24	0.70
2:C:5:ORN:OXT	3:C:101:HOH:O	2.15	0.65
1:A:310:GLU:OE1	1:A:409:ARG:NH2	2.28	0.65
1:A:297:LEU:HD22	1:A:301:THR:HG21	1.79	0.65
1:B:320:GLU:O	1:B:401:LYS:HE2	1.97	0.64
1:A:377:VAL:HG13	1:A:382:TYR:HB3	1.78	0.64
1:B:353:GLU:OE1	1:B:356:LYS:NZ	2.33	0.61
1:B:339:GLU:OE2	1:B:401:LYS:NZ	2.35	0.59
1:B:323:VAL:HG21	1:B:393:LEU:HD12	1.85	0.57
1:B:273:LEU:HB2	1:B:281:VAL:HG13	1.86	0.56
1:A:378:GLU:HG3	1:A:441:ILE:HG12	1.87	0.56
1:B:378:GLU:HG3	1:B:441:ILE:HG12	1.88	0.55
1:A:264:ARG:NH2	1:A:333:PRO:HD2	2.20	0.52
1:B:281:VAL:HG11	2:D:4:PPN:CZ	2.39	0.52
1:A:468:ASN:ND2	3:A:604:HOH:O	2.34	0.51
1:B:341:MET:HG3	1:B:393:LEU:HB3	1.92	0.51
1:B:419:ARG:NH2	1:B:435:LEU:O	2.45	0.50
1:B:297:LEU:HD22	1:B:301:THR:HG21	1.94	0.49
1:A:272:LYS:HE3	1:A:275:GLN:HB3	1.93	0.49
1:A:270:GLU:HG3	1:A:284:GLY:HA2	1.96	0.48
1:B:283:MET:HG3	1:B:340:TYR:CE1	2.48	0.47
1:A:388:ARG:HH22	2:C:1:FUM:H5	1.79	0.47
1:A:266:SER:OG	1:A:286:TRP:O	2.32	0.47
1:B:308:LEU:O	1:B:312:GLN:HG2	2.14	0.47
1:B:384:HIS:O	1:B:385:ARG:HB2	2.15	0.46
1:A:341:MET:HG3	1:A:393:LEU:HB3	1.98	0.46
1:B:351:LYS:HA	1:B:351:LYS:HD3	1.81	0.46
1:B:321:LYS:HD2	1:B:369:GLN:HB3	1.98	0.45
1:A:419:ARG:NH1	1:A:435:LEU:O	2.49	0.45
1:B:408:ALA:HA	2:D:2:ALC:HZ2	1.99	0.44
1:A:384:HIS:O	1:A:385:ARG:HB2	2.18	0.43
1:B:491:LEU:HA	1:B:491:LEU:HD12	1.90	0.43
1:A:323:VAL:HG21	1:A:393:LEU:HD12	2.01	0.43
1:B:467:VAL:O	1:B:471:VAL:HG23	2.20	0.42
1:A:281:VAL:HG11	2:C:4:PPN:CZ	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:441:ILE:O	1:B:445:VAL:HG23	2.20	0.42
1:B:411:ILE:HD11	2:D:2:ALC:HE13	2.02	0.41
1:A:276:GLY:N	1:A:279:GLY:O	2.45	0.41
1:A:353:GLU:O	1:A:356:LYS:HG2	2.20	0.41
1:B:347:LEU:HD12	1:B:347:LEU:HA	1.95	0.41
1:B:383:VAL:HG12	1:B:385:ARG:HG3	2.03	0.41
1:A:321:LYS:HD3	1:A:321:LYS:HA	1.87	0.41
1:B:500:ARG:HD3	1:B:505:GLU:HB3	2.02	0.40
1:B:347:LEU:HD23	2:D:5:ORN:HB2	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/286 (96%)	270 (99%)	3 (1%)	0	100	100
1	B	273/286 (96%)	270 (99%)	3 (1%)	0	100	100
All	All	546/572 (96%)	540 (99%)	6 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/245 (91%)	218 (98%)	5 (2%)	60	84
1	B	224/245 (91%)	223 (100%)	1 (0%)	93	98
All	All	447/490 (91%)	441 (99%)	6 (1%)	76	92

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	303	SER
1	A	317	LEU
1	A	419	ARG
1	A	472	LEU
1	A	510	GLU
1	B	419	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

8 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	ALC	C	2	2	9,11,12	0.80	1 (11%)	11,13,15	0.88	1 (9%)
2	PFF	C	3	2	10,12,13	1.10	1 (10%)	13,15,17	1.17	0
2	PPN	C	4	2	10,14,15	0.84	0	14,18,20	0.93	1 (7%)
2	ORN	C	5	2	5,8,8	0.76	0	3,9,9	0.40	0
2	ALC	D	2	2	9,11,12	0.74	1 (11%)	11,13,15	0.94	1 (9%)
2	PFF	D	3	2	10,12,13	1.14	1 (10%)	13,15,17	1.25	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PPN	D	4	2	10,14,15	0.88	0	14,18,20	1.19	2 (14%)
2	ORN	D	5	2	5,8,8	0.77	0	3,9,9	0.55	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	ALC	C	2	2	-	0/4/14/16	0/1/1/1
2	PFF	C	3	2	-	0/4/6/8	0/1/1/1
2	PPN	C	4	2	-	0/8/10/12	0/1/1/1
2	ORN	C	5	2	-	0/4/8/8	0/0/0/0
2	ALC	D	2	2	-	0/4/14/16	0/1/1/1
2	PFF	D	3	2	-	0/4/6/8	0/1/1/1
2	PPN	D	4	2	-	0/8/10/12	0/1/1/1
2	ORN	D	5	2	-	0/4/8/8	0/0/0/0

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	3	PFF	CB-CA	-2.31	1.48	1.53
2	C	3	PFF	CB-CA	-2.09	1.49	1.53
2	D	2	ALC	CB-CG	2.06	1.56	1.53
2	C	2	ALC	CB-CG	2.25	1.57	1.53

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	4	PPN	O-C-CA	-3.24	117.04	125.72
2	C	4	PPN	O-C-CA	-2.20	119.81	125.72
2	D	3	PFF	CG-CB-CA	-2.13	109.21	114.12
2	C	2	ALC	O-C-CA	-2.10	120.10	125.72
2	D	2	ALC	O-C-CA	-2.01	120.34	125.72
2	D	4	PPN	O1-N1-CZ	2.06	120.15	118.67

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	4	PPN	1	0
2	C	5	ORN	1	0
2	D	2	ALC	2	0
2	D	4	PPN	1	0
2	D	5	ORN	1	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	D	1
2	C	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	4:PPN	C	5:ORN	N	5.15
1	D	4:PPN	C	5:ORN	N	3.76

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/286 (96%)	-0.23	1 (0%) 93 93	20, 36, 56, 71	0
1	B	275/286 (96%)	-0.19	4 (1%) 76 79	20, 35, 54, 66	0
2	C	0/5	-	-	-	-
2	D	0/5	-	-	-	-
All	All	550/582 (94%)	-0.21	5 (0%) 85 88	20, 35, 56, 71	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	421	GLY	3.1
1	B	307	PHE	2.3
1	B	289	THR	2.3
1	A	421	GLY	2.2
1	B	465	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	PFF	C	3	12/13	0.93	0.15	-	32,34,36,36	0
2	PFF	D	3	12/13	0.93	0.20	-	26,30,32,32	0
2	ORN	D	5	9/9	0.83	0.30	-	45,58,63,65	0
2	PPN	C	4	14/15	0.91	0.15	-	28,30,37,39	0
2	PPN	D	4	14/15	0.94	0.17	-	22,25,41,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	ORN	C	5	9/9	0.86	0.20	-	45,58,65,65	0
2	ALC	D	2	11/12	0.93	0.17	-	28,32,37,37	0
2	ALC	C	2	11/12	0.91	0.20	-	35,39,43,43	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.