



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 12:48 AM GMT

PDB ID : 2BR5  
Title : CMCI-N160 SAH  
Authors : Oster, L.M.; Lester, D.R.; Terwisscha Van Scheltinga, A.; Svenda, M.;  
Genereux, C.; Andersson, I.  
Deposited on : 2005-05-01  
Resolution : 2.83 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.83 Å.


Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	3658 (2.88-2.80)
Ramachandran outliers	100387	3591 (2.88-2.80)
Sidechain outliers	100360	3594 (2.88-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	236	 78% 15% . .
1	B	236	 73% 21% 5% .
1	C	236	 69% 23% . .
1	D	236	 60% 19% . 18%
1	E	236	 71% 25% . .
1	F	236	 67% 17% . 13%

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 11055 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CEPHALOSPORIN HYDROXYLASE CMCI.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1861	1188	320	341	12			
1	B	232	Total	C	N	O	S	0	0	0
			1909	1216	328	353	12			
1	C	227	Total	C	N	O	S	0	0	0
			1860	1188	318	342	12			
1	D	194	Total	C	N	O	S	0	0	0
			1588	1009	274	293	12			
1	E	230	Total	C	N	O	S	0	0	0
			1896	1209	325	350	12			
1	F	206	Total	C	N	O	S	0	0	0
			1686	1079	288	307	12			

There are 15 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	10	GLN	LEU	ENGINEERED MUTATION	UNP O85726
A	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
A	200	PHE	LEU	ENGINEERED MUTATION	UNP O85726
B	10	GLN	LEU	ENGINEERED MUTATION	UNP O85726
B	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
B	200	PHE	LEU	ENGINEERED MUTATION	UNP O85726
C	10	GLN	LEU	ENGINEERED MUTATION	UNP O85726
C	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
C	200	PHE	LEU	ENGINEERED MUTATION	UNP O85726
D	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
E	10	GLN	LEU	ENGINEERED MUTATION	UNP O85726
E	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
E	200	PHE	LEU	ENGINEERED MUTATION	UNP O85726
F	160	ASN	ASP	ENGINEERED MUTATION	UNP O85726
F	200	PHE	LEU	ENGINEERED MUTATION	UNP O85726

- Molecule 2 is S-ADENOSYL-L-HOMOCYSTEINE (three-letter code: SAH) (formula:

C<sub>14</sub>H<sub>20</sub>N<sub>6</sub>O<sub>5</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	B	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	C	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	D	1	Total	C	N	O	S	0	0
			26	14	6	5	1		
2	E	1	Total	C	N	O	S	0	0
			26	14	6	5	1		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	17	Total	O	0	0
			17	17		
3	B	28	Total	O	0	0
			28	28		
3	C	35	Total	O	0	0
			35	35		
3	D	9	Total	O	0	0
			9	9		
3	E	14	Total	O	0	0
			14	14		
3	F	22	Total	O	0	0
			22	22		

### 3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI




#### • Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI




#### • Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI




#### • Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI






• Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI

Chain E: 71% 25%



• Molecule 1: CEPHALOSPORIN HYDROXYLASE CMCI

Chain F: 67% 17% 13%



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	90.91Å 102.52Å 181.41Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.39 – 2.83	Depositor
% Data completeness (in resolution range)	99.0 (49.39-2.83)	Depositor
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.231 , 0.301	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	11055	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	23.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SAH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.48	0/1915	0.84	6/2605 (0.2%)
1	B	0.54	0/1964	0.87	10/2672 (0.4%)
1	C	0.56	0/1914	0.87	9/2605 (0.3%)
1	D	0.45	0/1631	0.84	9/2216 (0.4%)
1	E	0.51	0/1951	0.86	14/2654 (0.5%)
1	F	0.53	0/1736	0.84	8/2364 (0.3%)
All	All	0.51	0/11111	0.86	56/15116 (0.4%)

There are no bond length outliers.

The worst 5 of 56 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	138	ASP	CB-CG-OD2	7.09	124.68	118.30
1	D	39	ASP	CB-CG-OD2	6.97	124.58	118.30
1	C	141	ASP	CB-CG-OD2	6.67	124.30	118.30
1	B	116	ASP	CB-CG-OD2	6.37	124.03	118.30
1	E	159	ASP	CB-CG-OD2	6.24	123.91	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1861	0	1771	16	0
1	B	1909	0	1805	23	0
1	C	1860	0	1765	26	0
1	D	1588	0	1508	17	0
1	E	1896	0	1794	17	0
1	F	1686	0	1599	16	0
2	A	26	0	19	1	0
2	B	26	0	19	0	0
2	C	26	0	19	1	0
2	D	26	0	19	1	0
2	E	26	0	19	3	0
3	A	17	0	0	0	0
3	B	28	0	0	0	0
3	C	35	0	0	3	0
3	D	9	0	0	0	0
3	E	14	0	0	3	0
3	F	22	0	0	0	0
All	All	11055	0	10337	104	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 104 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:164:ASN:HD21	1:E:167:ASN:HD22	1.22	0.85
1:B:66:ASP:HB2	1:B:67:PRO:HD2	1.66	0.76
1:E:88:LEU:HD22	1:E:139:CYS:SG	2.30	0.72
1:A:105:ILE:HD11	1:B:105:ILE:CD1	2.21	0.70
1:C:87:GLU:OE1	1:C:160:ASN:ND2	2.23	0.67

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/236 (95%)	215 (96%)	8 (4%)	1 (0%)	39	72
1	B	230/236 (98%)	218 (95%)	9 (4%)	3 (1%)	15	42
1	C	225/236 (95%)	217 (96%)	6 (3%)	2 (1%)	21	52
1	D	188/236 (80%)	170 (90%)	15 (8%)	3 (2%)	12	36
1	E	228/236 (97%)	211 (92%)	17 (8%)	0	100	100
1	F	202/236 (86%)	188 (93%)	13 (6%)	1 (0%)	34	67
All	All	1297/1416 (92%)	1219 (94%)	68 (5%)	10 (1%)	24	56

5 of 10 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	146	GLU
1	A	150	GLU
1	C	14	LEU
1	D	187	ASP
1	F	138	ASP

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	198/207 (96%)	176 (89%)	22 (11%)	8	21
1	B	203/207 (98%)	171 (84%)	32 (16%)	3	8
1	C	197/207 (95%)	169 (86%)	28 (14%)	4	11
1	D	170/207 (82%)	142 (84%)	28 (16%)	3	7
1	E	202/207 (98%)	174 (86%)	28 (14%)	4	12
1	F	178/207 (86%)	155 (87%)	23 (13%)	5	14
All	All	1148/1242 (92%)	987 (86%)	161 (14%)	4	12

5 of 161 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	147	HIS
1	D	88	LEU
1	F	87	GLU
1	C	169	MET
1	C	231	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 29 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	162	HIS
1	D	136	GLN
1	F	164	ASN
1	D	57	GLN
1	D	219	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	SAH	A	301	-	20,28,28	1.13	2 (10%)	19,40,40	2.77	2 (10%)
2	SAH	B	301	-	20,28,28	1.08	2 (10%)	19,40,40	2.97	1 (5%)
2	SAH	C	301	-	20,28,28	1.06	2 (10%)	19,40,40	3.26	5 (26%)
2	SAH	D	301	-	20,28,28	1.13	2 (10%)	19,40,40	2.98	2 (10%)
2	SAH	E	301	-	20,28,28	1.08	2 (10%)	19,40,40	3.39	5 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SAH	A	301	-	-	0/7/31/31	0/3/3/3
2	SAH	B	301	-	-	0/7/31/31	0/3/3/3
2	SAH	C	301	-	-	0/7/31/31	0/3/3/3
2	SAH	D	301	-	-	0/7/31/31	0/3/3/3
2	SAH	E	301	-	-	0/7/31/31	0/3/3/3

The worst 5 of 10 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	E	301	SAH	C2-N1	2.04	1.37	1.33
2	B	301	SAH	C2-N1	2.12	1.37	1.33
2	C	301	SAH	C2-N1	2.28	1.38	1.33
2	A	301	SAH	C2-N1	2.44	1.38	1.33
2	D	301	SAH	C2-N1	2.44	1.38	1.33

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	301	SAH	N3-C2-N1	-13.07	118.89	128.89
2	B	301	SAH	N3-C2-N1	-12.27	119.50	128.89
2	C	301	SAH	N3-C2-N1	-12.25	119.51	128.89
2	D	301	SAH	N3-C2-N1	-11.98	119.72	128.89
2	A	301	SAH	N3-C2-N1	-11.37	120.19	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 6 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	SAH	1	0
2	C	301	SAH	1	0
2	D	301	SAH	1	0
2	E	301	SAH	3	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates ⓘ

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands ⓘ

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers ⓘ

EDS was not executed - this section will therefore be empty.