



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:11 PM GMT

PDB ID : 4BRD  
Title : Legionella pneumophila NTPDase1 Q193E crystal form II, closed, Mg  
AMPPNP complex  
Authors : Zebisch, M.; Schaefer, P.; Lauble, P.; Straeter, N.  
Deposited on : 2013-06-04  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	<div> <div>6%</div> <div> <div></div> <div>86%</div> <div>11%</div> <div>••</div> </div> </div>
1	B	368	<div> <div>4%</div> <div> <div></div> <div>88%</div> <div>9%</div> <div>•</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	B	506	-	-	-	X

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6581 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	359	Total	C	N	O	S	0	18	0
			2956	1886	477	577	16			
1	B	358	Total	C	N	O	S	0	20	0
			2947	1880	477	574	16			


There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	34	MET	-	EXPRESSION TAG	UNP Q5ZUA2
A	394	LEU	-	EXPRESSION TAG	UNP Q5ZUA2
A	395	GLU	-	EXPRESSION TAG	UNP Q5ZUA2
A	396	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	397	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	398	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	399	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	400	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	401	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
A	137	ASP	GLU	CONFLICT	UNP Q5ZUA2
A	149	VAL	ALA	CONFLICT	UNP Q5ZUA2
A	193	GLU	GLN	ENGINEERED MUTATION	UNP Q5ZUA2
B	34	MET	-	EXPRESSION TAG	UNP Q5ZUA2
B	394	LEU	-	EXPRESSION TAG	UNP Q5ZUA2
B	395	GLU	-	EXPRESSION TAG	UNP Q5ZUA2
B	396	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	397	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	398	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	399	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	400	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	401	HIS	-	EXPRESSION TAG	UNP Q5ZUA2
B	137	ASP	GLU	CONFLICT	UNP Q5ZUA2
B	149	VAL	ALA	CONFLICT	UNP Q5ZUA2
B	193	GLU	GLN	ENGINEERED MUTATION	UNP Q5ZUA2

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	3	Total	Mg	0	0
			3	3		
2	A	2	Total	Mg	0	0
			2	2		

- Molecule 3 is PHOSPHOAMINOPHOSPHONIC ACID-ADENYLATE ESTER (three-letter code: ANP) (formula:  $C_{10}H_{17}N_6O_{12}P_3$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	1
			62	20	12	24	6		
3	B	1	Total	C	N	O	P	0	0
			31	10	6	12	3		

- Molecule 4 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (three-letter code: MES) (formula:  $C_6H_{13}NO_4S$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
4	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Cl	0	0
			1	1		
5	A	1	Total	Cl	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	276	Total	O	0	2
			278	278		
6	B	276	Total	O	0	0
			276	276		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



#### • Molecule 1: ECTONUCLEOSIDE TRIPHOSPHATE DIPHOSPHOHYDROLASE I



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	62.13Å 86.05Å 71.98Å 90.00° 106.78° 90.00°	Depositor
Resolution (Å)	29.20 – 1.50 28.94 – 1.50	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.20-1.50) 99.4 (28.94-1.50)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.73 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.140 , 0.197 0.137 , 0.193	Depositor DCC
$R_{free}$ test set	2315 reflections (2.05%)	DCC
Wilson B-factor (Å <sup>2</sup> )	14.8	Xtriage
Anisotropy	0.535	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 45.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 115329 reflections	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	6581	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ANP, MES, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.95	0/3088	0.93	8/4209 (0.2%)
1	B	0.97	2/3083 (0.1%)	0.96	5/4201 (0.1%)
All	All	0.96	2/6171 (0.0%)	0.95	13/8410 (0.2%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	279	ASN	CG-OD1	5.31	1.35	1.24
1	B	85	PHE	CG-CD1	5.01	1.46	1.38

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	122	ARG	NE-CZ-NH2	-15.09	112.75	120.30
1	A	234	MET	CG-SD-CE	-9.72	84.66	100.20
1	A	122	ARG	NE-CZ-NH1	9.50	125.05	120.30
1	B	122	ARG	NE-CZ-NH1	8.94	124.77	120.30
1	A	356	TYR	CB-CG-CD2	-8.41	115.95	121.00
1	A	122	ARG	NE-CZ-NH2	-7.89	116.36	120.30
1	B	116	TYR	CB-CG-CD2	-7.07	116.76	121.00
1	A	356	TYR	CB-CG-CD1	7.04	125.22	121.00
1	B	392	ARG	CG-CD-NE	-6.43	98.31	111.80
1	A	350	TYR	CB-CG-CD2	-6.40	117.16	121.00
1	A	350	TYR	CB-CG-CD1	6.38	124.83	121.00
1	A	234	MET	CB-CG-SD	6.29	131.28	112.40
1	B	96	ASP	CB-CG-OD1	5.77	123.49	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2956	0	2841	30	0
1	B	2947	0	2850	22	0
2	A	2	0	0	0	0
2	B	3	0	0	0	0
3	A	62	0	26	9	0
3	B	31	0	13	1	0
4	A	12	0	13	3	0
4	B	12	0	13	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	278	0	0	10	1
6	B	276	0	0	3	0
All	All	6581	0	5756	58	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (58) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:327[B]:ASN:ND2	1:A:328[B]:GLN:NE2	1.74	1.32
1:A:327[B]:ASN:HD22	1:A:328[B]:GLN:NE2	1.26	1.32
3:A:1393[B]:ANP:C5'	6:A:2125:HOH:O	1.87	1.22
3:A:1393[B]:ANP:H4'	6:A:2125:HOH:O	1.44	1.15
3:A:1393[B]:ANP:C4'	6:A:2125:HOH:O	1.97	0.99
1:B:342:PRO:HB3	6:B:2251:HOH:O	1.64	0.96
4:A:1395:MES:H82	6:A:2275:HOH:O	1.69	0.93
1:B:323[B]:GLN:OE1	1:B:327:ASN:ND2	2.02	0.92
1:A:327[B]:ASN:ND2	1:A:328[B]:GLN:HE22	1.55	0.92
1:A:327[B]:ASN:ND2	1:A:328[B]:GLN:HE21	1.47	0.88
3:A:1393[B]:ANP:H5'1	6:A:2125:HOH:O	1.63	0.86
1:A:158:ASP:OD2	6:A:2104:HOH:O	1.93	0.86
1:A:346[B]:TYR:CD2	3:A:1393[B]:ANP:C6	2.61	0.82
1:A:45:ILE:HG22	1:A:112:PRO:HG2	1.65	0.79
1:B:266[A]:GLU:OE1	6:B:2185:HOH:O	2.01	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:152:LYS:HD3	1:B:394:LEU:CD1	2.18	0.73
1:A:346[B]:TYR:CD2	3:A:1393[B]:ANP:N6	2.59	0.70
1:B:323[B]:GLN:CD	1:B:327:ASN:HD22	1.96	0.69
1:A:327[B]:ASN:HD21	1:A:328[B]:GLN:HE22	1.38	0.68
1:A:327[B]:ASN:HD22	1:A:328[B]:GLN:HE21	0.70	0.68
1:A:37:ASN:HB3	1:A:40:GLU:HG3	1.78	0.66
1:A:165:LEU:HD21	1:A:212:LEU:HD11	1.79	0.65
1:B:152:LYS:HD3	1:B:394:LEU:HD11	1.79	0.64
1:B:307[A]:GLN:HG3	1:B:308:LEU:HD22	1.81	0.62
1:B:109:HIS:HB2	1:B:145:GLN:O	2.00	0.61
1:A:66:ASP:OD1	1:A:66:ASP:N	2.33	0.61
1:A:302:ASN:HB3	3:A:1393[A]:ANP:C2	2.35	0.56
1:A:323[A]:GLN:OE1	1:A:327[A]:ASN:ND2	2.36	0.55
1:A:346[B]:TYR:CE2	3:A:1393[B]:ANP:N6	2.77	0.52
1:B:350:TYR:OH	3:B:1396:ANP:H2	2.10	0.51
1:B:78:ASN:HD22	1:B:78:ASN:C	2.14	0.51
1:B:125:PRO:HD2	1:B:128[B]:GLN:HE21	1.75	0.51
1:B:303:LEU:HD23	1:B:303:LEU:C	2.32	0.50
1:B:161:LEU:HD22	1:B:210[A]:VAL:HG11	1.93	0.49
1:B:78:ASN:C	1:B:78:ASN:ND2	2.66	0.49
4:A:1395:MES:H31	6:A:2083:HOH:O	2.13	0.49
1:A:201:ASN:HB3	1:A:204:ILE:HD12	1.95	0.48
1:A:213:ASN:HA	1:A:217:GLN:O	2.13	0.48
1:B:177:VAL:HA	1:B:217:GLN:OE1	2.14	0.48
1:A:231:GLN:NE2	3:A:1393[A]:ANP:O2'	2.47	0.47
1:A:206:LYS:HG2	6:A:2136:HOH:O	2.14	0.47
1:B:57:LEU:O	1:B:78:ASN:HA	2.16	0.46
1:A:126:GLN:HG2	6:A:2075:HOH:O	2.15	0.46
1:B:50:ALA:HB3	1:B:117:ALA:HA	1.98	0.45
1:A:276:HIS:ND1	4:A:1395:MES:O3S	2.51	0.44
1:A:323[B]:GLN:NE2	6:A:2249:HOH:O	2.42	0.44
1:A:39:CYS:HB2	1:A:108:ILE:HD13	2.00	0.44
1:A:50:ALA:HB3	1:A:117:ALA:HA	2.01	0.42
1:A:63:ASP:O	1:A:70:PRO:HA	2.19	0.42
1:B:165:LEU:HD21	1:B:212:LEU:HD11	2.01	0.42
1:B:37:ASN:HA	1:B:38:PRO:HD3	1.88	0.41
1:B:170:LYS:HD2	1:B:170:LYS:HA	1.92	0.41
1:A:324:GLN:O	1:A:328[A]:GLN:HB2	2.20	0.41
1:A:161:LEU:HD13	1:A:210[A]:VAL:HG11	2.02	0.41
1:B:266[A]:GLU:CD	6:B:2185:HOH:O	2.55	0.41
1:B:308:LEU:HD13	1:B:308:LEU:HA	1.93	0.40

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:229:LEU:HD13	1:A:272[B]:MET:HE2	2.03	0.40
1:A:296[B]:SER:OG	1:A:300:ILE:HG21	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2062:HOH:O	6:A:2252:HOH:O[2_755]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	375/368 (102%)	362 (96%)	13 (4%)	0	100	100
1	B	376/368 (102%)	366 (97%)	10 (3%)	0	100	100
All	All	751/736 (102%)	728 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	338/328 (103%)	330 (98%)	8 (2%)	57	22

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	338/328 (103%)	335 (99%)	3 (1%)	84	65
All	All	676/656 (103%)	665 (98%)	11 (2%)	70	41

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	ARG
1	A	66	ASP
1	A	180	LYS
1	A	234	MET
1	A	280	GLN
1	A	328[A]	GLN
1	A	328[B]	GLN
1	A	349	GLN
1	B	56	ARG
1	B	78	ASN
1	B	265	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	142	GLN
1	A	145	GLN
1	A	231	GLN
1	A	302	ASN
1	A	349	GLN
1	B	78	ASN
1	B	109	HIS
1	B	145	GLN
1	B	218	ASN
1	B	220	ASN
1	B	231	GLN
1	B	327	ASN
1	B	349	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry ⓘ

Of 12 ligands modelled in this entry, 7 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	ANP	A	1393[A]	2	27,33,33	2.13	8 (29%)	30,52,52	2.26	4 (13%)
3	ANP	A	1393[B]	2	27,33,33	2.11	8 (29%)	30,52,52	3.07	11 (36%)
4	MES	A	1395	-	11,12,12	1.35	2 (18%)	14,16,16	2.64	7 (50%)
3	ANP	B	1396	2	27,33,33	2.03	5 (18%)	30,52,52	3.20	10 (33%)
4	MES	B	1398	-	11,12,12	1.17	2 (18%)	14,16,16	3.50	4 (28%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ANP	A	1393[A]	2	-	1/12/38/38	0/3/3/3
3	ANP	A	1393[B]	2	-	0/12/38/38	0/3/3/3
4	MES	A	1395	-	-	0/6/14/14	0/1/1/1
3	ANP	B	1396	2	-	1/12/38/38	0/3/3/3
4	MES	B	1398	-	-	0/6/14/14	0/1/1/1

All (25) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1393[B]	ANP	PB-O2B	-2.77	1.49	1.56
3	A	1393[A]	ANP	PB-O2B	-2.75	1.49	1.56
3	B	1396	ANP	PG-O3G	-2.55	1.49	1.56
3	A	1393[B]	ANP	PG-O2G	-2.45	1.49	1.56
3	A	1393[B]	ANP	PG-O3G	-2.44	1.49	1.56
3	A	1393[A]	ANP	PG-O2G	-2.43	1.49	1.56
3	A	1393[A]	ANP	PG-O3G	-2.35	1.50	1.56
4	B	1398	MES	O2S-S	2.09	1.51	1.45
4	A	1395	MES	O1S-S	2.14	1.52	1.45
3	A	1393[A]	ANP	PB-O3A	2.16	1.61	1.59
3	A	1393[B]	ANP	PG-N3B	2.18	1.69	1.63
4	B	1398	MES	C5-N4	2.37	1.53	1.46
3	A	1393[B]	ANP	PB-O3A	2.44	1.62	1.59
3	B	1396	ANP	PG-N3B	2.51	1.70	1.63
3	B	1396	ANP	C4-N3	2.61	1.39	1.35
3	A	1393[A]	ANP	PG-N3B	2.62	1.70	1.63
3	B	1396	ANP	PB-O1B	2.76	1.49	1.46
4	A	1395	MES	O2S-S	3.02	1.54	1.45
3	A	1393[B]	ANP	PB-N3B	3.96	1.73	1.63
3	A	1393[A]	ANP	PB-N3B	4.00	1.73	1.63
3	A	1393[B]	ANP	PB-O1B	4.35	1.51	1.46
3	A	1393[A]	ANP	PB-O1B	4.97	1.51	1.46
3	A	1393[A]	ANP	PG-O1G	6.33	1.53	1.46
3	A	1393[B]	ANP	PG-O1G	6.40	1.53	1.46
3	B	1396	ANP	PG-O1G	8.06	1.55	1.46

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1393[B]	ANP	N3-C2-N1	-13.14	118.83	128.89
3	B	1396	ANP	N3-C2-N1	-12.42	119.38	128.89
3	A	1393[A]	ANP	N3-C2-N1	-9.08	121.94	128.89
3	B	1396	ANP	C2'-C1'-N9	-8.43	101.42	114.29
3	A	1393[A]	ANP	C2'-C1'-N9	-5.79	105.45	114.29
4	B	1398	MES	O3S-S-O1S	-5.54	98.70	111.61
4	A	1395	MES	O2S-S-C8	-4.88	102.74	106.91
3	B	1396	ANP	O1G-PG-N3B	-4.34	105.24	111.90
3	A	1393[B]	ANP	O1G-PG-N3B	-4.02	105.73	111.90
3	A	1393[B]	ANP	O3A-PB-N3B	-3.98	95.48	106.44
3	B	1396	ANP	C1'-N9-C4	-3.42	121.78	126.94
4	A	1395	MES	C2-C3-N4	-3.23	105.23	110.12
3	A	1393[B]	ANP	C1'-N9-C4	-3.18	122.15	126.94
4	A	1395	MES	C6-C5-N4	-2.72	106.00	110.12

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1396	ANP	O3A-PB-N3B	-2.51	99.53	106.44
3	A	1393[A]	ANP	PA-O3A-PB	-2.39	124.66	132.67
3	B	1396	ANP	O1B-PB-N3B	-2.25	108.45	111.90
3	A	1393[B]	ANP	C4'-O4'-C1'	-2.24	107.26	109.72
3	A	1393[B]	ANP	O3A-PA-O5'	-2.19	97.13	102.94
3	B	1396	ANP	O2'-C2'-C3'	-2.01	105.30	111.83
3	A	1393[B]	ANP	C2-N1-C6	2.05	122.43	118.77
4	A	1395	MES	O1-C6-C5	2.12	116.70	111.84
3	B	1396	ANP	O4'-C1'-N9	2.13	112.56	108.10
3	B	1396	ANP	O3G-PG-O2G	2.40	114.69	107.58
3	A	1393[B]	ANP	O3G-PG-O2G	2.45	114.83	107.58
4	B	1398	MES	C6-O1-C2	2.46	118.19	109.89
3	A	1393[A]	ANP	O2B-PB-O1B	2.47	115.15	110.00
3	A	1393[B]	ANP	O1B-PB-N3B	2.50	115.73	111.90
4	A	1395	MES	C5-N4-C3	2.69	114.73	108.90
4	B	1398	MES	C5-N4-C3	3.17	115.76	108.90
4	A	1395	MES	O1S-S-C8	3.17	109.61	106.91
3	A	1393[B]	ANP	C2'-C1'-N9	3.28	119.31	114.29
3	B	1396	ANP	O2B-PB-O1B	3.36	117.02	110.00
3	A	1393[B]	ANP	O2B-PB-O1B	4.43	119.25	110.00
4	A	1395	MES	C6-O1-C2	4.70	125.71	109.89
4	B	1398	MES	O2S-S-C8	10.50	115.86	106.91

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	1396	ANP	O1G-PG-N3B-PB
3	A	1393[A]	ANP	O1G-PG-N3B-PB

There are no ring outliers.

4 monomers are involved in 13 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1393[A]	ANP	2	0
3	A	1393[B]	ANP	7	0
4	A	1395	MES	3	0
3	B	1396	ANP	1	0



## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	359/368 (97%)	0.13	21 (5%) 26 28	10, 19, 44, 73	0
1	B	358/368 (97%)	0.09	13 (3%) 46 49	9, 17, 41, 60	0
All	All	717/736 (97%)	0.11	34 (4%) 35 36	9, 18, 44, 73	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	179	ASN	6.1
1	A	177	VAL	6.1
1	A	381	ASN	5.7
1	B	177	VAL	4.9
1	A	378	PRO	4.7
1	B	393	ALA	4.4
1	A	36	THR	4.3
1	A	176	SER	3.8
1	A	213	ASN	3.6
1	A	67	THR	3.5
1	B	66	ASP	3.4
1	B	381	ASN	3.4
1	A	216	GLY	3.4
1	B	179	ASN	3.3
1	B	394	LEU	3.2
1	B	37	ASN	3.2
1	A	66	ASP	3.2
1	B	343	ASP	3.0
1	A	218	ASN	3.0
1	B	377	PRO	2.9
1	A	180	LYS	2.8
1	A	379	GLU	2.6
1	A	35	ASP	2.6
1	B	175	LYS	2.6

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	138	TYR	2.6
1	A	343	ASP	2.5
1	A	217	GLN	2.5
1	B	203	GLU	2.4
1	B	378	PRO	2.3
1	A	34	MET	2.2
1	A	377	PRO	2.2
1	B	379	GLU	2.1
1	A	175	LYS	2.1
1	A	104	ALA	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	MG	B	506	1/1	0.98	0.29	13.48	44,44,44,44	0
4	MES	A	1395	12/12	0.92	0.10	1.38	29,36,48,49	0
4	MES	B	1398	12/12	0.97	0.09	0.97	21,29,36,39	0
2	MG	A	1394	1/1	1.00	0.10	0.77	12,12,12,12	0
3	ANP	A	1393[A]	31/31	0.98	0.10	0.66	10,20,43,45	31
3	ANP	A	1393[B]	31/31	0.98	0.10	0.46	10,22,38,42	31
5	CL	A	1396	1/1	1.00	0.09	0.19	20,20,20,20	0
3	ANP	B	1396	31/31	0.99	0.08	-0.43	9,23,39,50	0
5	CL	B	1395	1/1	1.00	0.07	-0.49	25,25,25,25	0
2	MG	B	1397	1/1	1.00	0.08	-0.96	12,12,12,12	0
2	MG	B	505	1/1	0.99	0.14	-	34,34,34,34	0
2	MG	A	505	1/1	1.00	0.04	-	16,16,16,16	0

## 6.5 Other polymers

There are no such residues in this entry.