



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:11 PM GMT

PDB ID : 4BSV  
Title : Heterodimeric Fc Antibody Azymetric Variant 1  
Authors : Suits, M.D.L.; Spreter, T.; Cabrera, E.E.; Dixit, S.B.; Lario, P.I.; Poon, D.K.Y.; D'Angelo, I.E.P.; Boulanger, M.J.  
Deposited on : 2013-06-11  
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	232	
2	B	232	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	1448[B]	-	-	X	-
4	GAL	A	1452[A]	-	-	-	X
4	GAL	A	1452[B]	-	-	-	X
6	EDO	A	1455	-	-	-	X
6	EDO	A	1456[B]	-	-	X	X
6	EDO	A	1461	-	-	X	X
6	EDO	A	1462	-	-	X	-
6	EDO	B	1455	-	-	X	X
6	EDO	B	1456[A]	-	-	X	-
6	EDO	B	1456[B]	-	-	X	X
6	EDO	B	1457	-	-	-	X
6	EDO	B	1458[A]	-	-	X	X
6	EDO	B	1458[B]	-	-	X	X
6	EDO	B	1459	-	-	X	X
7	NAG	B	1448[B]	-	-	X	-
7	GAL	B	1452[A]	-	-	-	X
7	GAL	B	1452[B]	-	-	-	X

## 2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 7945 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HETERODIMERIC FC ANTIBODY AZYMETRIC VARIANT 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	210	Total	C	N	O	S	0	210	0
			3350	2135	564	638	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	222	GLY	-	EXPRESSION TAG	UNP P01857
A	350	VAL	THR	ENGINEERED MUTATION	UNP P01857
A	351	TYR	LEU	MICROHETEROGENEITY	UNP P01857
A	366	LEU	THR	MICROHETEROGENEITY	UNP P01857
A	392	MET	LYS	MICROHETEROGENEITY	UNP P01857
A	394	TRP	THR	MICROHETEROGENEITY	UNP P01857
A	405	ALA	PHE	MICROHETEROGENEITY	UNP P01857
A	407	VAL	TYR	MICROHETEROGENEITY	UNP P01857

- Molecule 2 is a protein called HETERODIMERIC FC ANTIBODY AZYMETRIC VARIANT 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	210	Total	C	N	O	S	0	210	0
			3350	2135	564	638	13			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	222	GLY	-	EXPRESSION TAG	UNP P01857
B	350	VAL	THR	ENGINEERED MUTATION	UNP P01857
B	351	TYR	LEU	MICROHETEROGENEITY	UNP P01857
B	366	LEU	THR	MICROHETEROGENEITY	UNP P01857
B	392	MET	LYS	MICROHETEROGENEITY	UNP P01857
B	394	THR	TRP	MICROHETEROGENEITY	UNP P01857

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	405	ALA	PHE	MICROHETEROGENEITY	UNP P01857
B	407	VAL	TYR	MICROHETEROGENEITY	UNP P01857

- Molecule 3 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	2	Total	C	N	O	0	2
			48	28	2	18		


- Molecule 4 is a polymer of unknown type called SUGAR (5-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	5	Total	C	N	O	0	5
			122	68	4	50		

- Molecule 5 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	2	Total	C	N	O	0	2
			50	28	2	20		

- Molecule 6 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			4	2	2		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	1
			8	4	4		
6	A	1	Total	C	O	0	0
			4	2	2		
6	A	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	1
			8	4	4		
6	B	1	Total	C	O	0	0
			4	2	2		
6	B	1	Total	C	O	0	1
			8	4	4		
6	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 7 is a polymer of unknown type called SUGAR (9-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
7	B	9	Total	C	N	O	0	9
			220	124	8	88		

- Molecule 8 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	B	3	Total	I	0	1
			4	4		


- Molecule 9 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	342	Total	O	0	0
			342	342		
9	B	411	Total	O	0	0
			411	411		


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: HETERODIMERIC FC ANTIBODY AZYMETRIC VARIANT 2



#### • Molecule 2: HETERODIMERIC FC ANTIBODY AZYMETRIC VARIANT 2



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	49.54Å 74.92Å 148.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	41.38 – 1.75 41.38 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.1 (41.38-1.75) 99.0 (41.38-1.75)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	9.74 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.178 , 0.208 0.181 , 0.212	Depositor DCC
$R_{free}$ test set	2849 reflections (5.33%)	DCC
Wilson B-factor (Å <sup>2</sup> )	23.6	Xtriage
Anisotropy	0.055	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 58.4	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 56316 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	7945	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.51% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BMA, NAG, EDO, GAL, FUC, IOD, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.54	3/3439 (0.1%)	0.60	0/4677
2	B	0.52	0/3439	0.72	5/4677 (0.1%)
All	All	0.53	3/6878 (0.0%)	0.66	5/9354 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	4
2	B	0	4
All	All	0	8

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	270[A]	ASP	C-N	-11.34	1.12	1.34
1	A	270[B]	ASP	C-N	-11.34	1.12	1.34
1	A	405[B]	PHE	C-N	-8.19	1.15	1.34

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	405[B]	ALA	O-C-N	8.36	136.07	122.70
2	B	405[B]	ALA	C-N-CA	-6.84	104.61	121.70
2	B	405[B]	ALA	CA-C-N	-6.76	102.32	117.20
2	B	252[A]	MET	CG-SD-CE	-5.29	91.74	100.20
2	B	252[B]	MET	CG-SD-CE	-5.29	91.74	100.20

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	373[A]	TYR	Mainchain,Peptide
1	A	373[B]	TYR	Mainchain,Peptide
2	B	373[A]	TYR	Mainchain,Peptide
2	B	373[B]	TYR	Mainchain,Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3350	0	3269	197	0
2	B	3350	0	3273	182	0
3	A	48	0	43	3	0
4	A	122	0	104	32	0
5	A	50	0	44	12	0
6	A	20	0	27	20	0
6	B	28	0	38	33	0
7	B	220	0	189	31	0
8	B	4	0	0	0	0
9	A	342	0	0	78	0
9	B	411	0	0	83	0
All	All	7945	0	6987	438	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 33.

The worst 5 of 438 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:397[A]:VAL:CG1	9:A:4123:HOH:O	1.64	1.34
6:B:1458[A]:EDO:C2	9:B:2153:HOH:O	1.78	1.31
7:B:1454[B]:NAG:O3	7:B:1454[B]:NAG:H82	1.24	1.30
5:A:1454[A]:NAG:O3	5:A:1454[A]:NAG:H82	1.16	1.28
2:B:297[B]:ASN:HD21	7:B:1446[B]:NAG:C1	1.50	1.24

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	404/232 (174%)	390 (96%)	12 (3%)	2 (0%)	34	14
2	B	404/232 (174%)	386 (96%)	14 (4%)	4 (1%)	19	5
All	All	808/464 (174%)	776 (96%)	26 (3%)	6 (1%)	26	10

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	374[A]	PRO
1	A	374[B]	PRO
2	B	374[A]	PRO
2	B	374[B]	PRO
2	B	237[A]	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	389/213 (183%)	387 (100%)	2 (0%)	92	87
2	B	389/213 (183%)	380 (98%)	9 (2%)	58	33
All	All	778/426 (183%)	767 (99%)	11 (1%)	72	58

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	351[A]	LEU
2	B	398[A]	LEU
2	B	418[B]	GLN
2	B	308[B]	VAL
2	B	418[A]	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

36 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	NAG	A	1446[A]	1,3	14,14,15	0.60	0	15,19,21	1.09	2 (13%)
3	NAG	A	1446[B]	1,3	14,14,15	0.72	0	15,19,21	0.88	0
3	FUC	A	1447[A]	3	10,10,11	0.66	0	14,14,16	2.02	7 (50%)
3	FUC	A	1447[B]	3	10,10,11	0.67	0	14,14,16	2.09	7 (50%)
4	NAG	A	1448[A]	4	14,14,15	0.63	0	15,19,21	0.79	0
4	NAG	A	1448[B]	4	14,14,15	0.72	0	15,19,21	2.04	3 (20%)
4	BMA	A	1449[A]	4	11,11,12	0.32	0	14,15,17	1.29	1 (7%)
4	BMA	A	1449[B]	4	11,11,12	0.86	0	14,15,17	1.01	1 (7%)
4	MAN	A	1450[A]	4	11,11,12	0.66	0	14,15,17	1.34	2 (14%)
4	MAN	A	1450[B]	4	11,11,12	0.70	0	14,15,17	1.02	1 (7%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
4	NAG	A	1451[A]	4	14,14,15	0.68	0	15,19,21	0.75	0
4	NAG	A	1451[B]	4	14,14,15	0.71	0	15,19,21	0.68	0
4	GAL	A	1452[A]	4	11,11,12	0.59	0	14,15,17	0.51	0
4	GAL	A	1452[B]	4	11,11,12	0.58	0	14,15,17	1.50	2 (14%)
5	MAN	A	1453[A]	5	11,11,12	0.69	0	14,15,17	0.80	0
5	MAN	A	1453[B]	5	11,11,12	0.59	0	14,15,17	1.95	3 (21%)
5	NAG	A	1454[A]	5	14,14,15	0.50	0	15,19,21	0.77	0
5	NAG	A	1454[B]	5	14,14,15	0.50	0	15,19,21	0.76	0
7	NAG	B	1446[A]	2,7	14,14,15	0.66	0	15,19,21	1.09	0
7	NAG	B	1446[B]	7	14,14,15	0.36	0	15,19,21	0.83	0
7	FUC	B	1447[A]	7	10,10,11	0.34	0	14,14,16	1.09	2 (14%)
7	FUC	B	1447[B]	-	10,10,11	0.48	0	14,14,16	0.96	0
7	NAG	B	1448[A]	7	14,14,15	0.71	0	15,19,21	0.90	0
7	NAG	B	1448[B]	7	14,14,15	0.55	0	15,19,21	0.84	0
7	BMA	B	1449[A]	7	11,11,12	0.61	0	14,15,17	1.28	2 (14%)
7	BMA	B	1449[B]	7	11,11,12	0.39	0	14,15,17	1.13	1 (7%)
7	MAN	B	1450[A]	7	11,11,12	0.67	0	14,15,17	1.20	1 (7%)
7	MAN	B	1450[B]	7	11,11,12	0.64	0	14,15,17	1.32	3 (21%)
7	NAG	B	1451[A]	7	14,14,15	0.60	0	15,19,21	1.09	0
7	NAG	B	1451[B]	7	14,14,15	0.52	0	15,19,21	0.84	0
7	GAL	B	1452[A]	7	11,11,12	0.51	0	14,15,17	1.35	3 (21%)
7	GAL	B	1452[B]	7	11,11,12	0.50	0	14,15,17	1.38	3 (21%)
7	MAN	B	1453[A]	7	11,11,12	0.64	0	14,15,17	0.64	0
7	MAN	B	1453[B]	7	11,11,12	0.64	0	14,15,17	0.76	0
7	NAG	B	1454[A]	7	14,14,15	0.52	0	15,19,21	0.60	0
7	NAG	B	1454[B]	7	14,14,15	0.52	0	15,19,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	A	1446[A]	1,3	-	0/6/23/26	0/1/1/1
3	NAG	A	1446[B]	1,3	-	0/6/23/26	0/1/1/1
3	FUC	A	1447[A]	3	-	0/0/17/20	0/1/1/1
3	FUC	A	1447[B]	3	-	0/0/17/20	0/1/1/1
4	NAG	A	1448[A]	4	-	0/6/23/26	0/1/1/1
4	NAG	A	1448[B]	4	-	0/6/23/26	0/1/1/1
4	BMA	A	1449[A]	4	-	0/2/19/22	0/1/1/1
4	BMA	A	1449[B]	4	-	0/2/19/22	0/1/1/1

Continued on next page...

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	MAN	A	1450[A]	4	-	0/2/19/22	0/1/1/1
4	MAN	A	1450[B]	4	-	0/2/19/22	0/1/1/1
4	NAG	A	1451[A]	4	-	0/6/23/26	0/1/1/1
4	NAG	A	1451[B]	4	-	0/6/23/26	0/1/1/1
4	GAL	A	1452[A]	4	-	0/2/19/22	0/1/1/1
4	GAL	A	1452[B]	4	-	0/2/19/22	0/1/1/1
5	MAN	A	1453[A]	5	-	0/2/19/22	0/1/1/1
5	MAN	A	1453[B]	5	-	0/2/19/22	0/1/1/1
5	NAG	A	1454[A]	5	-	0/6/23/26	0/1/1/1
5	NAG	A	1454[B]	5	-	0/6/23/26	0/1/1/1
7	NAG	B	1446[A]	2,7	-	0/6/23/26	0/1/1/1
7	NAG	B	1446[B]	7	-	0/6/23/26	0/1/1/1
7	FUC	B	1447[A]	7	-	0/0/17/20	0/1/1/1
7	FUC	B	1447[B]	-	-	0/0/17/20	0/1/1/1
7	NAG	B	1448[A]	7	-	0/6/23/26	0/1/1/1
7	NAG	B	1448[B]	7	-	0/6/23/26	0/1/1/1
7	BMA	B	1449[A]	7	-	0/2/19/22	0/1/1/1
7	BMA	B	1449[B]	7	-	0/2/19/22	0/1/1/1
7	MAN	B	1450[A]	7	-	0/2/19/22	0/1/1/1
7	MAN	B	1450[B]	7	-	0/2/19/22	0/1/1/1
7	NAG	B	1451[A]	7	-	0/6/23/26	0/1/1/1
7	NAG	B	1451[B]	7	-	0/6/23/26	0/1/1/1
7	GAL	B	1452[A]	7	-	0/2/19/22	0/1/1/1
7	GAL	B	1452[B]	7	-	0/2/19/22	0/1/1/1
7	MAN	B	1453[A]	7	-	0/2/19/22	0/1/1/1
7	MAN	B	1453[B]	7	-	0/2/19/22	0/1/1/1
7	NAG	B	1454[A]	7	-	0/6/23/26	0/1/1/1
7	NAG	B	1454[B]	7	-	0/6/23/26	0/1/1/1

There are no bond length outliers.

The worst 5 of 44 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	1448[B]	NAG	C1-O5-C5	-5.87	104.80	112.25
4	A	1452[B]	GAL	C3-C4-C5	-4.31	102.68	110.20
3	A	1447[B]	FUC	C1-O5-C5	-3.53	106.93	112.38
3	A	1447[A]	FUC	C1-O5-C5	-3.39	107.14	112.38
7	B	1450[B]	MAN	O5-C1-C2	-2.73	106.43	110.86

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

21 monomers are involved in 68 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1446[B]	NAG	3	0
4	A	1448[A]	NAG	6	0
4	A	1448[B]	NAG	8	0
4	A	1449[A]	BMA	3	0
4	A	1449[B]	BMA	4	0
4	A	1451[A]	NAG	2	0
4	A	1451[B]	NAG	5	0
4	A	1452[A]	GAL	2	0
4	A	1452[B]	GAL	2	0
5	A	1453[A]	MAN	4	0
5	A	1453[B]	MAN	4	0
5	A	1454[A]	NAG	5	0
7	B	1446[B]	NAG	5	0
7	B	1447[A]	FUC	1	0
7	B	1447[B]	FUC	2	0
7	B	1448[A]	NAG	5	0
7	B	1448[B]	NAG	11	0
7	B	1449[B]	BMA	3	0
7	B	1452[B]	GAL	1	0
7	B	1454[A]	NAG	3	0
7	B	1454[B]	NAG	3	0

## 5.6 Ligand geometry

Of 16 ligands modelled in this entry, 4 are monoatomic - leaving 12 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	EDO	A	1455	-	3,3,3	0.46	0	2,2,2	0.47	0
6	EDO	A	1456[A]	-	3,3,3	0.49	0	2,2,2	0.40	0
6	EDO	A	1456[B]	-	3,3,3	0.50	0	2,2,2	0.43	0
6	EDO	A	1461	-	3,3,3	0.51	0	2,2,2	0.27	0
6	EDO	A	1462	-	3,3,3	0.49	0	2,2,2	0.42	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	EDO	B	1455	-	3,3,3	0.50	0	2,2,2	0.43	0
6	EDO	B	1456[A]	-	3,3,3	0.52	0	2,2,2	0.71	0
6	EDO	B	1456[B]	-	3,3,3	0.52	0	2,2,2	0.36	0
6	EDO	B	1457	-	3,3,3	0.49	0	2,2,2	0.42	0
6	EDO	B	1458[A]	-	3,3,3	0.62	0	2,2,2	0.18	0
6	EDO	B	1458[B]	-	3,3,3	0.58	0	2,2,2	0.45	0
6	EDO	B	1459	-	3,3,3	0.50	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	EDO	A	1455	-	-	0/1/1/1	0/0/0/0
6	EDO	A	1456[A]	-	-	0/1/1/1	0/0/0/0
6	EDO	A	1456[B]	-	-	0/1/1/1	0/0/0/0
6	EDO	A	1461	-	-	0/1/1/1	0/0/0/0
6	EDO	A	1462	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1455	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1456[A]	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1456[B]	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1457	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1458[A]	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1458[B]	-	-	0/1/1/1	0/0/0/0
6	EDO	B	1459	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

12 monomers are involved in 52 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1455	EDO	3	0
6	A	1456[A]	EDO	3	0
6	A	1456[B]	EDO	5	0
6	A	1461	EDO	5	0
6	A	1462	EDO	4	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1455	EDO	5	0
6	B	1456[A]	EDO	4	0
6	B	1456[B]	EDO	10	0
6	B	1457	EDO	1	0
6	B	1458[A]	EDO	5	0
6	B	1458[B]	EDO	4	0
6	B	1459	EDO	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	216/232 (93%)	0.79	26 (12%) 6 8	14, 23, 43, 60	12 (5%)
2	B	216/232 (93%)	0.58	10 (4%) 36 42	12, 21, 34, 43	12 (5%)
All	All	432/464 (93%)	0.68	36 (8%) 14 17	12, 22, 38, 60	24 (5%)

The worst 5 of 36 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	300[A]	TYR	6.1
1	A	296[A]	TYR	5.5
1	A	445[A]	PRO	4.8
1	A	444[A]	SER	4.4
2	B	236[A]	GLY	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	GAL	A	1452[B]	11/12	0.77	0.26	16.43	23,25,26,26	11
4	GAL	A	1452[A]	11/12	0.77	0.26	14.15	92,100,103,104	11
7	GAL	B	1452[A]	11/12	0.79	0.21	2.57	230,236,238,239	11

*Continued on next page...*

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
7	GAL	B	1452[B]	11/12	0.79	0.21	2.57	22,26,29,29	11
7	FUC	B	1447[A]	10/11	0.87	0.18	0.71	52,54,56,58	10
7	FUC	B	1447[B]	10/11	0.87	0.18	0.71	25,26,26,27	10
7	NAG	B	1451[A]	14/15	0.94	0.14	0.57	34,38,43,43	14
7	NAG	B	1451[B]	14/15	0.94	0.14	0.57	20,22,29,31	14
4	NAG	A	1448[B]	14/15	0.92	0.13	0.45	23,25,30,31	14
3	FUC	A	1447[A]	10/11	0.76	0.22	0.24	65,68,71,72	10
3	FUC	A	1447[B]	10/11	0.76	0.22	0.24	38,42,43,44	10
4	NAG	A	1451[A]	14/15	0.90	0.11	0.18	22,23,26,27	14
4	NAG	A	1448[A]	14/15	0.92	0.13	0.12	25,26,29,30	14
4	NAG	A	1451[B]	14/15	0.90	0.11	-0.20	39,41,43,44	14
7	NAG	B	1448[B]	14/15	0.94	0.10	-0.89	30,33,35,38	14
7	NAG	B	1448[A]	14/15	0.94	0.10	-0.89	24,27,28,29	14
3	NAG	A	1446[A]	14/15	0.89	0.13	-1.15	24,27,29,30	14
3	NAG	A	1446[B]	14/15	0.89	0.13	-1.15	27,29,30,30	14
7	NAG	B	1446[B]	14/15	0.93	0.11	-1.27	29,30,37,37	14
7	NAG	B	1446[A]	14/15	0.93	0.11	-1.27	32,34,37,37	14
7	MAN	B	1450[A]	11/12	0.91	0.14	-	29,31,33,35	11
5	NAG	A	1454[B]	14/15	0.70	0.21	-	111,116,118,119	14
5	MAN	A	1453[A]	11/12	0.85	0.13	-	34,35,36,37	11
7	MAN	B	1453[A]	11/12	0.84	0.13	-	106,109,111,115	11
5	NAG	A	1454[A]	14/15	0.70	0.21	-	100,104,106,106	14
4	BMA	A	1449[B]	11/12	0.93	0.10	-	33,35,37,38	11
7	MAN	B	1453[B]	11/12	0.84	0.13	-	40,42,45,46	11
4	BMA	A	1449[A]	11/12	0.93	0.10	-	20,21,22,24	11
7	BMA	B	1449[A]	11/12	0.91	0.11	-	41,43,43,45	11
7	BMA	B	1449[B]	11/12	0.91	0.11	-	22,26,27,28	11
4	MAN	A	1450[A]	11/12	0.96	0.10	-	20,21,25,25	11
7	NAG	B	1454[A]	14/15	0.67	0.23	-	109,111,113,113	14
4	MAN	A	1450[B]	11/12	0.96	0.10	-	33,34,38,40	11
7	NAG	B	1454[B]	14/15	0.67	0.23	-	112,116,119,120	14
5	MAN	A	1453[B]	11/12	0.85	0.13	-	133,135,137,138	11
7	MAN	B	1450[B]	11/12	0.91	0.14	-	29,29,31,32	11

## 6.4 Ligands ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
6	EDO	B	1455	4/4	0.66	0.33	14.90	33,35,37,37	4
6	EDO	A	1456[B]	4/4	0.89	0.20	14.77	28,30,31,31	4
6	EDO	A	1461	4/4	0.93	0.22	7.56	54,55,55,55	4
6	EDO	A	1455	4/4	0.94	0.22	6.67	25,26,28,32	4
6	EDO	B	1459	4/4	0.93	0.20	5.66	31,34,34,34	4
6	EDO	B	1458[A]	4/4	0.89	0.22	5.60	21,23,26,27	4
6	EDO	B	1458[B]	4/4	0.89	0.22	5.10	27,29,31,35	4
6	EDO	B	1457	4/4	0.94	0.17	2.97	41,43,44,47	4
6	EDO	B	1456[B]	4/4	0.93	0.15	2.01	23,24,26,29	4
6	EDO	B	1456[A]	4/4	0.93	0.15	1.92	21,21,22,22	4
8	IOD	B	1460	1/1	1.00	0.13	1.15	18,18,18,18	1
8	IOD	B	1461	1/1	1.00	0.13	1.15	17,17,17,17	1
6	EDO	A	1462	4/4	0.88	0.11	-1.18	34,34,35,37	4
8	IOD	B	1462[B]	1/1	1.00	0.06	-3.44	23,23,23,23	1
8	IOD	B	1462[A]	1/1	1.00	0.06	-3.45	23,23,23,23	1
6	EDO	A	1456[A]	4/4	0.89	0.20	-	32,34,34,35	4

## 6.5 Other polymers ⓘ

There are no such residues in this entry.