



Full wwPDB X-ray Structure Validation Report ⓘ

Jan 31, 2016 – 06:40 PM GMT

PDB ID : 1BXO
Title : ACID PROTEINASE (PENICILLOPEPSIN) (E.C.3.4.23.20) COMPLEX WITH PHOSPHONATE INHIBITOR: METHYL CYCLO[(2S)-2-[(1R)-1-(N-(L-N-(3-METHYLBUTANOYL)VALYL-L-ASPARTYL)AMINO)-3-METHYLBUTYL] HYDROXYPHOSPHINYLOXY]-3-(3-AMINOMETHYL)PHENYLPROPANOATE
Authors : Khan, A.R.; Parrish, J.C.; Fraser, M.E.; Smith, W.W.; Bartlett, P.A.; James, M.N.G.
Deposited on : 1998-10-07
Resolution : 0.95 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 0.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	1031 (1.08-0.84)
Ramachandran outliers	100387	1331 (1.10-0.82)
Sidechain outliers	100360	1330 (1.10-0.82)
RSRZ outliers	91569	1324 (1.10-0.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	323	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MAN	A	329	-	-	-	X
3	SO4	A	327	-	-	-	X
5	GOL	A	325	-	-	-	X
5	GOL	A	326	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5462 atoms, of which 2365 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PROTEIN (PENICILLOPEPSIN).

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
			Total	C	H	N	O	S			
1	A	323	4784	1541	2313	388	539	3	0	31	0

- Molecule 2 is SUGAR (ALPHA-D-MANNOSE) (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
2	A	1	22	6	11	5	0	0
2	A	1	22	6	11	5	0	0

- Molecule 3 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is METHYL CYCLO[(2S)-2-[[[(1R)-1-(N-(L-N-(3-METHYLBUTANOYL) VALYL-L-ASPARTYL)AMINO)-3-METHYLBUTYL]HYDROXYPHOSPHINYLOXY]-3-(3-AMINOMETHYL)PHENYLPROPANOATE (three-letter code: PP7) (formula: C₃₀H₄₇N₄O₉P).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
4	A	1	Total	C	H	N	O	P	0	0
			74	30	30	4	9	1		

- Molecule 5 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			6	3	3		
5	A	1	Total	C	O	0	0
			6	3	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	528	Total	O	0	18
			543	543		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

● Molecule 1: PROTEIN (PENICILLOPEPSIN)



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	96.98Å 46.65Å 65.71Å 90.00° 115.57° 90.00°	Depositor
Resolution (Å)	10.00 – 0.95 10.00 – 0.90	Depositor EDS
% Data completeness (in resolution range)	98.0 (10.00-0.95) 78.4 (10.00-0.90)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.06	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.02 (at 0.90Å)	Xtriage
Refinement program	SHELXL-97	Depositor
R, R_{free}	0.100 , 0.125 0.108 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	10.8	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 65.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 156181 reflections	Xtriage
F_o, F_c correlation	0.99	EDS
Total number of atoms	5462	wwPDB-VP
Average B, all atoms (Å ²)	17.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.67% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PP7, SO4, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	2.84	73/2649 (2.8%)	1.89	68/3612 (1.9%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	5

All (73) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	279	ASP	C-O	80.30	2.75	1.23
1	A	278	GLY	C-O	50.62	2.04	1.23
1	A	279	ASP	N-CA	31.81	2.10	1.46
1	A	279	ASP	CG-OD2	25.20	1.83	1.25
1	A	60[A]	SER	CB-OG	-20.68	1.15	1.42
1	A	60[B]	SER	CB-OG	-20.68	1.15	1.42
1	A	279	ASP	CA-C	19.64	2.04	1.52
1	A	279	ASP	CG-OD1	17.48	1.65	1.25
1	A	280	GLY	C-O	16.86	1.50	1.23
1	A	16[A]	GLU	CD-OE1	14.76	1.41	1.25
1	A	16[B]	GLU	CD-OE1	14.76	1.41	1.25
1	A	16[A]	GLU	CD-OE2	14.66	1.41	1.25
1	A	16[B]	GLU	CD-OE2	14.66	1.41	1.25
1	A	67[A]	SER	CA-CB	13.95	1.73	1.52
1	A	67[B]	SER	CA-CB	13.95	1.73	1.52
1	A	234[A]	SER	CB-OG	-12.84	1.25	1.42
1	A	234[B]	SER	CB-OG	-12.84	1.25	1.42
1	A	280	GLY	N-CA	-12.68	1.27	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	98[A]	HIS	CB-CG	-12.29	1.27	1.50
1	A	98[B]	HIS	CB-CG	-12.29	1.27	1.50
1	A	281	SER	CA-CB	-12.26	1.34	1.52
1	A	98[A]	HIS	CG-ND1	11.76	1.64	1.38
1	A	98[B]	HIS	CG-ND1	11.76	1.64	1.38
1	A	279	ASP	CA-CB	11.47	1.79	1.53
1	A	52[A]	SER	CA-CB	-10.39	1.37	1.52
1	A	52[B]	SER	CA-CB	-10.39	1.37	1.52
1	A	145	LYS	CE-NZ	-10.20	1.23	1.49
1	A	74[A]	SER	CB-OG	-9.98	1.29	1.42
1	A	74[B]	SER	CB-OG	-9.98	1.29	1.42
1	A	278	GLY	CA-C	-9.96	1.35	1.51
1	A	98[A]	HIS	ND1-CE1	9.67	1.58	1.34
1	A	98[B]	HIS	ND1-CE1	9.67	1.58	1.34
1	A	206	ASP	CG-OD1	9.54	1.47	1.25
1	A	60[A]	SER	CA-CB	9.48	1.67	1.52
1	A	60[B]	SER	CA-CB	9.48	1.67	1.52
1	A	279	ASP	C-N	9.35	1.49	1.33
1	A	52[A]	SER	CB-OG	9.25	1.54	1.42
1	A	52[B]	SER	CB-OG	9.25	1.54	1.42
1	A	278	GLY	C-N	-8.94	1.13	1.34
1	A	277	SER	CA-CB	8.79	1.66	1.52
1	A	64	LYS	CE-NZ	8.51	1.70	1.49
1	A	172[A]	SER	CA-CB	8.34	1.65	1.52
1	A	172[B]	SER	CA-CB	8.34	1.65	1.52
1	A	234[A]	SER	CA-CB	8.24	1.65	1.52
1	A	234[B]	SER	CA-CB	8.24	1.65	1.52
1	A	176[A]	THR	CB-OG1	-7.89	1.27	1.43
1	A	176[B]	THR	CB-OG1	-7.89	1.27	1.43
1	A	278	GLY	N-CA	7.46	1.57	1.46
1	A	16[A]	GLU	CG-CD	-7.18	1.41	1.51
1	A	16[B]	GLU	CG-CD	-7.18	1.41	1.51
1	A	65[A]	GLU	CD-OE2	-6.96	1.18	1.25
1	A	65[B]	GLU	CD-OE2	-6.96	1.18	1.25
1	A	203[A]	GLN	CG-CD	-6.50	1.36	1.51
1	A	203[B]	GLN	CG-CD	-6.50	1.36	1.51
1	A	161	GLN	CD-NE2	6.39	1.48	1.32
1	A	197	SER	CB-OG	-5.80	1.34	1.42
1	A	115	ASP	CB-CG	5.79	1.64	1.51
1	A	277	SER	CB-OG	5.71	1.49	1.42
1	A	323[A]	ALA	C-O	5.65	1.34	1.23
1	A	323[B]	ALA	C-O	5.65	1.34	1.23

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	67[A]	SER	CB-OG	-5.43	1.35	1.42
1	A	67[B]	SER	CB-OG	-5.43	1.35	1.42
1	A	67[A]	SER	N-CA	-5.40	1.35	1.46
1	A	67[B]	SER	N-CA	-5.40	1.35	1.46
1	A	289	SER	CB-OG	-5.33	1.35	1.42
1	A	98[A]	HIS	CG-CD2	5.30	1.44	1.35
1	A	98[B]	HIS	CG-CD2	5.30	1.44	1.35
1	A	74[A]	SER	CA-CB	5.13	1.60	1.52
1	A	74[B]	SER	CA-CB	5.13	1.60	1.52
1	A	313[A]	ASP	CG-OD1	5.09	1.37	1.25
1	A	313[B]	ASP	CG-OD1	5.09	1.37	1.25
1	A	98[A]	HIS	CE1-NE2	-5.03	1.21	1.32
1	A	98[B]	HIS	CE1-NE2	-5.03	1.21	1.32

All (68) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	279	ASP	CA-C-O	-35.82	44.88	120.10
1	A	279	ASP	CB-CG-OD1	32.81	147.83	118.30
1	A	279	ASP	N-CA-CB	-25.03	65.55	110.60
1	A	278	GLY	O-C-N	-20.25	90.30	122.70
1	A	279	ASP	CB-CG-OD2	-19.48	100.77	118.30
1	A	278	GLY	CA-C-N	18.77	158.49	117.20
1	A	206	ASP	CB-CG-OD1	-16.39	103.55	118.30
1	A	279	ASP	C-N-CA	-15.24	90.29	122.30
1	A	313[A]	ASP	CB-CG-OD1	13.72	130.65	118.30
1	A	313[B]	ASP	CB-CG-OD1	13.72	130.65	118.30
1	A	279	ASP	OD1-CG-OD2	-13.47	97.70	123.30
1	A	279	ASP	CA-C-N	-11.18	93.83	116.20
1	A	115	ASP	CB-CG-OD2	-10.48	108.86	118.30
1	A	98[A]	HIS	CG-ND1-CE1	-10.42	92.16	105.70
1	A	98[B]	HIS	CG-ND1-CE1	-10.42	92.16	105.70
1	A	67[A]	SER	N-CA-CB	9.73	125.10	110.50
1	A	67[B]	SER	N-CA-CB	9.73	125.10	110.50
1	A	278	GLY	CA-C-O	-9.43	103.63	120.60
1	A	279	ASP	CB-CA-C	9.18	128.75	110.40
1	A	176[A]	THR	OG1-CB-CG2	-8.95	89.43	110.00
1	A	176[B]	THR	OG1-CB-CG2	-8.95	89.43	110.00
1	A	279	ASP	O-C-N	-8.85	108.16	123.20
1	A	281	SER	CB-CA-C	8.34	125.95	110.10
1	A	248[A]	ASP	CB-CG-OD2	-8.28	110.85	118.30
1	A	248[B]	ASP	CB-CG-OD2	-8.28	110.85	118.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	64	LYS	CD-CE-NZ	-8.24	92.74	111.70
1	A	238[A]	GLN	CG-CD-OE1	-7.94	105.72	121.60
1	A	238[B]	GLN	CG-CD-OE1	-7.94	105.72	121.60
1	A	279	ASP	N-CA-C	-7.93	89.58	111.00
1	A	277	SER	CA-CB-OG	-7.93	89.80	111.20
1	A	280	GLY	N-CA-C	7.84	132.70	113.10
1	A	16[A]	GLU	CG-CD-OE1	7.83	133.96	118.30
1	A	16[B]	GLU	CG-CD-OE1	7.83	133.96	118.30
1	A	52[A]	SER	CB-CA-C	7.42	124.20	110.10
1	A	52[B]	SER	CB-CA-C	7.42	124.20	110.10
1	A	237[A]	GLN	N-CA-CB	7.27	123.69	110.60
1	A	237[B]	GLN	N-CA-CB	7.27	123.69	110.60
1	A	39	LEU	CB-CG-CD2	7.17	123.18	111.00
1	A	206	ASP	OD1-CG-OD2	7.03	136.66	123.30
1	A	279	ASP	CA-CB-CG	-6.97	98.07	113.40
1	A	145	LYS	CD-CE-NZ	6.92	127.61	111.70
1	A	277	SER	C-N-CA	-6.84	107.94	122.30
1	A	249[A]	CYS	CA-CB-SG	6.75	126.15	114.00
1	A	249[B]	CYS	CA-CB-SG	6.75	126.15	114.00
1	A	165	TYR	CB-CG-CD1	6.59	124.96	121.00
1	A	52[A]	SER	N-CA-CB	-6.40	100.90	110.50
1	A	52[B]	SER	N-CA-CB	-6.40	100.90	110.50
1	A	238[A]	GLN	CA-CB-CG	6.22	127.09	113.40
1	A	238[B]	GLN	CA-CB-CG	6.22	127.09	113.40
1	A	117	ASN	OD1-CG-ND2	6.19	136.13	121.90
1	A	165	TYR	CB-CG-CD2	-6.09	117.34	121.00
1	A	67[A]	SER	CA-CB-OG	-5.90	95.26	111.20
1	A	67[B]	SER	CA-CB-OG	-5.90	95.26	111.20
1	A	16[A]	GLU	OE1-CD-OE2	-5.77	116.38	123.30
1	A	16[B]	GLU	OE1-CD-OE2	-5.77	116.38	123.30
1	A	280	GLY	CA-C-O	-5.56	110.60	120.60
1	A	65[A]	GLU	OE1-CD-OE2	5.45	129.84	123.30
1	A	65[B]	GLU	OE1-CD-OE2	5.45	129.84	123.30
1	A	280	GLY	CA-C-N	5.37	129.02	117.20
1	A	98[A]	HIS	ND1-CE1-NE2	5.24	121.42	109.90
1	A	98[B]	HIS	ND1-CE1-NE2	5.24	121.42	109.90
1	A	238[A]	GLN	OE1-CD-NE2	5.18	133.81	121.90
1	A	238[B]	GLN	OE1-CD-NE2	5.18	133.81	121.90
1	A	222	ASP	CB-CG-OD1	5.15	122.94	118.30
1	A	313[A]	ASP	OD1-CG-OD2	-5.08	113.65	123.30
1	A	313[B]	ASP	OD1-CG-OD2	-5.08	113.65	123.30
1	A	98[A]	HIS	CA-CB-CG	5.06	122.20	113.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	98[B]	HIS	CA-CB-CG	5.06	122.20	113.60

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	161	GLN	Sidechain
1	A	278	GLY	Mainchain
1	A	279	ASP	Sidechain,Mainchain,Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2471	2313	2258	38	2
2	A	22	22	19	0	0
3	A	5	0	0	0	0
4	A	44	30	46	0	0
5	A	12	0	14	0	0
6	A	543	0	0	8	1
All	All	3097	2365	2337	38	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ASP:CA	1:A:279:ASP:CB	1.79	1.60
1:A:64:LYS:CE	1:A:64:LYS:NZ	1.70	1.53
1:A:67[B]:SER:OG	1:A:67[B]:SER:CB	1.64	1.42
1:A:279:ASP:OD1	1:A:279:ASP:CG	1.65	1.34
1:A:279:ASP:CA	1:A:279:ASP:C	2.04	1.26
1:A:279:ASP:O	1:A:279:ASP:HA	1.05	1.20
1:A:279:ASP:OD2	1:A:279:ASP:CG	1.83	1.17

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:279:ASP:CA	1:A:279:ASP:N	2.10	1.15
1:A:279:ASP:CA	1:A:279:ASP:O	1.95	1.14
1:A:279:ASP:N	1:A:279:ASP:CB	2.12	1.13
1:A:279:ASP:N	1:A:279:ASP:HB2	1.67	1.08
1:A:64:LYS:NZ	1:A:64:LYS:CD	2.27	0.97
1:A:278:GLY:C	1:A:278:GLY:O	2.04	0.96
1:A:279:ASP:CA	1:A:279:ASP:CG	2.46	0.84
1:A:249[A]:CYS:HB2	6:A:813:HOH:O	1.87	0.73
1:A:176[A]:THR:HG22	6:A:569:HOH:O	1.92	0.68
1:A:243:GLY:HA3	6:A:972:HOH:O	1.98	0.64
1:A:279:ASP:CA	1:A:280:GLY:N	2.60	0.63
1:A:278:GLY:O	1:A:279:ASP:N	2.34	0.61
1:A:176[A]:THR:HG21	6:A:771[A]:HOH:O	1.98	0.61
1:A:64:LYS:NZ	1:A:64:LYS:HD2	2.15	0.59
1:A:18:ILE:CG2	1:A:29:ASN:HB3	2.37	0.55
1:A:67[B]:SER:HG	1:A:67[B]:SER:CB	2.08	0.53
1:A:235:GLY:O	1:A:237[B]:GLN:HG3	2.09	0.52
1:A:279:ASP:OD2	1:A:279:ASP:CB	2.54	0.50
1:A:23[A]:ILE:CD1	1:A:28[A]:LEU:HD12	2.42	0.48
1:A:249[B]:CYS:HB2	6:A:813:HOH:O	2.15	0.46
1:A:58:ASN:ND2	6:A:839:HOH:O	2.50	0.45
1:A:150[A]:GLN:NE2	6:A:717:HOH:O	2.49	0.44
1:A:60[B]:SER:HB3	6:A:839:HOH:O	2.17	0.43

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:262:GLY:HA3	1:A:279:ASP:OD1[1_565]	1.56	0.04
1:A:279:ASP:O	6:A:718:HOH:O[1_545]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	351/323 (109%)	346 (99%)	4 (1%)	1 (0%)	46 16

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	279	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	288/259 (111%)	280 (97%)	8 (3%)	51 12

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	ASP
1	A	58	ASN
1	A	84	ASN
1	A	234[A]	SER
1	A	234[B]	SER
1	A	238[A]	GLN
1	A	238[B]	GLN
1	A	279	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	ASN
1	A	84	ASN
1	A	111	GLN
1	A	117	ASN
1	A	194	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	PP7	A	324	-	43,45,45	0.99	3 (6%)	50,63,63	1.22	3 (6%)
5	GOL	A	325	-	5,5,5	0.97	0	5,5,5	0.41	0
5	GOL	A	326	-	5,5,5	1.60	1 (20%)	5,5,5	2.50	3 (60%)
3	SO4	A	327	-	4,4,4	3.91	2 (50%)	6,6,6	0.95	0
2	MAN	A	328	1	11,11,12	1.40	1 (9%)	14,15,17	1.02	0
2	MAN	A	329	1	11,11,12	2.95	6 (54%)	14,15,17	2.70	8 (57%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	PP7	A	324	-	-	0/61/62/62	0/0/2/2
5	GOL	A	325	-	-	0/4/4/4	0/0/0/0
5	GOL	A	326	-	-	0/4/4/4	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	SO4	A	327	-	-	0/0/0/0	0/0/0/0
2	MAN	A	328	1	-	0/2/19/22	0/1/1/1
2	MAN	A	329	1	-	0/2/19/22	0/1/1/1

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	327	SO4	O4-S	-4.67	1.30	1.47
2	A	329	MAN	O4-C4	-4.52	1.32	1.43
2	A	329	MAN	C1-C2	-3.87	1.42	1.52
2	A	329	MAN	O5-C1	-3.06	1.38	1.43
5	A	326	GOL	O1-C1	-2.89	1.29	1.42
4	A	324	PP7	P-CAL	-2.63	1.82	1.84
4	A	324	PP7	P-O3	-2.10	1.51	1.56
4	A	324	PP7	C6-C5	2.15	1.55	1.52
2	A	329	MAN	O5-C5	2.26	1.48	1.43
2	A	329	MAN	C6-C5	2.67	1.61	1.51
2	A	328	MAN	C2-C3	3.62	1.57	1.52
2	A	329	MAN	C2-C3	5.56	1.60	1.52
3	A	327	SO4	O3-S	6.20	1.69	1.47

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	324	PP7	O5-P-O4	-5.34	105.23	114.91
2	A	329	MAN	C1-O5-C5	-5.08	105.81	112.25
2	A	329	MAN	C6-C5-C4	-4.34	102.31	113.02
2	A	329	MAN	O5-C5-C6	-3.95	98.79	107.35
2	A	329	MAN	C1-C2-C3	-3.63	105.25	109.54
5	A	326	GOL	O2-C2-C1	-3.37	93.19	108.65
2	A	329	MAN	O3-C3-C2	-2.73	105.08	110.00
4	A	324	PP7	O4-P-CAL	-2.36	110.03	114.29
2	A	329	MAN	C3-C4-C5	-2.16	106.42	110.20
2	A	329	MAN	O2-C2-C1	-2.00	105.19	109.21
2	A	329	MAN	O4-C4-C3	2.14	115.16	110.34
4	A	324	PP7	C2-C4-N2	2.53	119.72	116.10
5	A	326	GOL	C3-C2-C1	2.80	122.08	111.12
5	A	326	GOL	O2-C2-C3	3.31	123.82	108.65

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	323/323 (100%)	-0.86	2 (0%) 90 80	8, 12, 23, 83	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	ASP	9.6
1	A	278	GLY	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	SO4	A	327	5/5	0.97	0.17	15.43	22,28,52,59	0
5	GOL	A	325	6/6	0.90	0.09	8.04	16,19,20,27	0
2	MAN	A	329	11/12	0.91	0.10	5.92	18,29,52,65	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	GOL	A	326	6/6	0.95	0.13	3.78	24,36,45,58	0
2	MAN	A	328	11/12	0.98	0.05	0.24	12,15,22,28	0
4	PP7	A	324	44/44	0.99	0.04	0.15	8,11,23,31	2

6.5 Other polymers [i](#)

There are no such residues in this entry.