



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2C3O
Title : CRYSTAL STRUCTURE OF THE FREE RADICAL INTERMEDIATE OF
PYRUVATE:FERREDOXIN OXIDOREDUCTASE FROM DESULFOVIB-
RIO AFRICANUS
Authors : Cavazza, C.; Contreras-Martel, C.; Pieulle, L.; Chabriere, E.; Hatchikian, E.C.;
Fontecilla-Camps, J.C.
Deposited on : 2005-10-11
Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1231	<div> <div>6%</div> <div>56%</div> <div>41%</div> <div>.</div> </div>
1	B	1231	<div> <div>3%</div> <div>57%</div> <div>40%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
6	PYR	A	2239	-	-	-	X

2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 19451 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PYRUVATE-FERREDOXIN OXIDOREDUCTASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	1231	Total	C	N	O	S	0	0	0
			9383	5941	1599	1784	59			
1	B	1231	Total	C	N	O	S	0	0	0
			9383	5941	1599	1784	59			

- Molecule 2 is IRON/SULFUR CLUSTER (three-letter code: SF4) (formula: Fe₄S₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	A	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	Fe	S	0	0
			8	4	4		
2	B	1	Total	Fe	S	0	0
			8	4	4		

- Molecule 3 is THIAMINE DIPHOSPHATE (three-letter code: TPP) (formula: C₁₂H₁₉N₄O₇P₂S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		
3	B	1	Total	C	N	O	P	S	0	0
			26	12	4	7	2	1		

- Molecule 4 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	Mg	0	0
			1	1		
4	A	1	Total	Mg	0	0
			1	1		

- Molecule 5 is CALCIUM ION (three-letter code: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Ca	0	0
			1	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Ca	0	0
			1	1		

- Molecule 6 is PYRUVIC ACID (three-letter code: PYR) (formula: $C_3H_4O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			6	3	3		
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	262	Total	O	0	0
			262	262		
7	B	307	Total	O	0	0
			307	307		

3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: PYRUVATE-FERREDOXIN OXIDOREDUCTASE





● Molecule 1: PYRUVATE-FERREDOXIN OXIDOREDUCTASE



E1164	Q1084	Q1000
H1165	D1085	S1001
M1166	V1086	S1002
	M1087	F1003
T1169	M1088	A1004
	V1091	T1005
F1172	G1094	P1006
A1176		T1007
P1177	F1099	A1008
A1178	R1100	A1009
G1179	Y1101	
G1180	D1102	K1012
K1181	P1103	F1013
A1182	R1104	A1014
G1183	L1105	A1015
H1184		
S1185	M1111	R1019
D1199	Q1114	R1027
T1200	L1115	M1028
P1201		V1029
M1202		M1030
	K1118	
Q1214	A1119	V1037
M1215	P1120	
R1216		V1040
	S1123	
T1219	V1124	G1043
S1220	E1125	Y1044
E1221	F1126	S1045
Q1222	L1127	K1046
	L1128	Q1047
	M1132	Q1048
L1226		F1049
S1227	V1136	
K1228	L1137	V1052
A1229	D1138	
T1230	R1139	A1056
K1231		
K1232	E1143	F1059
	D1144	P1060
	A1145	G1061
	K1146	P1062
	R1147	S1063
	L1148	L1064
		V1065
	Q1151	L1066
	V1152	
	E1155	G1071
	L1156	
	D1157	Q1074
	V1158	
	R1159	R1077
	F1160	K1078
	K1161	M1080
	L1162	G1081
		K1082
		S1082

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	85.65Å 145.24Å 204.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	38.11 – 2.70 38.11 – 2.70	Depositor EDS
% Data completeness (in resolution range)	94.4 (38.11-2.70) 94.5 (38.11-2.70)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 2.69Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.205 , 0.268 0.203 , 0.265	Depositor DCC
R_{free} test set	3476 reflections (5.50%)	DCC
Wilson B-factor (Å ²)	22.5	Xtriage
Anisotropy	0.666	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 51.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.27$	Xtriage
Outliers	0 of 69293 reflections	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	19451	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, SF4, PYR, CA, TPP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.28	0/9585	0.52	0/12954
1	B	0.29	0/9585	0.54	0/12954
All	All	0.29	0/19170	0.53	0/25908

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	9383	0	9263	525	0
1	B	9383	0	9262	502	0
2	A	24	0	0	0	0
2	B	24	0	0	1	0
3	A	26	0	16	3	0
3	B	26	0	16	1	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
5	A	1	0	0	0	0
5	B	1	0	0	0	0
6	A	6	0	3	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	6	0	3	2	0
7	A	262	0	0	13	0
7	B	307	0	0	13	0
All	All	19451	0	18563	960	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 26.

The worst 5 of 960 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1200:THR:HG22	1:A:1202:MET:H	1.19	1.06
1:B:1077:ARG:HB2	1:B:1077:ARG:HH11	1.22	1.02
1:B:239:VAL:HG13	1:B:240:PRO:HD3	1.42	1.02
1:A:198:LYS:H	1:A:198:LYS:HD2	1.24	1.00
1:A:147:GLN:HE22	1:A:184:GLU:H	1.10	0.97

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	1229/1231 (100%)	1105 (90%)	96 (8%)	28 (2%)	8	20
1	B	1229/1231 (100%)	1120 (91%)	89 (7%)	20 (2%)	12	30
All	All	2458/2462 (100%)	2225 (90%)	185 (8%)	48 (2%)	9	24

5 of 48 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	1231	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	556	GLY
1	B	577	GLU
1	B	597	LYS
1	A	4	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	978/978 (100%)	946 (97%)	32 (3%)	45	76
1	B	978/978 (100%)	943 (96%)	35 (4%)	42	73
All	All	1956/1956 (100%)	1889 (97%)	67 (3%)	44	75

5 of 67 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1088	ASN
1	B	180	ILE
1	B	1047	GLN
1	A	1183	ASP
1	B	84	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 103 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1108	GLN
1	B	164	ASN
1	B	1048	GLN
1	A	1151	GLN
1	B	46	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SF4	A	2233	1	0,12,12	0.00	-	0,24,24	0.00	-
2	SF4	A	2234	1	0,12,12	0.00	-	0,24,24	0.00	-
2	SF4	A	2235	1	0,12,12	0.00	-	0,24,24	0.00	-
3	TPP	A	2236	4	20,27,27	3.95	8 (40%)	31,40,40	2.01	7 (22%)
6	PYR	A	2239	-	2,5,5	0.98	0	2,6,6	1.42	0
2	SF4	B	2233	1	0,12,12	0.00	-	0,24,24	0.00	-
2	SF4	B	2234	1	0,12,12	0.00	-	0,24,24	0.00	-
2	SF4	B	2235	1	0,12,12	0.00	-	0,24,24	0.00	-
3	TPP	B	2236	4	20,27,27	3.89	7 (35%)	31,40,40	2.10	7 (22%)
6	PYR	B	2239	-	2,5,5	0.77	0	2,6,6	1.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SF4	A	2233	1	-	0/0/48/48	0/6/5/5
2	SF4	A	2234	1	-	0/0/48/48	0/6/5/5
2	SF4	A	2235	1	-	0/0/48/48	0/6/5/5
3	TPP	A	2236	4	-	0/16/17/17	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PYR	A	2239	-	-	0/0/4/4	0/0/0/0
2	SF4	B	2233	1	-	0/0/48/48	0/6/5/5
2	SF4	B	2234	1	-	0/0/48/48	0/6/5/5
2	SF4	B	2235	1	-	0/0/48/48	0/6/5/5
3	TPP	B	2236	4	-	0/16/17/17	0/2/2/2
6	PYR	B	2239	-	-	0/0/4/4	0/0/0/0

The worst 5 of 15 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	2236	TPP	CM4-C4	2.50	1.55	1.49
3	B	2236	TPP	C5'-C4'	2.62	1.49	1.42
3	A	2236	TPP	C5'-C4'	3.43	1.51	1.42
3	A	2236	TPP	C6'-C5'	3.57	1.45	1.37
3	B	2236	TPP	C2'-N1'	3.71	1.40	1.34

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	2236	TPP	C6-C5-S1	-5.57	112.44	120.24
3	A	2236	TPP	C6-C5-S1	-3.45	115.41	120.24
3	A	2236	TPP	N1'-C2'-N3'	-2.85	120.32	125.60
3	B	2236	TPP	N1'-C2'-N3'	-2.65	120.70	125.60
3	B	2236	TPP	C6'-C5'-C4'	-2.07	112.75	115.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	2236	TPP	3	0
2	B	2233	SF4	1	0
3	B	2236	TPP	1	0
6	B	2239	PYR	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	1231/1231 (100%)	0.12	68 (5%)	29 27	3, 26, 71, 114	0
1	B	1231/1231 (100%)	-0.15	38 (3%)	52 52	2, 18, 55, 117	0
All	All	2462/2462 (100%)	-0.02	106 (4%)	39 38	2, 22, 64, 117	0

The worst 5 of 106 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	1180	GLY	7.7
1	B	1182	ALA	7.5
1	A	631	ALA	6.8
1	A	595	GLY	6.4
1	A	626	PRO	5.8

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	PYR	A	2239	6/6	0.82	0.29	3.42	50,55,57,58	0
6	PYR	B	2239	6/6	0.94	0.17	0.86	17,23,25,26	0
3	TPP	A	2236	26/26	0.96	0.14	-0.45	23,36,48,52	0
3	TPP	B	2236	26/26	0.97	0.13	-0.55	3,14,23,31	0
5	CA	B	2238	1/1	0.97	0.10	-1.36	47,47,47,47	0
5	CA	A	2238	1/1	0.92	0.10	-1.90	61,61,61,61	0
2	SF4	A	2233	8/8	0.97	0.10	-2.42	32,34,36,36	0
2	SF4	A	2235	8/8	0.97	0.07	-3.05	20,25,26,28	0
2	SF4	A	2234	8/8	0.97	0.05	-3.08	28,29,32,33	0
2	SF4	B	2233	8/8	0.98	0.04	-3.40	20,22,24,24	0
2	SF4	B	2234	8/8	0.98	0.05	-3.52	7,10,12,13	0
4	MG	B	2237	1/1	0.99	0.05	-3.78	3,3,3,3	0
4	MG	A	2237	1/1	0.94	0.07	-3.91	16,16,16,16	0
2	SF4	B	2235	8/8	0.98	0.05	-4.38	6,11,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.