



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 03:20 PM GMT

PDB ID : 4C5N
Title : Structure of the pyridoxal kinase from Staphylococcus aureus in complex with AMP-PCP and pyridoxal
Authors : Nodwell, M.; Alte, F.; Sieber, S.A.; Schneider, S.
Deposited on : 2013-09-12
Resolution : 1.75 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1609 (1.76-1.76)
Clashscore	102246	1730 (1.76-1.76)
Ramachandran outliers	100387	1711 (1.76-1.76)
Sidechain outliers	100360	1711 (1.76-1.76)
RSRZ outliers	91569	1610 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	276	<div> <div>3%</div> <div>87%</div> <div>9%</div> <div>••</div> </div>
1	B	276	<div> <div>4%</div> <div>87%</div> <div>11%</div> <div>•</div> </div>
1	C	276	<div> <div>%</div> <div>88%</div> <div>8%</div> <div>•</div> </div>
1	D	276	<div> <div>4%</div> <div>86%</div> <div>10%</div> <div>••</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	SO4	B	1277	-	-	-	X
4	SO4	D	1278	-	-	-	X
4	SO4	D	1279	-	-	-	X

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 8735 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PHOSPHOMETHYLPYRIMIDINE KINASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	268	Total	C	N	O	S	0	6	0
			2049	1310	326	402	11			
1	B	269	Total	C	N	O	S	0	7	0
			2060	1317	328	404	11			
1	C	266	Total	C	N	O	S	0	4	0
			2036	1298	326	400	12			
1	D	269	Total	C	N	O	S	0	1	0
			2039	1297	328	402	12			

- Molecule 2 is 3-HYDROXY-5-(HYDROXYMETHYL)-2-METHYLISONICOTINALDEHYDE (three-letter code: PXL) (formula: C₈H₉NO₃).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			12	8	1	3		
2	B	1	Total	C	N	O	0	0
			12	8	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	C	1	Total	C	N	O	0	0
			12	8	1	3		

- Molecule 3 is PHOSPHOMETHYLPHOSPHONIC ACID ADENYLATE ESTER (three-letter code: ACP) (formula: $C_{11}H_{18}N_5O_{12}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	1
			62	22	10	24	6		
3	B	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
3	C	1	Total	C	N	O	P	0	0
			31	11	5	12	3		
3	D	1	Total	C	N	O	P	0	0
			31	11	5	12	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O_4S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is 4,5-BIS(HYDROXYMETHYL)-2-METHYL-PYRIDIN-3-OL (three-letter code: UEG) (formula: C₈H₁₁NO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	D	1	Total	C	N	O	0	0
			12	8	1	3		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	81	Total	O	0	0
			81	81		
6	B	85	Total	O	0	0
			85	85		
6	C	88	Total	O	0	0
			88	88		
6	D	59	Total	O	0	0
			59	59		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: PHOSPHOMETHYLPYRIMIDINE KINASE




• Molecule 1: PHOSPHOMETHYLPYRIMIDINE KINASE



• Molecule 1: PHOSPHOMETHYLPYRIMIDINE KINASE



• Molecule 1: PHOSPHOMETHYLPYRIMIDINE KINASE





4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.19Å 100.66Å 168.21Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.26 – 1.75 48.22 – 1.75	Depositor EDS
% Data completeness (in resolution range)	99.7 (48.26-1.75) 99.8 (48.22-1.75)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.204 , 0.253 0.211 , 0.258	Depositor DCC
R_{free} test set	5333 reflections (5.24%)	DCC
Wilson B-factor (Å ²)	22.2	Xtriage
Anisotropy	0.470	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtriage
Outliers	1 of 107272 reflections (0.001%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8735	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 22.38 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 5.7692e-03.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: UEG, ACP, SO4, PXL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.92	0/2104	0.87	0/2851
1	B	0.88	0/2118	0.88	2/2871 (0.1%)
1	C	0.96	1/2085 (0.0%)	0.93	1/2823 (0.0%)
1	D	0.95	1/2079 (0.0%)	0.95	6/2816 (0.2%)
All	All	0.93	2/8386 (0.0%)	0.91	9/11361 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	94	TYR	CE1-CZ	-7.88	1.28	1.38
1	D	119	PRO	N-CD	5.42	1.55	1.47

The worst 5 of 9 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	59	ASP	CB-CG-OD1	8.11	125.60	118.30
1	D	270	ASP	CB-CG-OD1	6.85	124.47	118.30
1	B	167	ASP	CB-CG-OD1	6.38	124.04	118.30
1	C	89	ARG	NE-CZ-NH1	6.35	123.47	120.30
1	D	89	ARG	NE-CZ-NH1	5.78	123.19	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2049	0	2047	16	0
1	B	2060	0	2060	20	1
1	C	2036	0	2030	19	0
1	D	2039	0	2019	19	1
2	A	12	0	9	1	0
2	B	12	0	9	1	0
2	C	12	0	9	4	0
3	A	62	0	28	2	0
3	B	31	0	14	3	0
3	C	31	0	14	4	0
3	D	31	0	14	2	0
4	B	10	0	0	0	0
4	C	10	0	0	0	0
4	D	15	0	0	0	0
5	D	12	0	8	2	0
6	A	81	0	0	1	0
6	B	85	0	0	0	0
6	C	88	0	0	0	0
6	D	59	0	0	1	0
All	All	8735	0	8261	79	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:213:GLY:H	3:B:500:ACP:H3B2	1.05	1.09
1:D:118:ASN:O	1:D:120:GLY:N	1.90	1.05
1:B:213:GLY:N	3:B:500:ACP:H3B2	1.85	0.90
2:C:300:PXL:OP4	2:C:300:PXL:O4A	1.89	0.89
2:A:300:PXL:H5A1	2:A:300:PXL:O4A	1.82	0.79

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:270:ASP:OD2	1:D:118:ASN:O[2_454]	1.92	0.28

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	270/276 (98%)	263 (97%)	7 (3%)	0	100	100
1	B	272/276 (99%)	263 (97%)	9 (3%)	0	100	100
1	C	266/276 (96%)	260 (98%)	6 (2%)	0	100	100
1	D	266/276 (96%)	257 (97%)	8 (3%)	1 (0%)	39	19
All	All	1074/1104 (97%)	1043 (97%)	30 (3%)	1 (0%)	56	36

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	119	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/221 (98%)	210 (97%)	7 (3%)	46	20
1	B	219/221 (99%)	216 (99%)	3 (1%)	74	58
1	C	217/221 (98%)	213 (98%)	4 (2%)	66	46
1	D	215/221 (97%)	205 (95%)	10 (5%)	32	10
All	All	868/884 (98%)	844 (97%)	24 (3%)	52	25

5 of 24 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	20	MET
1	C	155	ILE
1	D	250	LYS
1	C	51	HIS
1	C	107	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	209	ASN
1	C	100	GLN
1	C	153	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

16 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PXL	A	300	-	12,12,12	2.31	2 (16%)	15,16,16	1.65	3 (20%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ACP	A	500[A]	-	25,33,33	2.48	6 (24%)	31,52,52	2.86	12 (38%)
3	ACP	A	500[B]	-	25,33,33	1.77	7 (28%)	31,52,52	1.98	3 (9%)
4	SO4	B	1277	-	4,4,4	0.73	0	6,6,6	0.32	0
4	SO4	B	1278	-	4,4,4	0.23	0	6,6,6	0.21	0
2	PXL	B	300	-	12,12,12	1.68	1 (8%)	15,16,16	2.33	6 (40%)
3	ACP	B	500	-	25,33,33	2.17	5 (20%)	31,52,52	1.96	10 (32%)
4	SO4	C	1277	-	4,4,4	1.00	0	6,6,6	1.13	1 (16%)
4	SO4	C	1278	-	4,4,4	0.34	0	6,6,6	0.19	0
2	PXL	C	300	-	12,12,12	0.92	0	15,16,16	1.24	1 (6%)
3	ACP	C	500	-	25,33,33	2.81	7 (28%)	31,52,52	2.68	10 (32%)
4	SO4	D	1277	-	4,4,4	0.45	0	6,6,6	0.28	0
4	SO4	D	1278	-	4,4,4	0.63	0	6,6,6	0.33	0
4	SO4	D	1279	-	4,4,4	1.21	0	6,6,6	0.32	0
5	UEG	D	300	1	12,12,12	2.67	2 (16%)	14,16,16	1.70	3 (21%)
3	ACP	D	500	-	25,33,33	2.13	8 (32%)	31,52,52	2.39	10 (32%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PXL	A	300	-	-	0/4/4/4	0/1/1/1
3	ACP	A	500[A]	-	-	0/15/38/38	0/3/3/3
3	ACP	A	500[B]	-	-	0/15/38/38	0/3/3/3
4	SO4	B	1277	-	-	0/0/0/0	0/0/0/0
4	SO4	B	1278	-	-	0/0/0/0	0/0/0/0
2	PXL	B	300	-	-	0/4/4/4	0/1/1/1
3	ACP	B	500	-	-	0/15/38/38	0/3/3/3
4	SO4	C	1277	-	-	0/0/0/0	0/0/0/0
4	SO4	C	1278	-	-	0/0/0/0	0/0/0/0
2	PXL	C	300	-	-	0/4/4/4	0/1/1/1
3	ACP	C	500	-	-	0/15/38/38	0/3/3/3
4	SO4	D	1277	-	-	0/0/0/0	0/0/0/0
4	SO4	D	1278	-	-	0/0/0/0	0/0/0/0
4	SO4	D	1279	-	-	0/0/0/0	0/0/0/0
5	UEG	D	300	1	-	0/4/4/4	0/1/1/1
3	ACP	D	500	-	-	0/15/38/38	0/3/3/3

The worst 5 of 38 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	D	300	UEG	C3-C2	-7.63	1.35	1.40
2	A	300	PXL	C3-C2	-6.49	1.36	1.40
5	D	300	UEG	O4A-C4A	-4.88	1.20	1.41
2	B	300	PXL	C4-C3	-4.28	1.35	1.40
3	A	500[B]	ACP	PG-O3G	-2.91	1.47	1.54

The worst 5 of 59 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	500[A]	ACP	N3-C2-N1	-9.46	121.65	128.89
3	A	500[B]	ACP	N3-C2-N1	-8.60	122.31	128.89
3	C	500	ACP	N3-C2-N1	-8.07	122.72	128.89
3	D	500	ACP	N3-C2-N1	-7.34	123.27	128.89
3	B	500	ACP	N3-C2-N1	-5.73	124.51	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

8 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	300	PXL	1	0
3	A	500[A]	ACP	2	0
2	B	300	PXL	1	0
3	B	500	ACP	3	0
2	C	300	PXL	4	0
3	C	500	ACP	4	0
5	D	300	UEG	2	0
3	D	500	ACP	2	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	268/276 (97%)	0.36	9 (3%) 49 55	16, 27, 43, 55	0
1	B	269/276 (97%)	0.22	10 (3%) 45 51	16, 26, 43, 75	0
1	C	266/276 (96%)	0.03	4 (1%) 76 82	13, 22, 35, 65	0
1	D	269/276 (97%)	0.25	10 (3%) 45 51	15, 24, 40, 60	0
All	All	1072/1104 (97%)	0.22	33 (3%) 52 58	13, 24, 41, 75	0

The worst 5 of 33 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1	GLY	8.3
1	B	109	VAL	5.8
1	D	109	VAL	4.8
1	A	276	VAL	4.7
1	A	98	ASN	4.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
4	SO4	B	1277	5/5	0.92	0.33	4.95	49,53,54,54	0
4	SO4	D	1278	5/5	0.92	0.26	4.34	53,54,55,57	0
4	SO4	D	1279	5/5	0.90	0.16	3.00	35,47,52,52	0
3	ACP	A	500[B]	31/31	0.92	0.13	1.67	16,24,29,30	31
3	ACP	A	500[A]	31/31	0.92	0.13	1.60	18,20,54,59	31
4	SO4	C	1278	5/5	0.91	0.19	1.52	34,37,39,39	5
2	PXL	B	300	12/12	0.90	0.14	0.31	21,29,40,43	0
3	ACP	D	500	31/31	0.92	0.12	0.20	20,28,60,76	0
3	ACP	C	500	31/31	0.94	0.09	0.20	14,18,42,61	0
3	ACP	B	500	31/31	0.90	0.12	0.17	26,29,52,52	0
2	PXL	A	300	12/12	0.92	0.13	-0.13	23,30,43,54	0
5	UEG	D	300	12/12	0.84	0.12	-0.15	24,33,38,43	0
4	SO4	C	1277	5/5	0.97	0.08	-0.35	27,33,35,36	0
2	PXL	C	300	12/12	0.94	0.09	-0.52	21,28,41,42	0
4	SO4	B	1278	5/5	0.89	0.19	-	41,42,43,44	5
4	SO4	D	1277	5/5	0.97	0.13	-	45,50,51,53	0

6.5 Other polymers [i](#)

There are no such residues in this entry.