



Full wwPDB X-ray Structure Validation Report ⓘ

May 26, 2016 – 10:53 PM EDT

PDB ID : 5C67
Title : Human Mesotrypsin in complex with amyloid precursor protein inhibitor variant APPI-M17G/I18F/F34V
Authors : Kayode, O.; Sankaran, B.; Radisky, E.S.
Deposited on : 2015-06-22
Resolution : 1.83 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : unknown
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027674
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027674


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.83 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2634 (1.86-1.82)
Clashscore	102246	2862 (1.86-1.82)
Ramachandran outliers	100387	2831 (1.86-1.82)
Sidechain outliers	100360	2832 (1.86-1.82)
RSRZ outliers	91569	2639 (1.86-1.82)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	224	 3% 92% 8%
1	B	224	 5% 92% 8%
2	C	81	 % 63% 35%
2	E	81	 63% 33%

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4289 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Trypsin-3.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	224	1700	1070	294	324	12	0	1	0
1	B	224	1690	1065	291	322	12	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	ALA	THR	variant	UNP P35030
A	195	ALA	SER	engineered mutation	UNP P35030
B	127	ALA	THR	variant	UNP P35030
B	195	ALA	SER	engineered mutation	UNP P35030

- Molecule 2 is a protein called Amyloid beta A4 protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	E	54	406	250	69	80	7	0	0	0
2	C	53	406	250	68	81	7	0	0	0

There are 62 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
E	-9	TYR	-	expression tag	UNP P05067
E	-8	VAL	-	expression tag	UNP P05067
E	-7	ASP	-	expression tag	UNP P05067
E	-6	TYR	-	expression tag	UNP P05067
E	-5	LYS	-	expression tag	UNP P05067
E	-4	ASP	-	expression tag	UNP P05067
E	-3	ASP	-	expression tag	UNP P05067
E	-2	ASP	-	expression tag	UNP P05067

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
E	-1	ASP	-	expression tag	UNP P05067
E	0	LYS	-	expression tag	UNP P05067
E	1	GLU	-	expression tag	UNP P05067
E	2	PHE	-	expression tag	UNP P05067
E	3	GLU	-	expression tag	UNP P05067
E	4	VAL	-	expression tag	UNP P05067
E	5	CYS	-	expression tag	UNP P05067
E	6	SER	-	expression tag	UNP P05067
E	7	GLU	-	expression tag	UNP P05067
E	17	GLY	MET	engineered mutation	UNP P05067
E	18	PHE	ILE	engineered mutation	UNP P05067
E	34	VAL	PHE	engineered mutation	UNP P05067
E	61	ARG	-	expression tag	UNP P05067
E	62	HIS	-	expression tag	UNP P05067
E	63	HIS	-	expression tag	UNP P05067
E	64	HIS	-	expression tag	UNP P05067
E	65	HIS	-	expression tag	UNP P05067
E	66	HIS	-	expression tag	UNP P05067
E	67	HIS	-	expression tag	UNP P05067
E	68	ALA	-	expression tag	UNP P05067
E	69	ALA	-	expression tag	UNP P05067
E	70	ALA	-	expression tag	UNP P05067
E	71	ASN	-	expression tag	UNP P05067
C	-9	TYR	-	expression tag	UNP P05067
C	-8	VAL	-	expression tag	UNP P05067
C	-7	ASP	-	expression tag	UNP P05067
C	-6	TYR	-	expression tag	UNP P05067
C	-5	LYS	-	expression tag	UNP P05067
C	-4	ASP	-	expression tag	UNP P05067
C	-3	ASP	-	expression tag	UNP P05067
C	-2	ASP	-	expression tag	UNP P05067
C	-1	ASP	-	expression tag	UNP P05067
C	0	LYS	-	expression tag	UNP P05067
C	1	GLU	-	expression tag	UNP P05067
C	2	PHE	-	expression tag	UNP P05067
C	3	GLU	-	expression tag	UNP P05067
C	4	VAL	-	expression tag	UNP P05067
C	5	CYS	-	expression tag	UNP P05067
C	6	SER	-	expression tag	UNP P05067
C	7	GLU	-	expression tag	UNP P05067
C	17	GLY	MET	engineered mutation	UNP P05067
C	18	PHE	ILE	engineered mutation	UNP P05067

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	34	VAL	PHE	engineered mutation	UNP P05067
C	61	ARG	-	expression tag	UNP P05067
C	62	HIS	-	expression tag	UNP P05067
C	63	HIS	-	expression tag	UNP P05067
C	64	HIS	-	expression tag	UNP P05067
C	65	HIS	-	expression tag	UNP P05067
C	66	HIS	-	expression tag	UNP P05067
C	67	HIS	-	expression tag	UNP P05067
C	68	ALA	-	expression tag	UNP P05067
C	69	ALA	-	expression tag	UNP P05067
C	70	ALA	-	expression tag	UNP P05067
C	71	ASN	-	expression tag	UNP P05067


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	46	Total O 46 46	0	0
3	E	7	Total O 7 7	0	0
3	B	26	Total O 26 26	0	0
3	C	8	Total O 8 8	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Trypsin-3




- Molecule 1: Trypsin-3



- Molecule 2: Amyloid beta A4 protein



- Molecule 2: Amyloid beta A4 protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	78.15Å 78.15Å 243.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	20.00 – 1.83 19.99 – 1.83	Depositor EDS
% Data completeness (in resolution range)	99.8 (20.00-1.83) 100.0 (19.99-1.83)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.47 (at 1.82Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.223 , 0.267 0.229 , 0.273	Depositor DCC
R_{free} test set	974 reflections (1.46%)	DCC
Wilson B-factor (Å ²)	27.9	Xtrriage
Anisotropy	0.075	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 38.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4289	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.18	1/1738 (0.1%)	1.07	5/2364 (0.2%)
1	B	0.94	3/1729 (0.2%)	0.96	6/2353 (0.3%)
2	C	1.00	0/416	0.99	1/561 (0.2%)
2	E	1.13	0/416	0.97	1/561 (0.2%)
All	All	1.07	4/4299 (0.1%)	1.01	13/5839 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	135	GLU	CD-OE2	-6.51	1.18	1.25
1	B	154	GLU	CG-CD	6.40	1.61	1.51
1	B	154	GLU	CB-CG	5.67	1.62	1.52
1	A	214	SER	CA-CB	5.66	1.61	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	102	ASP	CB-CG-OD1	6.96	124.56	118.30
1	A	189	ASP	CB-CG-OD1	6.73	124.35	118.30
1	B	154	GLU	OE1-CD-OE2	-6.24	115.81	123.30
1	A	156	LYS	CD-CE-NZ	-5.98	97.94	111.70
1	A	46	LEU	CB-CG-CD2	-5.83	101.09	111.00
1	B	194	ASP	CB-CG-OD1	5.83	123.54	118.30
1	A	119	SER	N-CA-CB	-5.63	102.05	110.50
2	E	15	ARG	NE-CZ-NH1	5.61	123.10	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	66	ARG	NE-CZ-NH2	-5.55	117.53	120.30
1	B	66	ARG	NE-CZ-NH1	5.46	123.03	120.30
1	A	100	ASP	CB-CG-OD1	5.22	123.00	118.30
2	C	20	ARG	NE-CZ-NH1	5.19	122.89	120.30
1	B	102	ASP	CB-CG-OD2	-5.07	113.74	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	24	GLU	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1700	0	1650	9	0
1	B	1690	0	1637	9	0
2	C	406	0	349	1	0
2	E	406	0	348	2	0
3	A	46	0	0	0	0
3	B	26	0	0	0	0
3	C	8	0	0	0	0
3	E	7	0	0	0	0
All	All	4289	0	3984	19	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (19) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:74:LYS:HD3	1:B:153:ASP:OD1	1.93	0.69
1:A:195:ALA:HB2	2:E:15:ARG:C	2.19	0.63
1:A:28:PRO:HB2	1:A:119:SER:HB3	1.80	0.62

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:75:VAL:O	1:A:75:VAL:HG12	2.07	0.53
1:A:213:VAL:HG22	1:A:228:TYR:HE1	1.75	0.52
1:A:82:PHE:CD1	1:A:82:PHE:N	2.77	0.52
2:E:4:VAL:HG11	2:E:42:ARG:CZ	2.40	0.51
1:A:82:PHE:HD1	1:A:82:PHE:N	2.09	0.51
1:A:81:GLN:C	1:A:82:PHE:HD1	2.15	0.49
1:B:153:ASP:HB2	1:B:154:GLU:OE1	2.13	0.47
1:B:195:ALA:HB2	2:C:15:ARG:C	2.35	0.46
1:B:101:ASN:HA	1:B:234:TYR:OH	2.18	0.44
1:B:154:GLU:HG2	1:B:156:LYS:HE2	2.01	0.42
1:B:72:ASN:HA	1:B:153:ASP:O	2.20	0.42
1:B:211:GLY:HA2	1:B:229:THR:O	2.20	0.41
1:A:105:LEU:HD11	1:A:238:ILE:HA	2.01	0.41
1:B:56:ALA:HA	1:B:104:MET:HB2	2.01	0.41
1:A:142:GLY:HA2	1:A:193:ARG:HB3	2.03	0.40
1:B:55:ALA:O	1:B:58:CYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	223/224 (100%)	215 (96%)	8 (4%)	0	100	100
1	B	222/224 (99%)	211 (95%)	11 (5%)	0	100	100
2	C	51/81 (63%)	48 (94%)	3 (6%)	0	100	100
2	E	52/81 (64%)	50 (96%)	2 (4%)	0	100	100
All	All	548/610 (90%)	524 (96%)	24 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	184/185 (100%)	184 (100%)	0	100	100
1	B	183/185 (99%)	183 (100%)	0	100	100
2	C	42/65 (65%)	42 (100%)	0	100	100
2	E	41/65 (63%)	41 (100%)	0	100	100
All	All	450/500 (90%)	450 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	224/224 (100%)	-0.01	6 (2%) 58 54	17, 27, 47, 67	0
1	B	224/224 (100%)	0.21	12 (5%) 29 26	20, 36, 57, 79	0
2	C	53/81 (65%)	-0.01	1 (1%) 70 67	24, 32, 51, 59	0
2	E	54/81 (66%)	-0.18	0 100 100	22, 29, 46, 49	0
All	All	555/610 (90%)	0.06	19 (3%) 49 45	17, 31, 55, 79	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	78	GLY	3.4
1	B	76	LEU	3.1
1	B	246	SER	3.1
1	B	147	PHE	3.0
1	A	246	SER	2.9
1	B	75	VAL	2.8
1	B	74	LYS	2.5
1	B	178	ASN	2.5
1	B	38	GLY	2.4
1	A	78	GLY	2.3
1	B	61	THR	2.3
1	B	52	VAL	2.2
1	B	77	GLU	2.2
1	A	97	ASP	2.2
1	A	38	GLY	2.1
1	A	147	PHE	2.1
1	B	96	ARG	2.0
1	A	62	ARG	2.0
2	C	3	GLU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.