



# Full wwPDB X-ray Structure Validation Report ⓘ

Jul 14, 2016 – 11:28 PM EDT

PDB ID : 5C8B  
Title : Structural insights into the redesign of a sucrose phosphorylase by induced loop repositioning  
Authors : Grimm, C.; Kraus, M.  
Deposited on : 2015-06-25  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027790  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027790


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	B	514	<div> <div>4%</div> <div>86%</div> <div>11% ..</div> </div>

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7946 atoms, of which 3826 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Sucrose phosphorylase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	B	504	Total	C	H	N	O	S	0	0	0
			7780	2511	3814	670	772	13			

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	MET	-	initiating methionine	UNP Q84HQ2
B	0	ALA	-	expression tag	UNP Q84HQ2
B	345	PHE	GLN	engineered mutation	UNP Q84HQ2
B	505	LEU	-	expression tag	UNP Q84HQ2
B	506	GLU	-	expression tag	UNP Q84HQ2
B	507	HIS	-	expression tag	UNP Q84HQ2
B	508	HIS	-	expression tag	UNP Q84HQ2
B	509	HIS	-	expression tag	UNP Q84HQ2
B	510	HIS	-	expression tag	UNP Q84HQ2
B	511	HIS	-	expression tag	UNP Q84HQ2
B	512	HIS	-	expression tag	UNP Q84HQ2

- Molecule 2 is BETA-D-GLUCOSE (three-letter code: BGC) (formula: C<sub>6</sub>H<sub>12</sub>O<sub>6</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	B	1	Total	C	H	O	0	0
			24	6	12	6		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	142	Total	O	0	0
			142	142		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Sucrose phosphorylase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	82.50Å 82.50Å 156.38Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.80 – 2.70 46.76 – 2.68	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.80-2.70) 92.4 (46.76-2.68)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.78 (at 2.69Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.9_1692)	Depositor
R, $R_{free}$	0.208 , 0.287 0.203 , 0.282	Depositor DCC
$R_{free}$ test set	726 reflections (5.02%)	DCC
Wilson B-factor (Å <sup>2</sup> )	77.2	Xtrriage
Anisotropy	0.258	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 51.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7946	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	104.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	B	0.77	0/4061	0.83	1/5523 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	50	ASP	CB-CG-OD2	-5.17	113.64	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	106	SER	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	3966	3814	3824	40	0
2	B	12	12	12	0	0
3	B	142	0	0	17	0
All	All	4120	3826	3836	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:300:ASP:N	3:B:705:HOH:O	2.18	0.77
1:B:391:LEU:N	3:B:704:HOH:O	2.18	0.76
1:B:403:ARG:O	3:B:701:HOH:O	2.06	0.73
1:B:13:ARG:NH1	3:B:701:HOH:O	2.08	0.72
1:B:48:GLY:O	3:B:702:HOH:O	2.07	0.71
1:B:483:GLU:OE1	1:B:492:ARG:NH1	2.23	0.71
1:B:360:HIS:ND1	3:B:709:HOH:O	2.25	0.69
1:B:397:ASN:ND2	3:B:710:HOH:O	2.25	0.67
1:B:307:LYS:NZ	1:B:312:ASP:OD2	2.31	0.58
1:B:393:ARG:O	3:B:703:HOH:O	2.17	0.58
1:B:448:THR:N	3:B:715:HOH:O	2.37	0.57
1:B:412:ASP:HA	1:B:415:LEU:HD12	1.89	0.53
1:B:50:ASP:C	3:B:707:HOH:O	2.47	0.53
1:B:402:ASN:HB3	3:B:768:HOH:O	2.10	0.52
1:B:391:LEU:O	1:B:394:LYS:N	2.42	0.52
1:B:54:ASP:OD1	3:B:707:HOH:O	2.20	0.49
1:B:133:ARG:HB3	1:B:135:ARG:O	2.13	0.49
1:B:447:ASP:N	3:B:715:HOH:O	2.47	0.47
1:B:269:GLU:N	1:B:270:PRO:CD	2.78	0.47
1:B:434:ASP:N	1:B:434:ASP:OD1	2.38	0.46
1:B:242:GLU:O	1:B:245:SER:OG	2.30	0.46
1:B:450:ILE:HG13	1:B:451:SER:N	2.31	0.46
1:B:495:ASP:OD2	3:B:706:HOH:O	2.19	0.46
1:B:283:VAL:HG13	1:B:376:GLN:HG3	1.97	0.45
1:B:400:ASP:OD2	3:B:708:HOH:O	2.20	0.45
1:B:266:GLY:HA2	1:B:467:PRO:HB2	1.99	0.45
1:B:269:GLU:O	1:B:272:ALA:N	2.50	0.44
1:B:109:TYR:HB3	1:B:110:PRO:HD3	1.99	0.43
1:B:370:PHE:CE1	1:B:463:LEU:HB2	2.54	0.43
1:B:29:ARG:NH2	1:B:412:ASP:OD2	2.41	0.43
1:B:503:VAL:HG12	1:B:504:ALA:N	2.34	0.43

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:492:ARG:HG2	1:B:493:SER:N	2.34	0.43
1:B:190:ARG:NH1	1:B:192:ASP:OD1	2.52	0.43
1:B:13:ARG:HD3	3:B:701:HOH:O	2.18	0.42
1:B:311:PRO:HB2	1:B:313:GLU:HG2	2.01	0.42
1:B:60:LYS:CG	3:B:793:HOH:O	2.66	0.42
1:B:118:VAL:HG21	1:B:153:TRP:CZ3	2.54	0.42
1:B:109:TYR:N	1:B:110:PRO:CD	2.82	0.41
1:B:255:LEU:HB3	1:B:256:PRO:HD3	2.03	0.41
1:B:268:VAL:HG22	1:B:447:ASP:O	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	502/514 (98%)	485 (97%)	17 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	428/437 (98%)	418 (98%)	10 (2%)	58	85

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	26	LEU
1	B	100	LEU
1	B	106	SER
1	B	276	ASP
1	B	397	ASN
1	B	434	ASP
1	B	445	ASP
1	B	446	ASP
1	B	447	ASP
1	B	450	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	397	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	B	601	-	12,12,12	1.26	1 (8%)	17,17,17	1.25	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	601	-	-	0/2/22/22	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	601	BGC	O3-C3	2.49	1.48	1.43

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	601	BGC	O1-C1-O5	-2.26	104.03	110.33
2	B	601	BGC	O3-C3-C4	2.54	116.09	110.36

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	B	504/514 (98%)	0.00	18 (3%)	46 46	60, 91, 139, 166	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	302	LEU	4.4
1	B	502	VAL	3.5
1	B	484	TRP	3.3
1	B	482	LEU	3.3
1	B	481	MET	3.3
1	B	122	GLY	3.0
1	B	455	ARG	3.0
1	B	494	ASP	2.9
1	B	301	GLN	2.7
1	B	460	GLN	2.7
1	B	504	ALA	2.6
1	B	339	SER	2.6
1	B	499	ASN	2.5
1	B	342	ASP	2.5
1	B	493	SER	2.5
1	B	496	LEU	2.4
1	B	461	ALA	2.2
1	B	333	ALA	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	B	601	12/12	0.96	0.18	0.74	62,87,105,106	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.