



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:28 PM GMT

PDB ID : 4CCP  
Title : X-RAY STRUCTURES OF RECOMBINANT YEAST CYTOCHROME C  
PEROXIDASE AND THREE HEME-CLEFT MUTANTS PREPARED BY  
SITE-DIRECTED MUTAGENESIS  
Authors : Wang, J.; Mauro, J.M.; Edwards, S.L.; Oatley, S.J.; Fishel, L.A.; Ashford,  
V.A.; Xuong, N.-H.; Kraut, J.  
Deposited on : 1990-02-28  
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : **NOT EXECUTED**  
EDS : **NOT EXECUTED**  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	296	 70% 25% . .

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2591 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YEAST CYTOCHROME C PEROXIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	S	0	0	0
			2336	1494	385	451	6			

There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	PHE	TRP	ENGINEERED MUTATION	UNP P00431
A	53	ILE	THR	VARIANT	UNP P00431
A	152	GLY	ASP	VARIANT	UNP P00431

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	212	Total 212	O 212	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

#### • Molecule 1: YEAST CYTOCHROME C PEROXIDASE



## 4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	104.76Å 74.21Å 45.67Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	(Not available) – 2.20	Depositor
% Data completeness (in resolution range)	(Not available) ((Not available)-2.20)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, $R_{free}$	0.160 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2591	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	38.0	wwPDB-VP

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.00	1/2402 (0.0%)	1.55	32/3257 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	2	THR	N-CA	10.41	1.67	1.46

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	143	ARG	NE-CZ-NH1	-13.88	113.36	120.30
1	A	166	ARG	NE-CZ-NH1	10.52	125.56	120.30
1	A	132	ASP	CB-CG-OD1	10.06	127.36	118.30
1	A	143	ARG	NE-CZ-NH2	8.74	124.67	120.30
1	A	130	ARG	NE-CZ-NH1	8.50	124.55	120.30
1	A	150	ASP	CB-CG-OD1	8.35	125.82	118.30
1	A	14	ARG	NE-CZ-NH1	-7.74	116.43	120.30
1	A	127	ARG	CD-NE-CZ	7.25	133.75	123.60
1	A	132	ASP	CB-CG-OD2	-7.17	111.85	118.30
1	A	279	ASP	CB-CG-OD2	-6.93	112.06	118.30
1	A	130	ARG	NE-CZ-NH2	-6.76	116.92	120.30
1	A	165	ASP	CB-CG-OD2	-6.75	112.22	118.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	235	ASP	CB-CG-OD2	-6.32	112.61	118.30
1	A	71	TYR	CB-CG-CD2	-5.92	117.45	121.00
1	A	254	ASP	CB-CG-OD1	5.90	123.61	118.30
1	A	71	TYR	CB-CG-CD1	5.86	124.52	121.00
1	A	150	ASP	CB-CG-OD2	-5.75	113.13	118.30
1	A	165	ASP	CB-CG-OD1	5.69	123.42	118.30
1	A	157	PHE	CB-CG-CD1	-5.68	116.82	120.80
1	A	31	ARG	NE-CZ-NH2	-5.56	117.52	120.30
1	A	34	ASP	CB-CG-OD1	5.53	123.28	118.30
1	A	162	ASN	N-CA-CB	-5.52	100.66	110.60
1	A	143	ARG	CD-NE-CZ	5.44	131.22	123.60
1	A	254	ASP	CB-CG-OD2	-5.42	113.42	118.30
1	A	144	LEU	CB-CA-C	5.24	120.15	110.20
1	A	136	ASP	CB-CG-OD1	5.18	122.97	118.30
1	A	155	ARG	NE-CZ-NH2	5.18	122.89	120.30
1	A	96	HIS	CA-CB-CG	-5.17	104.81	113.60
1	A	253	ASN	CA-CB-CG	-5.17	102.03	113.40
1	A	93	GLU	CA-CB-CG	5.15	124.73	113.40
1	A	67	TYR	CB-CG-CD2	-5.09	117.95	121.00
1	A	253	ASN	CB-CG-OD1	-5.07	111.46	121.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	143	ARG	Sidechain

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2336	0	2171	69	1
2	A	43	0	30	2	0
3	A	212	0	0	6	2
All	All	2591	0	2201	70	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 15.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:LEU:CD2	1:A:4:LEU:H	1.75	0.98
1:A:4:LEU:HD22	1:A:4:LEU:H	1.31	0.94
1:A:195:ASN:H	1:A:195:ASN:HD22	1.12	0.92
1:A:86:GLN:NE2	1:A:86:GLN:H	1.68	0.90
1:A:20:GLN:HE22	1:A:287:LYS:H	1.19	0.86
1:A:216:ASN:HD22	1:A:222:GLN:HE21	1.25	0.84
1:A:216:ASN:HD22	1:A:222:GLN:NE2	1.78	0.81
1:A:84:GLY:N	1:A:86:GLN:HE22	1.88	0.72
1:A:4:LEU:N	1:A:4:LEU:HD22	2.06	0.70
1:A:86:GLN:CG	3:A:684:HOH:O	2.39	0.69
1:A:4:LEU:HD23	1:A:61:ASP:O	1.93	0.69
1:A:294:LEU:N	1:A:294:LEU:HD12	2.06	0.68
1:A:84:GLY:H	1:A:86:GLN:NE2	1.91	0.68
1:A:133:THR:HB	1:A:134:PRO:HD2	1.77	0.67
1:A:4:LEU:CD2	1:A:61:ASP:O	2.43	0.66
1:A:84:GLY:N	1:A:86:GLN:NE2	2.44	0.66
1:A:84:GLY:H	1:A:86:GLN:HE22	1.44	0.66
1:A:294:LEU:N	1:A:294:LEU:CD1	2.60	0.64
1:A:20:GLN:HE22	1:A:287:LYS:N	1.96	0.63
1:A:255:GLN:HG2	3:A:556:HOH:O	1.99	0.63
1:A:293:GLY:C	1:A:294:LEU:HD12	2.18	0.62
1:A:84:GLY:C	1:A:86:GLN:HE21	2.03	0.62
1:A:213:LEU:HD11	1:A:221:GLU:HB3	1.82	0.61
1:A:195:ASN:ND2	1:A:195:ASN:H	1.93	0.60
1:A:86:GLN:HG2	3:A:684:HOH:O	1.99	0.59
1:A:161:LEU:O	1:A:162:ASN:CB	2.49	0.59
1:A:17:GLU:HG2	3:A:317:HOH:O	2.06	0.56
1:A:212:LYS:HB3	3:A:812:HOH:O	2.05	0.55
1:A:216:ASN:ND2	1:A:222:GLN:HE21	2.01	0.54
1:A:4:LEU:N	1:A:4:LEU:CD2	2.53	0.54
1:A:159:GLN:HA	1:A:159:GLN:NE2	2.22	0.54
1:A:288:THR:OG1	1:A:291:GLU:HG3	2.09	0.52
1:A:133:THR:HB	1:A:134:PRO:CD	2.39	0.52
1:A:73:PHE:CE1	1:A:135:GLU:HA	2.44	0.52
1:A:84:GLY:C	1:A:86:GLN:NE2	2.62	0.52
1:A:85:LEU:N	1:A:86:GLN:NE2	2.57	0.52
1:A:195:ASN:N	1:A:195:ASN:HD22	1.86	0.50
1:A:86:GLN:H	1:A:86:GLN:HE21	1.52	0.50

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:4:LEU:HD23	1:A:4:LEU:H	1.67	0.49
1:A:20:GLN:NE2	1:A:287:LYS:H	1.97	0.48
1:A:93:GLU:N	1:A:94:PRO:HD2	2.28	0.48
1:A:84:GLY:CA	1:A:86:GLN:NE2	2.77	0.47
1:A:10:VAL:HG12	1:A:11:GLU:O	2.14	0.47
1:A:188:GLU:H	1:A:222:GLN:HE22	1.62	0.47
1:A:34:ASP:O	1:A:39:TYR:HA	2.15	0.47
1:A:292:GLN:O	1:A:294:LEU:CD1	2.63	0.46
1:A:145:PRO:HD3	1:A:157:PHE:CZ	2.51	0.46
1:A:209:GLU:HB2	1:A:211:TRP:CE2	2.50	0.46
1:A:218:ALA:O	1:A:219:ASN:HB3	2.16	0.46
1:A:183:LYS:HG2	3:A:483:HOH:O	2.16	0.46
1:A:53:ILE:HG22	1:A:71:TYR:HB2	1.98	0.46
1:A:72:ARG:NH2	1:A:133:THR:O	2.48	0.45
1:A:163:MET:HA	1:A:167:GLU:OE2	2.17	0.45
1:A:184:ASN:HB2	2:A:296:HEM:O1A	2.16	0.44
1:A:86:GLN:H	1:A:86:GLN:CD	2.17	0.44
1:A:98:GLU:C	1:A:100:PRO:HD3	2.38	0.44
1:A:23:TYR:CD2	1:A:23:TYR:C	2.92	0.43
1:A:36:TYR:HB2	1:A:42:TYR:CE1	2.54	0.43
1:A:292:GLN:CB	1:A:294:LEU:HD13	2.49	0.43
1:A:243:LYS:HE3	1:A:243:LYS:H	1.83	0.43
1:A:292:GLN:HB2	1:A:294:LEU:HD13	2.02	0.42
1:A:267:GLU:OE1	1:A:271:GLU:OE1	2.37	0.42
1:A:35:GLU:CG	1:A:39:TYR:CE2	3.03	0.41
1:A:164:ASN:O	1:A:168:VAL:HG23	2.21	0.41
1:A:289:LEU:HD23	1:A:289:LEU:HA	1.85	0.41
2:A:296:HEM:HBB2	2:A:296:HEM:CMB	2.51	0.40
1:A:245:LEU:O	1:A:249:LYS:HG3	2.22	0.40
1:A:99:PHE:O	1:A:102:ILE:HG22	2.20	0.40
1:A:86:GLN:NE2	1:A:86:GLN:N	2.52	0.40
1:A:282:SER:CB	1:A:283:PRO:CD	2.99	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:431:HOH:O	3:A:526:HOH:O[4_566]	1.51	0.69
1:A:34:ASP:OD2	3:A:568:HOH:O[3_545]	2.08	0.12

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	291/296 (98%)	287 (99%)	4 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	241/254 (95%)	232 (96%)	9 (4%)	41	50

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	86	GLN
1	A	195	ASN
1	A	200	ASN
1	A	225	SER
1	A	237	SER
1	A	243	LYS
1	A	282	SER
1	A	287	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	20	GLN
1	A	24	ASN
1	A	60	HIS
1	A	86	GLN
1	A	87	ASN
1	A	141	ASN
1	A	195	ASN
1	A	200	ASN
1	A	208	ASN
1	A	222	GLN
1	A	292	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	HEM	A	296	1,3	30,50,50	2.29	7 (23%)	24,82,82	2.80	11 (45%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral

centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	296	1,3	-	0/10/54/54	0/0/8/8

All (7) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	296	HEM	C2D-C3D	-6.75	1.34	1.54
2	A	296	HEM	C3B-C4B	-5.47	1.47	1.51
2	A	296	HEM	C3D-C4D	-4.63	1.45	1.51
2	A	296	HEM	C2C-C1C	-3.76	1.45	1.52
2	A	296	HEM	CMA-C3A	2.18	1.56	1.51
2	A	296	HEM	C3C-CAC	2.52	1.56	1.51
2	A	296	HEM	C1C-NC	2.98	1.39	1.36

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	296	HEM	C3B-CAB-CBB	-4.29	117.87	124.46
2	A	296	HEM	CAA-C2A-C1A	-3.93	122.75	127.01
2	A	296	HEM	C3C-CAC-CBC	-3.70	118.78	124.46
2	A	296	HEM	CMA-C3A-C4A	-3.34	122.83	128.36
2	A	296	HEM	CMA-C3A-C2A	2.07	129.56	125.24
2	A	296	HEM	CMD-C2D-C3D	3.11	128.12	114.35
2	A	296	HEM	CAD-C3D-C4D	3.16	123.62	112.47
2	A	296	HEM	C2D-C3D-C4D	4.02	108.32	101.50
2	A	296	HEM	CMC-C2C-C3C	4.59	127.98	116.53
2	A	296	HEM	CMB-C2B-C3B	5.00	129.02	116.53
2	A	296	HEM	CAD-C3D-C2D	5.16	128.06	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	296	HEM	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.