



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 07:51 AM GMT

PDB ID : 3CEA
Title : Crystal structure of myo-inositol 2-dehydrogenase (NP_786804.1) from *Lactobacillus plantarum* at 2.40 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2008-02-28
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	2919 (2.40-2.40)
Clashscore	102246	3407 (2.40-2.40)
Ramachandran outliers	100387	3351 (2.40-2.40)
Sidechain outliers	100360	3352 (2.40-2.40)
RSRZ outliers	91569	2928 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 10991 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Myo-inositol 2-dehydrogenase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
1	A	342	Total	C 2626	N 1664	O 451	S 498	Se 3	10	0	2	0
1	B	342	Total	C 2629	N 1664	O 451	S 501	Se 3	10	0	1	0
1	C	339	Total	C 2593	N 1641	O 447	S 492	Se 3	10	0	1	0
1	D	339	Total	C 2609	N 1650	O 448	S 498	Se 3	10	0	1	0


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	LEADER SEQUENCE	UNP Q88S39
B	0	GLY	-	LEADER SEQUENCE	UNP Q88S39
C	0	GLY	-	LEADER SEQUENCE	UNP Q88S39
D	0	GLY	-	LEADER SEQUENCE	UNP Q88S39

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total Cl 1 1	0	0

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C₂₁H₂₇N₇O₁₄P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total		C	N	O	P	
			44		21	7	14	2	
3	B	1	Total		C	N	O	P	
			44		21	7	14	2	
3	C	1	Total		C	N	O	P	
			44		21	7	14	2	
3	D	1	Total		C	N	O	P	
			44		21	7	14	2	

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	C	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0
4	D	1	Total C O 4 2 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	88	Total O 89 89	0	1
5	B	85	Total O 85 85	0	0
5	C	81	Total O 82 82	0	1
5	D	89	Total O 89 89	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Myo-inositol 2-dehydrogenase




- Molecule 1: Myo-inositol 2-dehydrogenase




- Molecule 1: Myo-inositol 2-dehydrogenase



- Molecule 1: Myo-inositol 2-dehydrogenase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.33 Å 145.68 Å 100.21 Å 90.00° 91.92° 90.00°	Depositor
Resolution (Å)	29.45 – 2.40 29.45 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.0 (29.45-2.40) 99.1 (29.45-2.40)	Depositor EDS
R_{merge}	0.13	Depositor
R_{sym}	0.13	Depositor
$< I/\sigma(I) >$ ¹	2.12 (at 2.39 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R , R_{free}	0.193 , 0.243 0.196 , 0.244	Depositor DCC
R_{free} test set	2662 reflections (5.35%)	DCC
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.771	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 46.1	EDS
Estimated twinning fraction	0.041 for h,-k,-l	Xtriage
L-test for twinning ²	$< L > = 0.49$, $< L^2 > = 0.32$	Xtriage
Outliers	0 of 52383 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	10991	wwPDB-VP
Average B, all atoms (Å ²)	46.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $<|L|>$, $<|L^2|>$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, NAD, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.32	0/2672	0.47	0/3605
1	B	0.33	0/2672	0.47	0/3604
1	C	0.33	0/2636	0.46	0/3559
1	D	0.33	0/2651	0.47	0/3576
All	All	0.33	0/10631	0.46	0/14344

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2626	0	2558	14	0
1	B	2629	0	2566	16	0
1	C	2593	0	2512	14	0
1	D	2609	0	2538	25	0
2	B	1	0	0	0	0
3	A	44	0	26	1	0
3	B	44	0	26	1	0
3	C	44	0	26	0	0
3	D	44	0	26	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	C	4	0	6	0	0
4	D	8	0	12	2	0
5	A	89	0	0	0	0
5	B	85	0	0	0	0
5	C	82	0	0	0	0
5	D	89	0	0	0	0
All	All	10991	0	10296	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:91:MSE:HE2	1:C:117:ILE:HA	1.62	0.80
1:D:91:MSE:HE2	1:D:117:ILE:HA	1.61	0.80
1:B:91:MSE:HE2	1:B:117:ILE:HA	1.74	0.69
1:C:91:MSE:HE3	1:C:120:HIS:HB2	1.76	0.67
1:A:91:MSE:HE2	1:A:117:ILE:HA	1.81	0.62
1:D:143:ILE:HG23	1:D:148:ASP:HB2	1.82	0.62
1:A:155:MSE:HE3	1:A:255:VAL:HG22	1.83	0.60
1:D:26:LEU:HD23	1:D:35:LEU:HD13	1.84	0.59
1:D:158:TYR:CE1	1:D:240:ILE:HD12	2.39	0.57
1:B:158:TYR:CE1	1:B:240:ILE:HD12	2.40	0.57
1:A:340:LEU:HD13	1:D:340:LEU:HD13	1.88	0.54
1:C:8:LEU:HD12	1:C:305:VAL:HG13	1.89	0.54
1:D:158:TYR:HB2	1:D:252:GLU:HB3	1.89	0.54
1:D:283:ARG:HG2	4:D:402:EDO:H11	1.89	0.54
1:A:159:GLY:HA3	1:A:185:MSE:HE2	1.90	0.53
1:B:91:MSE:HE3	1:B:120:HIS:HB2	1.91	0.52
1:A:4:THR:HG23	1:A:5:ARG:H	1.75	0.51
1:B:229:LEU:HD11	1:B:239:LEU:HD11	1.93	0.50
1:B:17:ARG:NH1	3:B:400:NAD:O2N	2.42	0.50
1:A:333:GLN:O	1:A:337:ILE:HG12	2.13	0.49
1:B:333:GLN:O	1:B:337:ILE:HG12	2.13	0.49
1:A:209:SER:HB3	1:A:224:THR:HB	1.95	0.48
1:D:199:GLN:HG3	1:D:233:ASP:CG	2.34	0.48
1:C:27:VAL:HG21	1:C:53:LEU:HB3	1.95	0.48
1:A:154:TYR:CZ	1:D:240:ILE:HD11	2.48	0.48
1:B:143:ILE:HG23	1:B:148:ASP:HB2	1.94	0.48
1:B:340:LEU:HD13	1:C:340:LEU:HD13	1.95	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:158:TYR:HB2	1:A:252:GLU:HB3	1.95	0.47
1:B:268:PRO:HG2	1:B:285:SER:HB3	1.96	0.47
1:A:143:ILE:HG23	1:A:148:ASP:HB2	1.97	0.47
1:B:155:MSE:HE3	1:B:255:VAL:HG22	1.96	0.46
1:D:226:VAL:HG22	1:D:240:ILE:HG12	1.97	0.46
1:C:158:TYR:CE1	1:C:240:ILE:HD12	2.51	0.46
1:C:155:MSE:HE3	1:C:255:VAL:HG22	1.98	0.46
1:A:91:MSE:CE	1:A:120:HIS:HB2	2.46	0.45
1:A:17:ARG:NH1	3:A:400:NAD:O2N	2.49	0.45
1:D:87:THR:O	1:D:91:MSE:HG2	2.17	0.45
1:B:155:MSE:HE2	1:B:253:LEU:HD11	1.99	0.45
1:D:204:ALA:HB2	1:D:343:ILE:HD11	1.99	0.44
1:D:229:LEU:HB3	1:D:231:MSE:CE	2.47	0.44
1:C:254:GLU:OE2	1:C:256:MSE:HE3	2.18	0.44
1:D:88:ILE:HG12	1:D:116:VAL:HG21	1.99	0.43
1:A:186:ASN:HD21	1:A:240:ILE:C	2.21	0.43
1:B:83:HIS:HB2	1:B:84:PRO:HD3	2.00	0.43
1:A:240:ILE:HD11	1:D:154:TYR:CZ	2.54	0.43
1:D:209:SER:HB3	1:D:224:THR:HB	2.01	0.42
1:C:83:HIS:HB2	1:C:84:PRO:HD3	2.02	0.42
1:D:162:PRO:HA	1:D:245:ALA:O	2.20	0.42
1:B:77:VAL:HG13	1:B:100:GLU:HG3	2.02	0.42
1:D:281:VAL:HG12	4:D:402:EDO:H12	2.03	0.41
1:D:185:MSE:HE2	1:D:242:GLY:O	2.20	0.41
1:B:130:MSE:H	1:B:130:MSE:SE	2.53	0.41
1:D:229:LEU:HB3	1:D:231:MSE:HE1	2.03	0.41
1:D:182:PHE:CE1	1:D:227:ALA:HB2	2.56	0.41
1:C:71:ILE:HG13	1:C:95:LEU:HD11	2.03	0.41
1:C:101:LYS:HD3	1:C:187:ILE:HG21	2.02	0.41
1:D:283:ARG:HA	1:D:283:ARG:HD2	1.94	0.41
1:B:153:ILE:HD11	1:C:212:ALA:HB2	2.03	0.40
1:D:333:GLN:O	1:D:337:ILE:HG12	2.20	0.40
1:B:8:LEU:HD12	1:B:305:VAL:HG13	2.03	0.40
1:C:154:TYR:HA	1:C:236:ILE:O	2.22	0.40
1:D:26:LEU:CD1	1:D:301:VAL:HG11	2.52	0.40
1:D:144:VAL:HG11	1:D:152:ILE:HD11	2.04	0.40
1:C:130:MSE:SE	1:C:130:MSE:H	2.54	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	342/346 (99%)	335 (98%)	7 (2%)	0	100 100
1	B	341/346 (99%)	336 (98%)	5 (2%)	0	100 100
1	C	338/346 (98%)	331 (98%)	7 (2%)	0	100 100
1	D	338/346 (98%)	332 (98%)	6 (2%)	0	100 100
All	All	1359/1384 (98%)	1334 (98%)	25 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	272/275 (99%)	268 (98%)	4 (2%)	72 87
1	B	275/275 (100%)	271 (98%)	4 (2%)	72 87
1	C	268/275 (98%)	262 (98%)	6 (2%)	60 79
1	D	273/275 (99%)	269 (98%)	4 (2%)	72 87
All	All	1088/1100 (99%)	1070 (98%)	18 (2%)	68 85

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	26	LEU
1	A	57	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	101	LYS
1	A	130	MSE
1	B	26	LEU
1	B	57	THR
1	B	101	LYS
1	B	130	MSE
1	C	20	GLU
1	C	26	LEU
1	C	57	THR
1	C	101	LYS
1	C	130	MSE
1	C	244	HIS
1	D	57	THR
1	D	101	LYS
1	D	130	MSE
1	D	244	HIS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	186	ASN
1	B	186	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
3	NAD	A	400	-	38,48,48	1.67	3 (7%)	47,73,73	2.02	4 (8%)
3	NAD	B	400	-	38,48,48	1.64	3 (7%)	47,73,73	2.02	3 (6%)
3	NAD	C	400	-	38,48,48	1.66	3 (7%)	47,73,73	2.06	3 (6%)
4	EDO	C	401	-	3,3,3	0.42	0	2,2,2	0.57	0
3	NAD	D	400	-	38,48,48	1.68	3 (7%)	47,73,73	2.00	3 (6%)
4	EDO	D	401	-	3,3,3	0.46	0	2,2,2	0.41	0
4	EDO	D	402	-	3,3,3	0.50	0	2,2,2	0.27	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	400	-	-	0/22/62/62	0/5/5/5
3	NAD	B	400	-	-	0/22/62/62	0/5/5/5
3	NAD	C	400	-	-	0/22/62/62	0/5/5/5
4	EDO	C	401	-	-	0/1/1/1	0/0/0/0
3	NAD	D	400	-	-	0/22/62/62	0/5/5/5
4	EDO	D	401	-	-	0/1/1/1	0/0/0/0
4	EDO	D	402	-	-	0/1/1/1	0/0/0/0

All (12) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	400	NAD	C2A-N1A	2.50	1.38	1.33
3	B	400	NAD	C2A-N1A	2.56	1.38	1.33
3	D	400	NAD	C2A-N1A	2.63	1.38	1.33
3	A	400	NAD	C2A-N1A	2.70	1.39	1.33
3	B	400	NAD	C2A-N3A	3.59	1.38	1.32
3	C	400	NAD	C2A-N3A	3.73	1.38	1.32
3	A	400	NAD	C2A-N3A	3.75	1.38	1.32
3	D	400	NAD	C2A-N3A	3.76	1.38	1.32
3	B	400	NAD	O7N-C7N	7.96	1.41	1.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	400	NAD	O7N-C7N	8.05	1.41	1.24
3	C	400	NAD	O7N-C7N	8.05	1.41	1.24
3	D	400	NAD	O7N-C7N	8.15	1.41	1.24

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	400	NAD	N3A-C2A-N1A	-12.05	119.67	128.89
3	C	400	NAD	N3A-C2A-N1A	-11.81	119.86	128.89
3	A	400	NAD	N3A-C2A-N1A	-11.65	119.97	128.89
3	D	400	NAD	N3A-C2A-N1A	-11.56	120.04	128.89
3	A	400	NAD	PN-O3-PA	-3.85	121.92	132.73
3	D	400	NAD	PN-O3-PA	-3.83	121.96	132.73
3	C	400	NAD	PN-O3-PA	-3.67	122.42	132.73
3	B	400	NAD	PN-O3-PA	-3.50	122.90	132.73
3	A	400	NAD	C3N-C7N-N7N	2.04	120.05	117.82
3	B	400	NAD	O4D-C1D-N1N	2.19	110.54	108.13
3	A	400	NAD	O4D-C1D-N1N	2.80	111.20	108.13
3	D	400	NAD	O4D-C1D-N1N	2.98	111.41	108.13
3	C	400	NAD	O4D-C1D-N1N	3.94	112.46	108.13

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	400	NAD	1	0
3	B	400	NAD	1	0
4	D	402	EDO	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	332/346 (95%)	0.59	27 (8%) 15 14	36, 46, 57, 64	0
1	B	332/346 (95%)	0.51	22 (6%) 22 22	37, 46, 57, 61	0
1	C	329/346 (95%)	0.71	31 (9%) 11 10	35, 46, 58, 61	0
1	D	329/346 (95%)	0.66	33 (10%) 9 9	36, 46, 58, 63	0
All	All	1322/1384 (95%)	0.62	113 (8%) 13 13	35, 46, 57, 64	0

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	67	ASP	5.2
1	C	107	PHE	4.9
1	C	51	ASN	4.7
1	C	44	ASN	4.7
1	D	217	ALA	4.6
1	D	216	LEU	4.5
1	D	173	ALA	4.4
1	B	4	THR	4.4
1	D	218	ASP	4.0
1	B	345	LEU	4.0
1	C	52	GLU	3.9
1	D	167	GLU	3.9
1	A	219	ILE	3.7
1	D	48	TRP	3.6
1	B	58	THR	3.6
1	B	344	GLN	3.5
1	A	108	ASN	3.5
1	C	76	ILE	3.5
1	A	44	ASN	3.4
1	D	215	GLN	3.4
1	B	52	GLU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	80	THR	3.4
1	D	51	ASN	3.3
1	B	50	LYS	3.3
1	B	51	ASN	3.3
1	A	175	GLU	3.2
1	B	107	PHE	3.2
1	D	50	LYS	3.2
1	B	5	ARG	3.2
1	B	76	ILE	3.2
1	A	345	LEU	3.1
1	A	107	PHE	3.1
1	D	94	GLY	3.0
1	C	344	GLN	3.0
1	D	43	SER	3.0
1	A	76	ILE	3.0
1	D	219	ILE	2.9
1	A	237	ALA	2.9
1	A	174	THR	2.9
1	A	127[A]	SER	2.8
1	B	237	ALA	2.8
1	D	76	ILE	2.8
1	A	5	ARG	2.8
1	D	174	THR	2.8
1	A	77	VAL	2.8
1	A	128	GLY	2.8
1	B	215	GLN	2.8
1	C	237	ALA	2.7
1	A	159	GLY	2.7
1	C	187	ILE	2.7
1	A	43	SER	2.6
1	B	128	GLY	2.6
1	C	108	ASN	2.5
1	C	111	ASP	2.5
1	C	14	GLY	2.5
1	A	99	CYS	2.5
1	C	56	GLU	2.5
1	A	238	THR	2.5
1	C	315	GLU	2.5
1	C	12	ILE	2.4
1	B	43	SER	2.4
1	C	310	VAL	2.4
1	C	238	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	240	ILE	2.4
1	C	122	ASN	2.4
1	D	186	ASN	2.4
1	B	127[A]	SER	2.4
1	B	42	ASP	2.3
1	D	344	GLN	2.3
1	D	158	TYR	2.3
1	C	251	VAL	2.3
1	D	156	ARG	2.3
1	A	121	PRO	2.3
1	A	64	ASP	2.3
1	B	311	GLY	2.3
1	D	56	GLU	2.3
1	B	64	ASP	2.3
1	C	177	ASP	2.3
1	C	115	LYS	2.3
1	C	239	LEU	2.3
1	C	36	VAL	2.3
1	A	106	ASP	2.3
1	A	239	LEU	2.3
1	D	160	ILE	2.2
1	C	7	PRO	2.2
1	D	157	GLY	2.2
1	D	54	GLY	2.2
1	A	172	PHE	2.2
1	D	119	SER	2.1
1	C	157	GLY	2.1
1	C	164	SER	2.1
1	B	26	LEU	2.1
1	B	99	CYS	2.1
1	C	101	LYS	2.1
1	C	156	ARG	2.1
1	A	218	ASP	2.1
1	C	218	ASP	2.1
1	D	88	ILE	2.1
1	D	336	ASN	2.1
1	D	49	ALA	2.1
1	D	145	ASP	2.1
1	D	177	ASP	2.1
1	C	173	ALA	2.1
1	B	108	ASN	2.1
1	A	169	PHE	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	72	ASP	2.1
1	A	318	VAL	2.0
1	D	18	LEU	2.0
1	D	214	PRO	2.0
1	D	42	ASP	2.0
1	A	170	THR	2.0
1	B	48	TRP	2.0
1	C	48	TRP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	EDO	C	401	4/4	0.78	0.19	0.92	57,58,58,60	0
4	EDO	D	402	4/4	0.87	0.15	0.22	44,45,45,46	0
4	EDO	D	401	4/4	0.96	0.14	-0.25	45,45,46,46	0
3	NAD	C	400	44/44	0.91	0.16	-0.83	49,60,64,64	0
3	NAD	A	400	44/44	0.93	0.15	-1.06	46,49,56,57	0
3	NAD	D	400	44/44	0.94	0.12	-1.21	40,46,52,52	0
3	NAD	B	400	44/44	0.94	0.13	-1.48	40,43,50,51	0
2	CL	B	346	1/1	0.92	0.10	-	52,52,52,52	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.