



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:51 AM GMT

PDB ID : 3CES
Title : Crystal Structure of E.coli MnmG (GidA), a Highly-Conserved tRNA Modifying Enzyme
Authors : Shi, R.; Matte, A.; Cygler, M.; Montreal-Kingston Bacterial Structural Genomics Initiative (BSGI)
Deposited on : 2008-02-29
Resolution : 2.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3386 (2.44-2.40)
Clashscore	102246	3897 (2.44-2.40)
Ramachandran outliers	100387	3837 (2.44-2.40)
Sidechain outliers	100360	3838 (2.44-2.40)
RSRZ outliers	91569	3396 (2.44-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	651	 62% 15% 18%
1	B	651	 62% 15% 20%
1	C	651	 59% 17% 21%
1	D	651	 59% 18% 22%

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 15980 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called tRNA uridine 5-carboxymethylaminomethyl modification enzyme gidA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	531	4098	2562	739	778	19	0	1	0
1	B	519	3895	2447	696	735	17	0	1	0
1	C	516	3884	2433	699	734	18	0	0	0
1	D	507	3771	2370	678	705	18	0	1	0

There are 88 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	EXPRESSION TAG	UNP P0A6U3
A	-20	GLY	-	EXPRESSION TAG	UNP P0A6U3
A	-19	SER	-	EXPRESSION TAG	UNP P0A6U3
A	-18	SER	-	EXPRESSION TAG	UNP P0A6U3
A	-17	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-16	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-15	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-14	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-13	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-12	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-11	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-10	HIS	-	EXPRESSION TAG	UNP P0A6U3
A	-9	SER	-	EXPRESSION TAG	UNP P0A6U3
A	-8	SER	-	EXPRESSION TAG	UNP P0A6U3
A	-7	GLY	-	EXPRESSION TAG	UNP P0A6U3
A	-6	LEU	-	EXPRESSION TAG	UNP P0A6U3
A	-5	VAL	-	EXPRESSION TAG	UNP P0A6U3
A	-4	PRO	-	EXPRESSION TAG	UNP P0A6U3
A	-3	ARG	-	EXPRESSION TAG	UNP P0A6U3
A	-2	GLY	-	EXPRESSION TAG	UNP P0A6U3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	SER	-	EXPRESSION TAG	UNP P0A6U3
A	0	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-21	MET	-	EXPRESSION TAG	UNP P0A6U3
B	-20	GLY	-	EXPRESSION TAG	UNP P0A6U3
B	-19	SER	-	EXPRESSION TAG	UNP P0A6U3
B	-18	SER	-	EXPRESSION TAG	UNP P0A6U3
B	-17	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-16	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-15	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-14	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-13	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-12	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-11	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-10	HIS	-	EXPRESSION TAG	UNP P0A6U3
B	-9	SER	-	EXPRESSION TAG	UNP P0A6U3
B	-8	SER	-	EXPRESSION TAG	UNP P0A6U3
B	-7	GLY	-	EXPRESSION TAG	UNP P0A6U3
B	-6	LEU	-	EXPRESSION TAG	UNP P0A6U3
B	-5	VAL	-	EXPRESSION TAG	UNP P0A6U3
B	-4	PRO	-	EXPRESSION TAG	UNP P0A6U3
B	-3	ARG	-	EXPRESSION TAG	UNP P0A6U3
B	-2	GLY	-	EXPRESSION TAG	UNP P0A6U3
B	-1	SER	-	EXPRESSION TAG	UNP P0A6U3
B	0	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-21	MET	-	EXPRESSION TAG	UNP P0A6U3
C	-20	GLY	-	EXPRESSION TAG	UNP P0A6U3
C	-19	SER	-	EXPRESSION TAG	UNP P0A6U3
C	-18	SER	-	EXPRESSION TAG	UNP P0A6U3
C	-17	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-16	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-15	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-14	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-13	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-12	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-11	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-10	HIS	-	EXPRESSION TAG	UNP P0A6U3
C	-9	SER	-	EXPRESSION TAG	UNP P0A6U3
C	-8	SER	-	EXPRESSION TAG	UNP P0A6U3
C	-7	GLY	-	EXPRESSION TAG	UNP P0A6U3
C	-6	LEU	-	EXPRESSION TAG	UNP P0A6U3
C	-5	VAL	-	EXPRESSION TAG	UNP P0A6U3
C	-4	PRO	-	EXPRESSION TAG	UNP P0A6U3


Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-3	ARG	-	EXPRESSION TAG	UNP P0A6U3
C	-2	GLY	-	EXPRESSION TAG	UNP P0A6U3
C	-1	SER	-	EXPRESSION TAG	UNP P0A6U3
C	0	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-21	MET	-	EXPRESSION TAG	UNP P0A6U3
D	-20	GLY	-	EXPRESSION TAG	UNP P0A6U3
D	-19	SER	-	EXPRESSION TAG	UNP P0A6U3
D	-18	SER	-	EXPRESSION TAG	UNP P0A6U3
D	-17	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-16	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-15	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-14	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-13	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-12	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-11	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-10	HIS	-	EXPRESSION TAG	UNP P0A6U3
D	-9	SER	-	EXPRESSION TAG	UNP P0A6U3
D	-8	SER	-	EXPRESSION TAG	UNP P0A6U3
D	-7	GLY	-	EXPRESSION TAG	UNP P0A6U3
D	-6	LEU	-	EXPRESSION TAG	UNP P0A6U3
D	-5	VAL	-	EXPRESSION TAG	UNP P0A6U3
D	-4	PRO	-	EXPRESSION TAG	UNP P0A6U3
D	-3	ARG	-	EXPRESSION TAG	UNP P0A6U3
D	-2	GLY	-	EXPRESSION TAG	UNP P0A6U3
D	-1	SER	-	EXPRESSION TAG	UNP P0A6U3
D	0	HIS	-	EXPRESSION TAG	UNP P0A6U3

- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	147	Total O 147 147	0	0
2	B	114	Total O 114 114	0	0
2	C	41	Total O 41 41	0	0
2	D	30	Total O 30 30	0	0



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	85.90Å 144.33Å 147.75Å 90.00° 106.63° 90.00°	Depositor
Resolution (Å)	141.42 – 2.41 47.75 – 2.41	Depositor EDS
% Data completeness (in resolution range)	97.4 (141.42-2.41) 97.4 (47.75-2.41)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.08 (at 2.42Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R, R_{free}	0.201 , 0.238 0.201 , 0.237	Depositor DCC
R_{free} test set	6511 reflections (5.31%)	DCC
Wilson B-factor (Å ²)	41.1	Xtriage
Anisotropy	0.531	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 52.7	EDS
Estimated twinning fraction	0.014 for h,-k,-h-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 129227 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	15980	wwPDB-VP
Average B, all atoms (Å ²)	54.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.01% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.78	2/4174 (0.0%)	0.87	13/5651 (0.2%)
1	B	0.68	0/3970	0.85	14/5391 (0.3%)
1	C	0.54	1/3949 (0.0%)	0.85	10/5354 (0.2%)
1	D	0.47	0/3843	0.84	6/5213 (0.1%)
All	All	0.63	3/15936 (0.0%)	0.85	43/21609 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	1	1
1	C	0	2
1	D	0	3
All	All	1	6

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	0	HIS	C-N	14.28	1.66	1.34
1	C	500	THR	C-N	-6.51	1.19	1.34
1	A	506	GLU	CB-CG	-5.92	1.40	1.52

The worst 5 of 43 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	487	PRO	O-C-N	-35.11	66.53	122.70
1	C	500	THR	O-C-N	-32.11	71.33	122.70
1	D	487	PRO	CA-C-N	15.40	151.09	117.20
1	A	447	ARG	NE-CZ-NH2	-9.57	115.52	120.30
1	C	500	THR	CA-C-N	-8.81	97.82	117.20

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	B	348	PHE	CA

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	89	GLY	Peptide
1	C	500	THR	Mainchain
1	C	89	GLY	Peptide
1	D	487	PRO	Mainchain
1	D	89	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4098	0	4061	95	0
1	B	3895	0	3741	71	0
1	C	3884	0	3757	78	0
1	D	3771	0	3604	73	0
2	A	147	0	0	5	0
2	B	114	0	0	2	0
2	C	41	0	0	2	0
2	D	30	0	0	2	0
All	All	15980	0	15163	309	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 309 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:230:MET:HE2	1:C:230:MET:HA	1.25	1.12
1:D:154:THR:HG21	2:D:647:HOH:O	1.51	1.10
1:B:49:PRO:HB3	1:B:101:ARG:NH1	1.67	1.09
1:C:288:ALA:O	1:C:289:ASP:HB2	1.52	1.03

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:90:PRO:HB2	1:D:440:ARG:HD2	1.39	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	526/651 (81%)	507 (96%)	14 (3%)	5 (1%)	19	27
1	B	514/651 (79%)	476 (93%)	33 (6%)	5 (1%)	19	27
1	C	504/651 (77%)	461 (92%)	28 (6%)	15 (3%)	5	4
1	D	496/651 (76%)	466 (94%)	22 (4%)	8 (2%)	12	15
All	All	2040/2604 (78%)	1910 (94%)	97 (5%)	33 (2%)	12	15

5 of 33 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	160	ASP
1	B	348	PHE
1	B	437	LEU
1	B	504	SER
1	C	279	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	429/535 (80%)	389 (91%)	40 (9%)	11	16
1	B	383/535 (72%)	349 (91%)	34 (9%)	12	17
1	C	389/535 (73%)	361 (93%)	28 (7%)	18	27
1	D	365/535 (68%)	339 (93%)	26 (7%)	18	28
All	All	1566/2140 (73%)	1438 (92%)	128 (8%)	14	21

5 of 128 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	204	ARG
1	B	413	LEU
1	D	227	PHE
1	B	227	PHE
1	B	314	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 52 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	323	GLN
1	C	123	GLN
1	D	310	ASN
1	B	383	GLN
1	C	38	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	531/651 (81%)	0.07	4 (0%) 87 87	20, 34, 63, 86	0
1	B	519/651 (79%)	0.43	43 (8%) 14 13	23, 38, 125, 136	0
1	C	516/651 (79%)	0.71	56 (10%) 7 7	36, 58, 105, 148	4 (0%)
1	D	507/651 (77%)	0.75	66 (13%) 5 4	40, 61, 128, 160	1 (0%)
All	All	2073/2604 (79%)	0.49	169 (8%) 14 14	20, 49, 114, 160	5 (0%)

The worst 5 of 169 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	504	SER	28.3
1	C	503	LEU	22.1
1	C	502	PRO	15.0
1	D	259	LEU	8.7
1	D	526	LEU	7.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers

There are no such residues in this entry.