wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 07:03 PM BST

PDB ID	:	2CH0
Title	:	SOLUTION STRUCTURE OF THE HUMAN MAN1 C-TERMINAL DOMAIN (RESIDUES 655-775)
Authors	:	Caputo, S.; Couprie, J.; Duband-Goulet, I.; Lin, F.; Braud, S.; Gondry, M.; Worman, H.J.; Gilquin, B.; Zinn-Justin, S.
Deposited on	:	2006-03-10

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	133	41% 18% • 38%

2 Ensemble composition and analysis (i)

This entry contains 20 models. Model 7 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:18-A:100 (83)	0.62	7

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 3, 4, 5, 6, 7, 8, 9, 13, 14, 15, 18, 19
2	10, 11, 12, 16, 17
Single-model clusters	20

3 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 2225 atoms, of which 1107 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called INNER NUCLEAR MEMBRANE PROTEIN MAN1.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	133	2225	702	1107	212	198	6	0

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: INNER NUCLEAR MEMBRANE PROTEIN MAN1

Chain A: 41% 18% • 38%

G1 S2 P3 E4 F5 R6 W7 T8 K9 E10 E11 E12 E13 T14 R15 Q16 M17 Y18 D19 M20 K23 I24 V27 L28 H31 C35 K39 D40 L41 Q42 M45 P48 H49 V50 R51 L54 I55 Q56 P57 R60 K61 V81 T85 I88 W96 R97 W98

I99 Q100 P101 S102 A103 S104 C105 D106 K107 I108 L109 V110 I111 P112 S113 K114 V115 W116 Q117 G118 Q119 A120 F121 H122 L123 D124 R125 R126 L127 E128 R129 P130 H131 R132 D133

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 7. Colouring as in section 4.1 above.

- Molecule 1: INNER NUCLEAR MEMBRANE PROTEIN MAN1

Chain A: 43% 16% • • 38%

G1 S2 P3 E4 F5 R6 W7 T8 K9 E10 E11 E12 E13 T14 R15 Q16 M17 Y18 D19 M20 V21 I24 V27 L28 H31 K39 D40 L41 Q42 M45 P48 H49 V50 L54 I55 Q56 P57 R60 K61 W67 V71 V81 R86 R87 I88 F93

W98 P101 S102 A103 S104 C105 D106 K107 I108 L109 V110 I111 P112 S113 K114 V115 W116 Q117 G118 Q119 A120 F121 H122 L123 D124 R125 R126 L127 E128 R129 P130 H131 R132 D133

5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *DISTANCE GEOMETRY, SIMULATED ANNEALING.*

Of the 1000 calculated structures, 20 were deposited, based on the following criterion: *BACK CALCULATED DATA AGREE WITH EXPERIMENTAL NOESY SPECTRUM, STRUCTURES WITH ACCEPTABLE COVALENT GEOMETRY, STRUCTURES WITH FAVORABLE NON-BOND ENERGY, STRUCTURES WITH THE LEAST RESTRAINT VIOLATIONS, STRUCTURES WITH THE LOWEST ENERGY.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS 1.0	refinement	
XWINNMR 2.5	structure solution	
NMRPIPE 2.0	structure solution	
FELIX 2000.1	structure solution	
CNS 1.0	structure solution	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 6919
Number of chemical shift lists	1
Total number of shifts	1220
Number of shifts mapped to atoms	1220
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.14±0.02	0±0/715 (0.0±0.0%)	1.62±0.06	10±3/968 (1.1±0.3%)
All	All	1.15	0/14300 (0.0%)	1.62	206/19360 (1.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.1±0.3	1.1±1.2
All	All	2	23

There are no bond-length outliers.

5 of 71 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
1	A	98	TRP	CB-CG-CD2	-17.75	103.52	126.60	2	6
1	A	18	TYR	CB-CG-CD1	9.55	126.73	121.00	8	2
1	A	98	TRP	CB-CG-CD1	-9.47	114.69	127.00	6	4
1	A	73	PHE	CB-CG-CD1	9.22	127.26	120.80	20	3
1	A	35	CYS	CA-CB-SG	8.81	129.86	114.00	3	3

All unique chiral outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Models (Total)
1	A	61	LYS	CA	1
1	A	98	TRP	CA	1

5 of 6 unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	44	TYR	Sidechain	5
1	A	18	TYR	Sidechain	5
1	A	49	HIS	Sidechain	5
1	A	42	GLN	Peptide	4
1	A	31	HIS	Sidechain	3

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	698	701	701	6±3
All	All	13960	14020	14020	118

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

5 of 76 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:27:VAL:HG11	1:A:53:SER:HB3	0.79	1.53	15	2
1:A:81:VAL:HG22	1:A:98:TRP:HB2	0.74	1.60	18	3
1:A:49:HIS:CE1	1:A:50:VAL:HG12	0.72	2.19	6	2
1:A:81:VAL:HG12	1:A:98:TRP:HB3	0.70	1.63	17	4
1:A:27:VAL:HG21	1:A:50:VAL:HB	0.66	1.67	16	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	83/133 (62%)	73±2 (88±2%)	7±1 (9±2%)	3±1 (3±1%)	8 38
All	All	1660/2660 (62%)	1457 (88%)	147 (9%)	56 (3%)	8 38

5 of 8 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	42	GLN	20
1	A	57	PRO	20
1	A	78	GLU	6
1	A	19	ASP	4
1	A	39	LYS	3

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	76/122 (62%)	58±3 (76±4%)	18±3 (24±4%)	3 28
All	All	1520/2440 (62%)	1154 (76%)	366 (24%)	3 28

5 of 57 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	39	LYS	20
1	A	54	LEU	20
1	A	60	ARG	20
1	A	88	ILE	20
1	A	23	LYS	17

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

There are no ligands in this entry.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 61% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 6919

Chemical shift list name: *assigned_ chem_ shift_ list_ 1*

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1220
Number of shifts mapped to atoms	1220
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.1.2 Chemical shift referencing (i)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction ± precision, *ppm*	Suggested action
$^{13}C_{\alpha}$	95	-0.31 ± 0.15	None needed (< 0.5 ppm)
$^{13}C_{\beta}$	87	0.21 ± 0.16	None needed (< 0.5 ppm)
$^{13}C'$	90	-0.18 ± 0.12	None needed (< 0.5 ppm)
^{15}N	89	0.17 ± 0.52	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 869 atoms were assigned a chemical shift out of a possible 1126. 12 out of 13 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^{1}H	^{13}C	^{15}N
Backbone	366/407 (90%)	148/162 (91%)	148/166 (89%)	70/79 (89%)
Sidechain	448/628 (71%)	283/370 (76%)	159/218 (73%)	6/40 (15%)

Continued on next page...

Continued from previous page...

	Total	^{1}H	^{13}C	^{15}N
Aromatic	55/91 (60%)	29/48 (60%)	23/37 (62%)	3/6 (50%)
Overall	869/1126 (77%)	460/580 (79%)	330/421 (78%)	79/125 (63%)

7.1.4 Statistically unusual chemical shifts (i)

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

Mol	Chain	Res	Type	Atom	Shift, *ppm*	Expected range, *ppm*	Z-score
1	A	96	TRP	CZ3	131.45	129.20 – 113.60	6.4
1	A	47	ILE	HG12	-1.23	3.27 – -0.73	-6.2
1	A	67	TRP	CZ3	113.32	129.20 – 113.60	-5.2

7.1.5 Random Coil Index (RCI) plots (i)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A: