



wwPDB EM Map/Model Validation Report ⓘ

Apr 10, 2016 – 01:57 PM BST

PDB ID : 4CK5
EMDB ID: : EMD-2537
Title : Pseudo-atomic model of microtubule-bound human kinesin-5 motor domain in the ADP state, based on cryo-electron microscopy experiment.
Authors : Goulet, A.; Major, J.; Jun, Y.; Gross, S.; Rosenfeld, S.; Moores, C.
Deposited on : 2013-12-30
Resolution : 10.00 Å(reported)
Based on PDB ID : 3HQD,1JFF

This is a wwPDB EM Map/Model Validation Report for a publicly released PDB/EMDB entry.
For rigid body fitted models, validation errors reported here could stem from errors in the original structure(s) used in the fitting.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/EMValidationReportHelp>

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et. al. (1996)
Validation Pipeline (wwPDB-VP) : trunk27241


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY




The reported resolution of this entry is 10.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	114402	924

The table below summarises the geometric issues observed across the polymeric chains. The red, orange, yellow and green segments on the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	452	 91% 9%
2	B	445	 95% .
3	C	373	 92% . 6%

2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 1338 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TUBULIN ALPHA-1D CHAIN.

Mol	Chain	Residues	Atoms		AltConf	Trace
1	A	412	Total	C	0	412
			412	412		

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	7	ILE	VAL	CONFLICT	UNP Q2HJ86
A	114	ILE	LEU	CONFLICT	UNP Q2HJ86
A	136	SER	LEU	CONFLICT	UNP Q2HJ86
A	137	VAL	ILE	CONFLICT	UNP Q2HJ86
A	358	GLU	ASP	CONFLICT	UNP Q2HJ86
A	437	VAL	MET	CONFLICT	UNP Q2HJ86

- Molecule 2 is a protein called TUBULIN BETA-2B CHAIN.

Mol	Chain	Residues	Atoms		AltConf	Trace
2	B	426	Total	C	0	426
			426	426		

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	55	ALA	THR	CONFLICT	UNP Q6B856
B	170	VAL	MET	CONFLICT	UNP Q6B856
B	296	ALA	SER	CONFLICT	UNP Q6B856
B	316	VAL	ILE	CONFLICT	UNP Q6B856


- Molecule 3 is a protein called KINESIN-LIKE PROTEIN KIF11.

Mol	Chain	Residues	Atoms		AltConf	Trace
3	C	350	Total	C	0	350
			350	350		

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	43	SER	CYS	ENGINEERED MUTATION	UNP P52732
C	87	ALA	CYS	ENGINEERED MUTATION	UNP P52732
C	99	ALA	CYS	ENGINEERED MUTATION	UNP P52732
C	126	CYS	THR	ENGINEERED MUTATION	UNP P52732
C	361	HIS	-	EXPRESSION TAG	UNP P52732
C	362	HIS	-	EXPRESSION TAG	UNP P52732
C	363	HIS	-	EXPRESSION TAG	UNP P52732
C	364	HIS	-	EXPRESSION TAG	UNP P52732
C	365	HIS	-	EXPRESSION TAG	UNP P52732
C	366	HIS	-	EXPRESSION TAG	UNP P52732

- Molecule 4 is GUANOSINE-5'-TRIPHOSPHATE (three-letter code: GTP) (formula: $C_{10}H_{16}N_5O_{14}P_3$).



Mol	Chain	Residues	Atoms					AltConf
4	A	1	Total	C	N	O	P	0
			32	10	5	14	3	

- Molecule 5 is GUANOSINE-5'-DIPHOSPHATE (three-letter code: GDP) (formula: $C_{10}H_{15}N_5O_{11}P_2$).



Mol	Chain	Residues	Atoms					AltConf
5	B	1	Total 28	C 10	N 5	O 11	P 2	0

- Molecule 6 is TAXOL (three-letter code: TA1) (formula: $C_{47}H_{51}NO_{14}$).




Mol	Chain	Residues	Atoms				AltConf
6	B	1	Total	C	N	O	0
			62	47	1	14	

- Molecule 7 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
7	C	1	Total	Mg	0
			1	1	

- Molecule 8 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).




Mol	Chain	Residues	Atoms					AltConf
8	C	1	Total	C	N	O	P	0
			27	10	5	10	2	

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: TUBULIN ALPHA-1D CHAIN

Chain A:  91% 9%




- Molecule 2: TUBULIN BETA-2B CHAIN

Chain B:  95% .



- Molecule 3: KINESIN-LIKE PROTEIN KIF11

Chain C:  92% . 6%



4 Experimental information

Property	Value	Source
Reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	Depositor
Number of images	Not provided	Depositor
Resolution determination method	Not provided	Depositor
CTF correction method	FREALIGN, Not provided	Depositor
Microscope	FEI TECNAI F20	Depositor
Voltage (kV)	200	Depositor
Electron dose ($e^-/\text{\AA}^2$)	18	Depositor
Minimum defocus (nm)	700	Depositor
Maximum defocus (nm)	2400	Depositor
Magnification	68000	Depositor
Image detector	GATAN ULTRASCAN 4000 (4K X 4K)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GDP, GTP, MG, TA1, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

There are no protein, RNA or DNA chains available to summarize Z scores of covalent bonds and angles.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	412	0	0	1	0
2	B	426	0	0	1	0
3	C	350	0	0	3	0
4	A	32	0	12	0	0
5	B	28	0	12	0	0
6	B	62	0	51	3	0
7	C	1	0	0	0	0
8	C	27	0	12	0	0
All	All	1338	0	87	8	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (8) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:402:ARG:CA	1:A:403:ALA:CA	2.54	0.85
3:C:27:PRO:CA	3:C:28:PHE:CA	2.65	0.74
6:B:1187:TA1:H463	6:B:1187:TA1:H261	1.80	0.63
3:C:303:PRO:CA	3:C:304:TYR:CA	2.81	0.58
2:B:284:LEU:CA	2:B:285:THR:CA	2.86	0.53
3:C:130:ASP:CA	3:C:131:PRO:CA	2.92	0.47
6:B:1187:TA1:H463	6:B:1187:TA1:C26	2.46	0.46
6:B:1187:TA1:H101	6:B:1187:TA1:C25	2.52	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein backbone outliers to report in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	GTP	A	1185	-	26,34,34	1.36	2 (7%)	29,54,54	2.28	4 (13%)
5	GDP	B	1186	-	24,30,30	2.64	8 (33%)	26,47,47	3.29	9 (34%)
6	TA1	B	1187	-	68,68,68	1.92	19 (27%)	102,105,105	1.30	8 (7%)
8	ADP	C	1184	7	24,29,29	1.23	3 (12%)	23,45,45	2.34	6 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GTP	A	1185	-	-	0/18/38/38	0/3/3/3
5	GDP	B	1186	-	-	0/12/32/32	0/3/3/3
6	TA1	B	1187	-	-	0/41/127/127	0/5/7/7
8	ADP	C	1184	7	-	0/12/32/32	0/3/3/3

All (32) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	B	1187	TA1	C08-C07	-4.84	1.26	1.38
5	B	1186	GDP	PB-O2B	-4.15	1.40	1.54
8	C	1184	ADP	C8-N7	-3.26	1.28	1.34
6	B	1187	TA1	C04-C03	-2.42	1.44	1.49
8	C	1184	ADP	C5-C4	-2.10	1.35	1.40
6	B	1187	TA1	C10-C02	2.02	1.62	1.57
6	B	1187	TA1	C37-C29	2.06	1.54	1.51
4	A	1185	GTP	O4'-C1'	2.10	1.44	1.41
5	B	1186	GDP	O3'-C3'	2.12	1.48	1.43
6	B	1187	TA1	C18-C20	2.14	1.62	1.56
6	B	1187	TA1	C11-C10	2.18	1.61	1.55
6	B	1187	TA1	C16-C15	2.20	1.56	1.52
6	B	1187	TA1	C01-C45	2.22	1.66	1.56
6	B	1187	TA1	C26-C25	2.28	1.56	1.51
6	B	1187	TA1	C43-C26	2.46	1.58	1.52
5	B	1186	GDP	C5-C4	2.52	1.46	1.40
8	C	1184	ADP	C2'-C3'	2.52	1.60	1.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	B	1186	GDP	PB-O3B	2.64	1.63	1.54
6	B	1187	TA1	C43-C01	2.79	1.60	1.54
6	B	1187	TA1	C46-C45	2.80	1.59	1.53
6	B	1187	TA1	C25-C24	3.16	1.39	1.34
5	B	1186	GDP	C8-N7	3.33	1.41	1.34
6	B	1187	TA1	O02-C03	3.37	1.41	1.34
6	B	1187	TA1	C36-C31	3.41	1.44	1.39
6	B	1187	TA1	C45-C24	3.45	1.61	1.54
5	B	1186	GDP	O6-C6	4.03	1.34	1.24
6	B	1187	TA1	C18-C10	4.04	1.68	1.57
6	B	1187	TA1	C05-C04	4.63	1.46	1.39
4	A	1185	GTP	C6-N1	4.81	1.41	1.33
6	B	1187	TA1	C06-C05	5.59	1.49	1.38
5	B	1186	GDP	O4'-C1'	5.93	1.49	1.41
5	B	1186	GDP	C2-N1	7.24	1.49	1.35

All (27) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	B	1186	GDP	C6-C5-C4	-9.91	109.54	120.86
4	A	1185	GTP	C5-C6-N1	-7.77	113.36	123.52
8	C	1184	ADP	O2A-PA-O3A	-6.35	78.07	105.27
5	B	1186	GDP	N2-C2-N1	-5.75	107.72	117.20
5	B	1186	GDP	N3-C2-N1	-5.33	120.31	127.56
6	B	1187	TA1	C06-C05-C04	-4.70	114.75	120.35
6	B	1187	TA1	C05-C04-C03	-3.91	111.56	120.38
8	C	1184	ADP	N3-C2-N1	-3.60	126.04	128.87
4	A	1185	GTP	N3-C2-N1	-3.44	122.88	127.56
5	B	1186	GDP	C1'-N9-C4	-2.66	123.84	126.81
6	B	1187	TA1	O04-C11-C14	-2.50	101.73	108.08
8	C	1184	ADP	C1'-N9-C4	-2.24	124.31	126.81
4	A	1185	GTP	C6-C5-C4	-2.15	118.41	120.86
5	B	1186	GDP	C6-N1-C2	2.02	118.25	115.88
8	C	1184	ADP	O5'-C5'-C4'	2.08	116.60	109.09
5	B	1186	GDP	O2'-C2'-C3'	2.25	119.14	111.86
6	B	1187	TA1	O01-C01-C43	2.49	113.33	106.82
6	B	1187	TA1	C17-C18-C20	2.55	109.77	102.14
6	B	1187	TA1	C45-C01-C02	3.11	115.23	111.64
5	B	1186	GDP	C2'-C3'-C4'	3.20	109.19	102.64
6	B	1187	TA1	C09-C04-C03	3.58	128.45	120.38
5	B	1186	GDP	C4'-O4'-C1'	4.07	113.96	109.64
8	C	1184	ADP	O2A-PA-O1A	4.26	134.72	112.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
6	B	1187	TA1	C07-C08-C09	5.11	127.30	120.20
8	C	1184	ADP	C4'-O4'-C1'	5.24	115.20	109.64
4	A	1185	GTP	C6-N1-C2	7.45	124.61	115.88
5	B	1186	GDP	N2-C2-N3	7.59	131.96	117.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	B	1187	TA1	3	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.