



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:18 PM GMT

PDB ID : 5CK7  
Title : Mouse ADP-dependent Glucokinase; AMP bound  
Authors : Richter, J.P.; Ronimus, R.S.; Sutherland-Smith, A.J.  
Deposited on : 2015-07-15  
Resolution : 2.99 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026982  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20026982


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.99 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	486	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	AMP	A	601	-	-	X	-

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 3497 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ADP-dependent glucokinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	448	Total	C	N	O	S	0	0	0
			3447	2188	601	646	12			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	38	MET	-	initiating methionine	UNP Q8VDL4
A	39	GLY	-	expression tag	UNP Q8VDL4
A	40	SER	-	expression tag	UNP Q8VDL4
A	41	GLY	-	expression tag	UNP Q8VDL4
A	42	SER	-	expression tag	UNP Q8VDL4
A	43	GLY	-	expression tag	UNP Q8VDL4
A	44	ASP	-	expression tag	UNP Q8VDL4
A	45	ASP	-	expression tag	UNP Q8VDL4
A	46	ASP	-	expression tag	UNP Q8VDL4
A	47	ASP	-	expression tag	UNP Q8VDL4
A	48	LYS	-	expression tag	UNP Q8VDL4
A	49	LEU	-	expression tag	UNP Q8VDL4
A	50	ALA	-	expression tag	UNP Q8VDL4
A	372	SER	ALA	conflict	UNP Q8VDL4
A	496	LYS	-	expression tag	UNP Q8VDL4
A	497	GLY	-	expression tag	UNP Q8VDL4
A	498	GLN	-	expression tag	UNP Q8VDL4
A	499	LEU	-	expression tag	UNP Q8VDL4
A	500	GLN	-	expression tag	UNP Q8VDL4
A	501	GLY	-	expression tag	UNP Q8VDL4
A	502	LYS	-	expression tag	UNP Q8VDL4
A	503	PRO	-	expression tag	UNP Q8VDL4
A	504	ILE	-	expression tag	UNP Q8VDL4
A	505	PRO	-	expression tag	UNP Q8VDL4
A	506	ASN	-	expression tag	UNP Q8VDL4
A	507	PRO	-	expression tag	UNP Q8VDL4
A	508	LEU	-	expression tag	UNP Q8VDL4

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	509	LEU	-	expression tag	UNP Q8VDL4
A	510	GLY	-	expression tag	UNP Q8VDL4
A	511	LEU	-	expression tag	UNP Q8VDL4
A	512	ASP	-	expression tag	UNP Q8VDL4
A	513	SER	-	expression tag	UNP Q8VDL4
A	514	THR	-	expression tag	UNP Q8VDL4
A	515	ARG	-	expression tag	UNP Q8VDL4
A	516	THR	-	expression tag	UNP Q8VDL4
A	517	GLY	-	expression tag	UNP Q8VDL4
A	518	HIS	-	expression tag	UNP Q8VDL4
A	519	HIS	-	expression tag	UNP Q8VDL4
A	520	HIS	-	expression tag	UNP Q8VDL4
A	521	HIS	-	expression tag	UNP Q8VDL4
A	522	HIS	-	expression tag	UNP Q8VDL4
A	523	HIS	-	expression tag	UNP Q8VDL4

- Molecule 2 is ADENOSINE MONOPHOSPHATE (three-letter code: AMP) (formula: C<sub>10</sub>H<sub>14</sub>N<sub>5</sub>O<sub>7</sub>P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			23	10	5	7	1		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	27	Total	O	0	0
			27	27		

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ADP-dependent glucokinase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	45.68 Å 58.49 Å 161.71 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.64 – 2.99 39.64 – 2.99	Depositor EDS
% Data completeness (in resolution range)	96.7 (39.64-2.99) 96.7 (39.64-2.99)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	0.09	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 3.01 Å)	Xtriage
Refinement program	REFMAC 5.8.0124	Depositor
R, $R_{free}$	0.227 , 0.243 0.228 , 0.254	Depositor DCC
$R_{free}$ test set	449 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	50.9	Xtriage
Anisotropy	0.527	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 40.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 8994 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	3497	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	64.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.57% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: AMP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.77	1/3529 (0.0%)	0.91	15/4799 (0.3%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	474	ARG	CD-NE	5.48	1.55	1.46

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	471	ASP	N-CA-CB	-12.07	88.87	110.60
1	A	277	ARG	NE-CZ-NH2	-11.99	114.31	120.30
1	A	456	ARG	NE-CZ-NH2	10.82	125.71	120.30
1	A	62	ASP	CB-CG-OD2	-7.28	111.75	118.30
1	A	331	LEU	CB-CG-CD2	-6.93	99.21	111.00
1	A	300	SER	CA-CB-OG	6.90	129.83	111.20
1	A	312	HIS	CB-CA-C	6.74	123.88	110.40
1	A	474	ARG	NE-CZ-NH2	6.11	123.36	120.30
1	A	471	ASP	CB-CG-OD2	-6.05	112.85	118.30
1	A	350	ASP	CB-CG-OD2	-5.95	112.95	118.30
1	A	453	GLU	CA-CB-CG	5.67	125.87	113.40
1	A	367	ARG	NE-CZ-NH1	5.56	123.08	120.30
1	A	373	SER	N-CA-C	-5.54	96.04	111.00
1	A	448	ASP	CB-CG-OD1	5.46	123.22	118.30
1	A	56	ARG	NE-CZ-NH1	5.28	122.94	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3447	0	3377	56	1
2	A	23	0	12	7	0
3	A	27	0	0	1	0
All	All	3497	0	3389	56	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (56) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:368:SER:OG	1:A:370:ASN:OD1	1.86	0.92
1:A:336:SER:OG	1:A:360:TRP:CZ2	2.28	0.85
1:A:228:ARG:O	1:A:228:ARG:NH1	2.13	0.81
1:A:336:SER:OG	1:A:360:TRP:CE2	2.36	0.79
1:A:305:GLU:OE1	3:A:701:HOH:O	2.04	0.75
1:A:478:LEU:HD23	2:A:601:AMP:O4'	1.87	0.75
1:A:337:GLY:HA3	1:A:340:SER:OG	1.88	0.74
1:A:338:PRO:HB3	1:A:437:LEU:HD11	1.70	0.74
1:A:459:ILE:HG22	1:A:461:PHE:CE1	2.27	0.69
1:A:493:ARG:HB2	1:A:494:PRO:CD	2.21	0.69
1:A:459:ILE:HG22	1:A:461:PHE:HE1	1.57	0.68
1:A:58:ALA:CB	1:A:419:ILE:HG21	2.26	0.65
1:A:336:SER:OG	1:A:360:TRP:NE1	2.29	0.65
1:A:338:PRO:HB3	1:A:437:LEU:CD1	2.26	0.64
1:A:442:ARG:HG3	1:A:442:ARG:O	1.97	0.64
1:A:298:LEU:HD22	1:A:321:LEU:HD11	1.79	0.64
1:A:228:ARG:HG2	1:A:228:ARG:HH11	1.64	0.62
1:A:299:ALA:O	1:A:301:MET:HG2	2.00	0.61
1:A:478:LEU:HD23	2:A:601:AMP:C1'	2.32	0.60
1:A:199:SER:HA	1:A:242:MET:HE2	1.82	0.60
1:A:228:ARG:HG2	1:A:228:ARG:NH1	2.15	0.60
1:A:478:LEU:HD23	2:A:601:AMP:N9	2.18	0.59
1:A:370:ASN:OD1	1:A:370:ASN:N	2.29	0.57
1:A:356:ASP:HB3	1:A:437:LEU:HD12	1.86	0.57

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:199:SER:HA	1:A:242:MET:CE	2.34	0.57
1:A:392:ASP:OD1	1:A:460:THR:OG1	2.23	0.56
1:A:371:ARG:HG3	1:A:371:ARG:NH1	2.20	0.55
1:A:438:GLU:OE1	1:A:456:ARG:NE	2.39	0.55
1:A:58:ALA:HB1	1:A:419:ILE:HG21	1.88	0.55
1:A:383:LEU:HA	2:A:601:AMP:H2'	1.88	0.54
1:A:461:PHE:CD1	1:A:461:PHE:N	2.76	0.53
1:A:359:PHE:CE1	1:A:456:ARG:HG2	2.43	0.53
1:A:336:SER:HG	1:A:360:TRP:HZ2	1.46	0.53
1:A:333:GLN:HA	1:A:340:SER:HB3	1.90	0.52
1:A:350:ASP:OD1	1:A:436:HIS:CE1	2.62	0.52
1:A:112:ARG:NH1	1:A:138:THR:HG23	2.24	0.52
1:A:459:ILE:CG2	1:A:461:PHE:CE1	2.93	0.52
1:A:455:HIS:N	1:A:455:HIS:CD2	2.78	0.51
1:A:342:LEU:HD21	1:A:344:SER:O	2.10	0.50
1:A:456:ARG:HG3	1:A:457:GLU:OE1	2.13	0.48
1:A:478:LEU:CD2	2:A:601:AMP:C4	2.97	0.47
1:A:478:LEU:HD23	2:A:601:AMP:C4	2.49	0.47
1:A:461:PHE:HD1	1:A:461:PHE:N	2.14	0.46
1:A:371:ARG:HG3	1:A:371:ARG:HH11	1.80	0.46
1:A:359:PHE:CZ	1:A:457:GLU:OE1	2.69	0.46
1:A:493:ARG:HB2	1:A:494:PRO:HD2	1.96	0.45
1:A:371:ARG:CG	1:A:371:ARG:HH11	2.28	0.45
1:A:299:ALA:O	1:A:300:SER:C	2.55	0.45
1:A:58:ALA:HB2	1:A:419:ILE:HG21	1.99	0.45
1:A:454:TRP:C	1:A:455:HIS:CD2	2.90	0.45
1:A:382:THR:O	2:A:601:AMP:O2'	2.36	0.44
1:A:335:ALA:HB3	1:A:360:TRP:HH2	1.82	0.44
1:A:456:ARG:CG	1:A:457:GLU:OE1	2.67	0.42
1:A:366:GLY:O	1:A:371:ARG:HD2	2.20	0.41
1:A:323:LEU:HD12	1:A:323:LEU:N	2.35	0.41
1:A:342:LEU:HD11	1:A:344:SER:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:364:GLU:OE2	1:A:417:GLU:OE2[1_455]	1.87	0.33

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	446/486 (92%)	432 (97%)	12 (3%)	2 (0%)	39 80

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	300	SER
1	A	337	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	368/401 (92%)	350 (95%)	18 (5%)	31 71

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	ASP
1	A	64	LEU
1	A	192	ASN
1	A	212	TYR
1	A	228	ARG
1	A	244	GLU
1	A	323	LEU
1	A	326	GLN
1	A	340	SER

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	341	SER
1	A	370	ASN
1	A	371	ARG
1	A	419	ILE
1	A	453	GLU
1	A	455	HIS
1	A	461	PHE
1	A	473	VAL
1	A	493	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	264	HIS
1	A	436	HIS
1	A	455	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	AMP	A	601	-	22,25,25	1.36	3 (13%)	22,38,38	2.19	8 (36%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	AMP	A	601	-	-	0/6/26/26	0/3/3/3

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	601	AMP	C2'-C1'	-3.57	1.48	1.53
2	A	601	AMP	C5-C4	2.42	1.46	1.40
2	A	601	AMP	O4'-C1'	2.55	1.44	1.41

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	AMP	N3-C2-N1	-5.80	124.32	128.87
2	A	601	AMP	O2'-C2'-C1'	-2.56	103.59	111.61
2	A	601	AMP	O5'-P-O1P	-2.24	101.45	107.08
2	A	601	AMP	O4'-C1'-N9	-2.19	103.96	108.11
2	A	601	AMP	N6-C6-N1	2.00	121.88	118.52
2	A	601	AMP	C2'-C3'-C4'	2.44	107.63	102.64
2	A	601	AMP	C1'-N9-C4	2.51	129.61	126.81
2	A	601	AMP	C4'-O4'-C1'	4.48	114.39	109.64

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	AMP	7	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	448/486 (92%)	0.09	15 (3%)	50 22	23, 56, 115, 152	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	338	PRO	4.0
1	A	336	SER	3.8
1	A	335	ALA	3.8
1	A	438	GLU	3.4
1	A	51	LEU	2.7
1	A	495	ASP	2.6
1	A	452	VAL	2.5
1	A	422	ASN	2.4
1	A	334	SER	2.3
1	A	436	HIS	2.3
1	A	431	GLU	2.2
1	A	437	LEU	2.2
1	A	339	HIS	2.1
1	A	445	LEU	2.1
1	A	429	PRO	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	AMP	A	601	23/23	0.89	0.19	-0.02	53,61,68,68	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.