



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 19, 2016 – 10:13 PM GMT

PDB ID : 5CNJ
Title : mGluR2 with glutamate analog
Authors : Monn, J.A.; Clawson, D.K.
Deposited on : 2015-07-17
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026982
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3152 (2.70-2.62)
Clashscore	102246	3524 (2.70-2.62)
Ramachandran outliers	100387	3469 (2.70-2.62)
Sidechain outliers	100360	3469 (2.70-2.62)
RSRZ outliers	91569	3161 (2.70-2.62)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	503	
1	B	503	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7184 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Metabotropic glutamate receptor 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	445	Total	C	N	O	S	0	0	0
			3496	2220	625	638	13			
1	B	445	Total	C	N	O	S	0	0	0
			3489	2217	621	638	13			

There are 24 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	MET	-	expression tag	UNP Q14416
A	0	ALA	-	expression tag	UNP Q14416
A	1	LEU	-	expression tag	UNP Q14416
A	234	SER	CYS	conflict	UNP Q14416
A	494	GLU	-	expression tag	UNP Q14416
A	495	GLY	-	expression tag	UNP Q14416
A	496	HIS	-	expression tag	UNP Q14416
A	497	HIS	-	expression tag	UNP Q14416
A	498	HIS	-	expression tag	UNP Q14416
A	499	HIS	-	expression tag	UNP Q14416
A	500	HIS	-	expression tag	UNP Q14416
A	501	HIS	-	expression tag	UNP Q14416
B	-1	MET	-	expression tag	UNP Q14416
B	0	ALA	-	expression tag	UNP Q14416
B	1	LEU	-	expression tag	UNP Q14416
B	234	SER	CYS	conflict	UNP Q14416
B	494	GLU	-	expression tag	UNP Q14416
B	495	GLY	-	expression tag	UNP Q14416
B	496	HIS	-	expression tag	UNP Q14416
B	497	HIS	-	expression tag	UNP Q14416
B	498	HIS	-	expression tag	UNP Q14416
B	499	HIS	-	expression tag	UNP Q14416
B	500	HIS	-	expression tag	UNP Q14416
B	501	HIS	-	expression tag	UNP Q14416

- Molecule 2 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is (1R,2S,4R,5R,6R)-2-amino-4-(1H-1,2,4-triazol-3-ylsulfanyl)bicyclo[3.1.0]hexane-2,6-dicarboxylic acid (three-letter code: 52Q) (formula: $C_{10}H_{12}N_4O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			19	10	4	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			19	10	4	4	1		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	81	Total	O	0	0
			81	81		
4	B	66	Total	O	0	0
			66	66		

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	80.39 Å 155.09 Å 92.71 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.04 – 2.65 92.71 – 2.65	Depositor EDS
% Data completeness (in resolution range)	97.9 (28.04-2.65) 97.8 (92.71-2.65)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.65 Å)	Xtriage
Refinement program	BUSTER 2.11.5	Depositor
R, R_{free}	0.193 , 0.241 0.202 , 0.244	Depositor DCC
R_{free} test set	1005 reflections (3.08%)	DCC
Wilson B-factor (Å ²)	51.6	Xtriage
Anisotropy	0.552	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 65.6	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	2 of 33680 reflections (0.006%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7184	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.26% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, 52Q

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/3580	0.71	0/4852
1	B	0.53	0/3574	0.71	0/4846
All	All	0.54	0/7154	0.71	0/9698

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3496	0	3406	19	0
1	B	3489	0	3387	30	0
2	A	14	0	13	0	0
3	A	19	0	9	0	0
3	B	19	0	9	0	0
4	A	81	0	0	0	0
4	B	66	0	0	2	0
All	All	7184	0	6824	49	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:99:LEU:HD22	1:B:150:GLN:CG	2.23	0.69
1:B:99:LEU:HD22	1:B:150:GLN:HG3	1.77	0.67
1:A:482:THR:HA	1:A:485:ILE:HD12	1.78	0.65
1:A:24:LYS:HD3	1:A:343:PRO:HB2	1.79	0.64
1:A:355:CYS:HB2	1:A:361:ASP:HB3	1.78	0.63
1:B:279:LEU:HD21	1:B:311:ALA:HB2	1.81	0.61
1:B:210:VAL:HG12	1:B:211:ALA:N	2.17	0.60
1:A:95:ASP:HB2	1:A:150:GLN:HG3	1.86	0.57
1:B:209:THR:OG1	1:B:238:SER:HA	2.06	0.56
1:B:209:THR:HG1	1:B:238:SER:HA	1.72	0.55
1:B:210:VAL:HG12	1:B:211:ALA:H	1.71	0.55
1:A:142:GLY:HA3	1:A:148:SER:OG	2.09	0.52
1:A:187:PRO:HD3	1:A:448:ASP:HB3	1.92	0.52
1:B:219:THR:HA	1:B:222:GLU:HG2	1.91	0.52
1:B:142:GLY:HA3	1:B:148:SER:OG	2.10	0.51
1:A:210:VAL:HG11	1:A:278:LEU:HD11	1.92	0.51
1:B:25:VAL:HG21	1:B:87:HIS:CE1	2.46	0.51
1:A:141:GLY:HA2	1:A:165:TYR:CE2	2.47	0.50
1:B:482:THR:HG22	1:B:485:ILE:HD12	1.94	0.49
1:A:221:ILE:HG12	1:A:269:PHE:HB2	1.94	0.49
1:B:163:ILE:HA	1:B:182:ALA:O	2.14	0.48
1:A:26:LEU:HD13	1:A:62:LEU:HD11	1.95	0.47
1:B:304:VAL:HG11	1:B:458:TYR:CE2	2.49	0.47
1:B:212:SER:HA	1:B:241:VAL:O	2.14	0.47
1:A:199:LEU:HD13	1:A:207:VAL:HG11	1.96	0.47
1:B:273:GLU:CD	1:B:273:GLU:H	2.18	0.47
1:B:206:TYR:C	1:B:207:VAL:HG13	2.35	0.47
1:B:265:VAL:HG22	1:B:290:THR:HB	1.97	0.47
1:B:177:ARG:NH1	4:B:701:HOH:O	2.38	0.46
1:A:207:VAL:HG12	1:A:265:VAL:HB	1.98	0.46
1:A:225:GLU:HG2	1:A:235:VAL:HG21	1.99	0.45
1:B:159:GLN:O	1:B:415:GLY:HA3	2.16	0.45
1:B:31:ASP:HB2	1:B:82:VAL:HG13	1.99	0.45
1:B:207:VAL:HG12	1:B:265:VAL:HB	2.00	0.44
1:A:193:LYS:HE3	1:A:231:ARG:HH22	1.84	0.43
1:A:324:ILE:HD12	1:A:379:MET:HA	2.01	0.43
1:A:36:GLY:HA3	1:A:88:ILE:HG12	2.00	0.43
1:B:304:VAL:HG11	1:B:458:TYR:CZ	2.55	0.42
1:A:273:GLU:CD	1:A:273:GLU:H	2.23	0.42
1:A:298:GLY:HA2	1:A:317:ILE:O	2.19	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:26:LEU:HD11	1:B:66:LEU:HD11	2.00	0.42
1:B:26:LEU:HB3	1:B:88:ILE:HB	2.01	0.41
1:B:89:LEU:HD13	1:B:101:GLN:HB3	2.02	0.41
1:A:195:MET:CE	1:A:294:SER:HB3	2.50	0.41
1:B:405:ARG:NH1	4:B:702:HOH:O	2.50	0.41
1:B:141:GLY:HA2	1:B:165:TYR:CE2	2.56	0.41
1:B:297:TRP:C	1:B:297:TRP:CD1	2.94	0.41
1:B:210:VAL:CG1	1:B:211:ALA:N	2.85	0.40
1:B:295:ASP:HA	1:B:317:ILE:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	441/503 (88%)	423 (96%)	14 (3%)	4 (1%)	21	44
1	B	441/503 (88%)	416 (94%)	25 (6%)	0	100	100
All	All	882/1006 (88%)	839 (95%)	39 (4%)	4 (0%)	34	59

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	364	ALA
1	A	325	SER
1	A	360	ARG
1	A	451	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	359/402 (89%)	349 (97%)	10 (3%)	51	79
1	B	358/402 (89%)	348 (97%)	10 (3%)	51	79
All	All	717/804 (89%)	697 (97%)	20 (3%)	51	79

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	24	LYS
1	A	25	VAL
1	A	93	SER
1	A	179	ASP
1	A	183	ARG
1	A	245	MET
1	A	268	LEU
1	A	269	PHE
1	A	361	ASP
1	A	368	ARG
1	B	60	GLN
1	B	143	SER
1	B	176	SER
1	B	208	SER
1	B	212	SER
1	B	215	ASP
1	B	269	PHE
1	B	297	TRP
1	B	335	ASP
1	B	358	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	439	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	NAG	A	601	1	14,14,15	0.33	0	15,19,21	0.78	1 (6%)
3	52Q	A	602	-	11,21,21	2.21	5 (45%)	4,33,33	1.85	1 (25%)
3	52Q	B	601	-	11,21,21	1.99	6 (54%)	4,33,33	2.02	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	601	1	-	0/6/23/26	0/1/1/1
3	52Q	A	602	-	-	0/2/38/38	0/2/3/3
3	52Q	B	601	-	-	0/2/38/38	0/2/3/3

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	52Q	C9-S19	-3.80	1.77	1.83
3	A	602	52Q	C7-C9	-3.52	1.46	1.55
3	B	601	52Q	C7-C9	-3.17	1.46	1.55
3	B	601	52Q	C9-S19	-3.07	1.79	1.83
3	A	602	52Q	C5-C10	-2.53	1.49	1.54
3	B	601	52Q	C5-C10	-2.48	1.49	1.54
3	A	602	52Q	C5-C9	-2.47	1.50	1.53
3	A	602	52Q	C6-C8	-2.31	1.47	1.50
3	B	601	52Q	C6-C8	-2.22	1.47	1.50
3	B	601	52Q	C6-C7	-2.21	1.48	1.50
3	B	601	52Q	C5-C9	-2.01	1.50	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	601	52Q	C3-C6-C7	-3.42	115.19	122.61
3	A	602	52Q	C3-C6-C7	-3.19	115.69	122.61
2	A	601	NAG	C1-O5-C5	2.60	115.96	112.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	445/503 (88%)	0.21	10 (2%) 65 64	33, 50, 79, 106	0
1	B	445/503 (88%)	0.32	19 (4%) 39 37	32, 58, 90, 110	0
All	All	890/1006 (88%)	0.26	29 (3%) 50 48	32, 54, 87, 110	0

All (29) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	435	ALA	5.9
1	B	233	ILE	5.0
1	B	436	ASP	3.9
1	A	461	ALA	3.8
1	B	437	THR	3.4
1	B	476	GLU	3.3
1	B	459	LEU	2.9
1	B	279	LEU	2.8
1	B	206	TYR	2.7
1	A	304	VAL	2.6
1	A	362	CYS	2.5
1	B	403	THR	2.5
1	B	111	SER	2.5
1	B	228	ALA	2.5
1	B	207	VAL	2.4
1	B	250	PHE	2.3
1	A	473	TYR	2.3
1	B	466	TYR	2.3
1	B	245	MET	2.3
1	A	49	ASP	2.2
1	A	437	THR	2.2
1	A	359	GLN	2.1
1	A	336	PRO	2.1
1	B	304	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	361	ASP	2.1
1	B	477	GLY	2.1
1	B	49	ASP	2.1
1	A	324	ILE	2.1
1	B	134	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	52Q	A	602	19/19	0.97	0.18	-0.35	35,46,59,60	0
3	52Q	B	601	19/19	0.98	0.15	-1.46	29,43,61,61	0
2	NAG	A	601	14/15	0.88	0.18	-	71,85,90,91	0

6.5 Other polymers [i](#)

There are no such residues in this entry.