



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 07:57 AM GMT

PDB ID : 3CS2  
Title : Crystal structure of PTE G60A mutant  
Authors : Kim, J.; Almo, S.C.  
Deposited on : 2008-04-08  
Resolution : 1.95 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.95 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1833 (1.96-1.96)
Clashscore	102246	1953 (1.96-1.96)
Ramachandran outliers	100387	1936 (1.96-1.96)
Sidechain outliers	100360	1936 (1.96-1.96)
RSRZ outliers	91569	1835 (1.96-1.96)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	331	<div> <div>2%</div> <div>82% 17%</div> </div>
1	B	331	<div> <div>86% 12%</div> </div>
1	K	331	<div> <div>3%</div> <div>87% 11%</div> </div>
1	P	331	<div> <div>85% 12%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11352 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Parathion hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	331	Total	C	N	O	S	0	3	0
			2548	1609	454	478	7			
1	B	330	Total	C	N	O	S	0	3	0
			2546	1608	453	478	7			
1	K	331	Total	C	N	O	S	0	2	0
			2542	1606	453	476	7			
1	P	330	Total	C	N	O	S	0	0	0
			2525	1597	450	471	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	60	ALA	GLY	ENGINEERED	UNP P0A434
B	60	ALA	GLY	ENGINEERED	UNP P0A434
K	60	ALA	GLY	ENGINEERED	UNP P0A434
P	60	ALA	GLY	ENGINEERED	UNP P0A434

- Molecule 2 is CACODYLATE ION (three-letter code: CAC) (formula: C<sub>2</sub>H<sub>6</sub>AsO<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	As	C	O	0	0
			5	1	2	2		
2	B	1	Total	As	C	O	0	0
			5	1	2	2		
2	K	1	Total	As	C	O	0	0
			5	1	2	2		
2	P	1	Total	As	C	O	0	0
			5	1	2	2		

- Molecule 3 is COBALT (II) ION (three-letter code: CO) (formula: Co).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	P	2	Total	Co	0	0
			2	2		
3	B	2	Total	Co	0	0
			2	2		
3	A	2	Total	Co	0	0
			2	2		
3	K	2	Total	Co	0	0
			2	2		

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	305	Total	O	0	0
			305	305		

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	276	Total 276	O 276	0	0
4	K	312	Total 312	O 312	0	0
4	P	270	Total 270	O 270	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: Parathion hydrolase




#### • Molecule 1: Parathion hydrolase



#### • Molecule 1: Parathion hydrolase



#### • Molecule 1: Parathion hydrolase



I333	I336	I337	I338	I339	Q343	E344	I353	P354	A355	R356	P360	T361	L362	R363	ALA

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	55.30 Å 68.30 Å 90.03 Å 90.05° 100.42° 89.96°	Depositor
Resolution (Å)	31.64 – 1.95 31.65 – 1.95	Depositor EDS
% Data completeness (in resolution range)	97.1 (31.64-1.95) 96.9 (31.65-1.95)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.92 (at 1.95 Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, $R_{free}$	0.167 , 0.224 0.167 , 0.225	Depositor DCC
$R_{free}$ test set	4597 reflections (5.29%)	DCC
Wilson B-factor (Å <sup>2</sup> )	19.9	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	(Not available) , (Not available)	EDS
Estimated twinning fraction	0.467 for -h,k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Outliers	0 of 91514 reflections	Xtriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	11352	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.96% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CAC, CO, KCX

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	0/2583	0.74	2/3509 (0.1%)
1	B	0.62	0/2581	0.69	0/3504
1	K	0.68	0/2577	0.77	4/3500 (0.1%)
1	P	0.63	0/2560	0.70	1/3476 (0.0%)
All	All	0.66	0/10301	0.73	7/13989 (0.1%)

There are no bond length outliers.

The worst 5 of 7 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	K	225	ARG	NE-CZ-NH2	-9.04	115.78	120.30
1	K	225	ARG	NE-CZ-NH1	7.16	123.88	120.30
1	K	89	ARG	NE-CZ-NH2	-6.36	117.12	120.30
1	A	131	TRP	CA-CB-CG	5.31	123.78	113.70
1	A	356	ARG	NE-CZ-NH2	-5.31	117.65	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2548	0	2564	46	0
1	B	2546	0	2558	30	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	K	2542	0	2559	42	0
1	P	2525	0	2545	34	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	K	5	0	0	0	0
2	P	5	0	0	0	0
3	A	2	0	0	0	0
3	B	2	0	0	0	0
3	K	2	0	0	0	0
3	P	2	0	0	0	0
4	A	305	0	0	8	0
4	B	276	0	0	4	0
4	K	312	0	0	13	0
4	P	270	0	0	5	0
All	All	11352	0	10226	152	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 152 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:259:ALA:HB2	1:K:272:LEU:HB2	1.26	1.11
1:A:259:ALA:HB2	1:A:272:LEU:HB2	1.15	1.10
1:B:337:ARG:NH2	1:B:343:GLN:HG2	1.67	1.08
1:B:337:ARG:HH21	1:B:343:GLN:HG2	0.90	1.05
1:K:225:ARG:HD3	4:K:2451:HOH:O	1.58	1.02

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	331/331 (100%)	317 (96%)	14 (4%)	0	100	100
1	B	330/331 (100%)	316 (96%)	14 (4%)	0	100	100
1	K	330/331 (100%)	319 (97%)	10 (3%)	1 (0%)	46	35
1	P	327/331 (99%)	317 (97%)	10 (3%)	0	100	100
All	All	1318/1324 (100%)	1269 (96%)	48 (4%)	1 (0%)	56	48

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	259	ALA

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	267/264 (101%)	260 (97%)	7 (3%)	54	43
1	B	267/264 (101%)	255 (96%)	12 (4%)	34	18
1	K	266/264 (101%)	258 (97%)	8 (3%)	48	36
1	P	264/264 (100%)	250 (95%)	14 (5%)	28	13
All	All	1064/1056 (101%)	1023 (96%)	41 (4%)	40	24

5 of 41 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	363	ARG
1	K	303	LEU
1	P	343	GLN
1	K	196	VAL
1	K	274	ILE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 10 such sidechains are listed below:

Mol	Chain	Res	Type
1	K	212	GLN
1	K	295	GLN
1	K	343	GLN
1	B	312	ASN
1	K	312	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
1	KCX	A	169	1,3	7,11,12	0.79	0	7,12,14	1.06	1 (14%)
1	KCX	B	169	1,3	7,11,12	0.57	0	7,12,14	1.50	1 (14%)
1	KCX	K	169	1,3	7,11,12	0.76	0	7,12,14	0.98	0
1	KCX	P	169	1,3	7,11,12	0.33	0	7,12,14	0.91	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	KCX	A	169	1,3	-	0/6/10/12	0/0/0/0
1	KCX	B	169	1,3	-	0/6/10/12	0/0/0/0
1	KCX	K	169	1,3	-	0/6/10/12	0/0/0/0
1	KCX	P	169	1,3	-	0/6/10/12	0/0/0/0

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	169	KCX	CE-NZ-CX	-3.27	119.79	123.49
1	A	169	KCX	O-C-CA	-2.06	120.13	125.49
1	P	169	KCX	O-C-CA	-2.05	120.15	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 12 ligands modelled in this entry, 8 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	CAC	A	1	3	0,4,4	0.00	-	0,6,6	0.00	-
2	CAC	B	2	3	0,4,4	0.00	-	0,6,6	0.00	-
2	CAC	K	3	3	0,4,4	0.00	-	0,6,6	0.00	-
2	CAC	P	4	3	0,4,4	0.00	-	0,6,6	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CAC	A	1	3	-	0/0/0/0	0/0/0/0
2	CAC	B	2	3	-	0/0/0/0	0/0/0/0
2	CAC	K	3	3	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	CAC	P	4	3	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	330/331 (99%)	-0.29	7 (2%) 67 75	11, 17, 37, 62	0
1	B	329/331 (99%)	-0.26	1 (0%) 94 96	11, 21, 36, 44	2 (0%)
1	K	330/331 (99%)	-0.25	10 (3%) 54 64	11, 17, 37, 64	0
1	P	329/331 (99%)	-0.30	1 (0%) 94 96	12, 21, 37, 45	3 (0%)
All	All	1318/1324 (99%)	-0.28	19 (1%) 78 85	11, 19, 37, 64	5 (0%)

The worst 5 of 19 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	K	260	ILE	6.7
1	K	262	LEU	6.1
1	A	263	GLU	5.2
1	A	259	ALA	5.0
1	A	262	LEU	4.5

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
1	KCX	P	169	12/13	0.98	0.07	-	13,15,16,16	0
1	KCX	B	169	12/13	0.94	0.10	-	13,16,17,17	0
1	KCX	A	169	12/13	0.95	0.08	-	11,13,16,17	0
1	KCX	K	169	12/13	0.97	0.07	-	12,14,14,15	0

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	CAC	K	3	5/5	0.99	0.10	1.82	16,17,22,23	5
2	CAC	P	4	5/5	0.99	0.10	1.00	20,22,24,24	5
2	CAC	A	1	5/5	0.99	0.09	-0.04	17,18,21,23	5
2	CAC	B	2	5/5	0.99	0.08	-0.51	16,17,21,21	5
3	CO	B	3	1/1	1.00	0.05	-	14,14,14,14	0
3	CO	K	6	1/1	1.00	0.07	-	14,14,14,14	0
3	CO	A	365	1/1	1.00	0.05	-	12,12,12,12	0
3	CO	B	4	1/1	0.99	0.05	-	17,17,17,17	0
3	CO	P	7	1/1	1.00	0.07	-	15,15,15,15	0
3	CO	A	2	1/1	1.00	0.06	-	13,13,13,13	0
3	CO	P	8	1/1	0.99	0.04	-	17,17,17,17	0
3	CO	K	5	1/1	1.00	0.06	-	11,11,11,11	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.