



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:38 PM GMT

PDB ID : 4CWY
Title : Structure of bovine endothelial nitric oxide synthase heme domain in complex with 6-(5-(((3R,4R)-4-((6-AMINO-4-METHYLPYRIDIN-2-YL)METHYL) P YRROLIDIN-3-YL)OXY)PENTYL)-4-METHYLPYRIDIN-2-AMINE
Authors : Li, H.; Poulos, T.L.
Deposited on : 2014-04-03
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	443	<div> <div>12%</div> <div>83%</div> <div>7% • 9%</div> </div>
1	B	443	<div> <div>9%</div> <div>80%</div> <div>9% • 9%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	ACT	B	860	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6896 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called NITRIC OXIDE SYNTHASE, ENDOTHELIAL.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	404	Total	As	C	N	O	S	0	0	0
			3212	1	2043	564	588	16			
1	B	402	Total	As	C	N	O	S	0	0	0
			3201	1	2036	562	586	16			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	100	ARG	CYS	SEE REMARK 999	UNP P29473
B	100	ARG	CYS	SEE REMARK 999	UNP P29473

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		

- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (three-letter code: H4B) (formula: $C_9H_{15}N_5O_3$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	9	5	3		
3	B	1	Total	C	N	O	0	0
			17	9	5	3		

- Molecule 4 is 6-[5-({(3R,4R)-4-[(6-AMINO-4-METHYLPYRIDIN-2-YL)METHYL]PYRROLIDIN-3-YL}OXY)PENTYL]-4-METHYLPYRIDIN-2-AMINE (three-letter code: HW1) (formula: $C_{22}H_{33}N_5O$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 28	C 22	N 5	O 1	0	0
4	B	1	Total 28	C 22	N 5	O 1	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: $\text{C}_2\text{H}_3\text{O}_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 4 2 2	0	0
5	A	1	Total C O 4 2 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	B	1	Total	Zn	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	159	Total	O	0	0
			159	159		
7	B	131	Total	O	0	0
			131	131		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



• Molecule 1: NITRIC OXIDE SYNTHASE, ENDOTHELIAL



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.88Å 106.47Å 156.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.08 – 2.15 38.01 – 2.15	Depositor EDS
% Data completeness (in resolution range)	99.4 (39.08-2.15) 99.4 (38.01-2.15)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.68 (at 2.14Å)	Xtriage
Refinement program	REFMAC 5.6.0117	Depositor
R, R_{free}	0.167 , 0.204 0.166 , 0.204	Depositor DCC
R_{free} test set	2683 reflections (5.28%)	DCC
Wilson B-factor (Å ²)	42.0	Xtriage
Anisotropy	0.478	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	1 of 53603 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6896	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, H4B, CAS, ACT, HEM, HW1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	4/3292 (0.1%)	0.65	0/4483
1	B	0.64	6/3280 (0.2%)	0.63	0/4466
All	All	0.65	10/6572 (0.2%)	0.64	0/8949

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	447	TRP	CD2-CE2	5.95	1.48	1.41
1	A	396	TRP	CD2-CE2	5.77	1.48	1.41
1	A	281	TRP	CD2-CE2	5.55	1.48	1.41
1	A	358	TRP	CD2-CE2	5.53	1.48	1.41
1	B	281	TRP	CD2-CE2	5.41	1.47	1.41
1	B	358	TRP	CD2-CE2	5.35	1.47	1.41
1	B	332	TRP	CD2-CE2	5.21	1.47	1.41
1	B	396	TRP	CD2-CE2	5.11	1.47	1.41
1	B	447	TRP	CD2-CE2	5.06	1.47	1.41
1	B	76	TRP	CD2-CE2	5.02	1.47	1.41

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3212	0	3114	20	0
1	B	3201	0	3104	22	0
2	A	43	0	30	2	0
2	B	43	0	30	6	0
3	A	17	0	15	2	0
3	B	17	0	15	1	0
4	A	28	0	33	3	0
4	B	28	0	33	3	0
5	A	8	0	6	0	0
5	B	8	0	6	0	0
6	B	1	0	0	0	0
7	A	159	0	0	1	0
7	B	131	0	0	1	0
All	All	6896	0	6386	47	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (47) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:384:CAS:AS	1:A:384:CAS:SG	2.58	1.22
1:B:384:CAS:AS	1:B:384:CAS:SG	2.64	1.16
1:A:384:CAS:CB	1:A:384:CAS:AS	2.99	0.71
1:A:240:ARG:HD3	1:A:298:PRO:HB3	1.74	0.69
1:A:384:CAS:HB2	1:A:384:CAS:AS	2.59	0.62
1:B:71:PRO:HG2	1:B:84:ASP:HB3	1.82	0.62
1:B:476:ARG:NH1	7:B:2126:HOH:O	2.29	0.61
2:B:500:HEM:C1C	4:B:800:HW1:H34	2.37	0.60
1:B:259:GLN:NE2	1:B:260:ASP:OD2	2.33	0.60
1:B:233:PHE:HB3	1:B:234:PRO:CD	2.35	0.55
1:B:99:ARG:HG2	1:B:100:ARG:HD2	1.88	0.54
1:A:246:TRP:HB2	1:A:294:LEU:HB3	1.89	0.53
2:A:500:HEM:HBA2	4:A:800:HW1:H24	1.91	0.52
2:B:500:HEM:HBC2	2:B:500:HEM:HMC1	1.92	0.52
2:B:500:HEM:HBC2	2:B:500:HEM:CMC	2.40	0.52
1:A:449:TRP:HA	3:A:600:H4B:N1	2.26	0.51
1:A:240:ARG:HD3	1:A:298:PRO:CB	2.41	0.51
1:A:378:LEU:HB2	7:A:2118:HOH:O	2.10	0.51
1:B:449:TRP:HA	3:B:600:H4B:N1	2.26	0.51
1:A:370:CYS:SG	1:A:378:LEU:HD13	2.51	0.51
1:A:126:LEU:HD11	1:A:156:GLU:HA	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:72:ARG:HB2	1:A:83:TYR:CE1	2.50	0.46
1:A:240:ARG:HD2	1:A:241:GLY:O	2.16	0.46
1:A:102:LEU:HD11	1:B:71:PRO:HB3	1.98	0.46
1:B:178:GLN:HE22	1:B:181:ARG:HH11	1.63	0.46
1:B:358:TRP:H	2:B:500:HEM:HAB	1.82	0.45
1:B:231:THR:O	1:B:353:ALA:HA	2.16	0.45
1:B:236:ARG:HD2	1:B:242:ASP:OD1	2.17	0.45
1:A:472:SER:HA	1:A:473:PRO:C	2.38	0.43
1:A:70:PHE:HB3	1:A:84:ASP:O	2.18	0.43
1:B:337:ALA:HB2	1:B:356:SER:HB3	2.00	0.43
1:A:185:ARG:NH1	1:A:477:TYR:OH	2.51	0.43
1:B:311:LEU:HD23	1:B:311:LEU:HA	1.91	0.42
1:B:281:TRP:HB2	1:B:304:LEU:HD21	2.00	0.42
1:B:236:ARG:CD	1:B:242:ASP:OD1	2.67	0.42
1:A:338:VAL:HG23	4:A:800:HW1:H31	2.02	0.42
1:B:366:THR:O	1:B:370:CYS:HB2	2.19	0.42
1:B:481:PRO:HD2	1:B:482:TRP:CZ3	2.55	0.42
1:B:384:CAS:AS	1:B:384:CAS:CB	3.28	0.42
1:A:105:LEU:HD22	1:B:465:GLU:HB3	2.02	0.42
2:A:500:HEM:O1A	3:A:600:H4B:N3	2.49	0.42
1:A:68:PRO:HB2	1:A:70:PHE:HD1	1.85	0.41
2:B:500:HEM:HBA2	4:B:800:HW1:H24	2.03	0.41
1:B:240:ARG:HD2	1:B:241:GLY:O	2.21	0.41
1:A:392:THR:HB	1:B:423:HIS:HB2	2.03	0.40
2:B:500:HEM:NC	4:B:800:HW1:H34	2.35	0.40
4:A:800:HW1:H19	4:A:800:HW1:H16	1.80	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	399/443 (90%)	391 (98%)	8 (2%)	0	100	100
1	B	397/443 (90%)	387 (98%)	10 (2%)	0	100	100
All	All	796/886 (90%)	778 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	342/375 (91%)	331 (97%)	11 (3%)	46	45
1	B	341/375 (91%)	328 (96%)	13 (4%)	40	37
All	All	683/750 (91%)	659 (96%)	24 (4%)	43	41

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	100	ARG
1	A	102	LEU
1	A	105	LEU
1	A	123	GLU
1	A	125	LEU
1	A	293	LEU
1	A	294	LEU
1	A	328	LEU
1	A	331	ARG
1	A	378	LEU
1	A	416	LYS
1	B	92	GLN
1	B	100	ARG
1	B	102	LEU
1	B	105	LEU
1	B	149	GLU
1	B	207	GLN
1	B	223	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	225	ASN
1	B	240	ARG
1	B	278	GLN
1	B	330	LEU
1	B	378	LEU
1	B	389	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	146	GLN
1	A	191	GLN
1	A	376	ASN
1	A	413	GLN
1	A	468	ASN
1	B	178	GLN
1	B	191	GLN
1	B	222	ASN
1	B	225	ASN
1	B	278	GLN
1	B	376	ASN
1	B	405	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	CAS	A	384	1	5,8,9	1.11	0	2,9,11	1.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CAS	B	384	1	5,8,9	1.21	0	2,9,11	1.58	1 (50%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	A	384	1	-	0/0/7/9	0/0/0/0
1	CAS	B	384	1	-	0/0/7/9	0/0/0/0

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	384	CAS	O-C-CA	-2.23	119.69	125.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	384	CAS	3	0
1	B	384	CAS	2	0

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 1 is monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	HEM	A	500	1	30,50,50	2.57	10 (33%)	24,82,82	2.81	11 (45%)
3	H4B	A	600	-	13,18,18	0.66	0	11,26,26	2.66	6 (54%)
4	HW1	A	800	-	27,30,30	0.69	0	31,40,40	2.00	5 (16%)
5	ACT	A	860	-	1,3,3	1.31	0	0,3,3	0.00	-
5	ACT	A	861	-	1,3,3	2.55	1 (100%)	0,3,3	0.00	-
2	HEM	B	500	1	30,50,50	2.72	13 (43%)	24,82,82	2.89	9 (37%)
3	H4B	B	600	-	13,18,18	0.63	0	11,26,26	2.32	4 (36%)
4	HW1	B	800	-	27,30,30	0.67	0	31,40,40	2.01	7 (22%)
5	ACT	B	860	-	1,3,3	1.29	0	0,3,3	0.00	-
5	ACT	B	861	-	1,3,3	1.91	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	HEM	A	500	1	-	0/10/54/54	0/0/8/8
3	H4B	A	600	-	-	0/8/17/17	0/2/2/2
4	HW1	A	800	-	-	0/13/23/23	0/3/3/3
5	ACT	A	860	-	-	0/0/0/0	0/0/0/0
5	ACT	A	861	-	-	0/0/0/0	0/0/0/0
2	HEM	B	500	1	-	0/10/54/54	0/0/8/8
3	H4B	B	600	-	-	0/8/17/17	0/2/2/2
4	HW1	B	800	-	-	0/13/23/23	0/3/3/3
5	ACT	B	860	-	-	0/0/0/0	0/0/0/0
5	ACT	B	861	-	-	0/0/0/0	0/0/0/0

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C3B-C4B	-5.55	1.46	1.51
2	A	500	HEM	C3B-C4B	-3.18	1.49	1.51
2	A	500	HEM	C2D-C3D	-2.85	1.46	1.54
2	B	500	HEM	C2D-C3D	-2.77	1.46	1.54
2	B	500	HEM	C3D-C4D	-2.24	1.48	1.51
2	A	500	HEM	C2C-C1C	-2.22	1.48	1.52
2	B	500	HEM	CHD-C1D	2.19	1.45	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	500	HEM	C4A-CHB	2.23	1.46	1.39
2	B	500	HEM	FE-NB	2.24	2.09	1.97
2	A	500	HEM	FE-NB	2.32	2.09	1.97
2	A	500	HEM	CHC-C4B	2.38	1.45	1.38
2	B	500	HEM	FE-ND	2.48	2.10	1.97
5	A	861	ACT	CH3-C	2.55	1.52	1.48
2	B	500	HEM	CHC-C4B	3.00	1.47	1.38
2	B	500	HEM	C2A-C3A	3.12	1.46	1.37
2	A	500	HEM	C2A-C3A	3.41	1.47	1.37
2	B	500	HEM	CHC-C1C	3.82	1.45	1.36
2	A	500	HEM	CHC-C1C	3.85	1.45	1.36
2	B	500	HEM	CHD-C4C	3.86	1.45	1.36
2	A	500	HEM	CHD-C4C	4.05	1.45	1.36
2	B	500	HEM	C1C-NC	6.52	1.44	1.36
2	A	500	HEM	C1C-NC	6.73	1.44	1.36
2	A	500	HEM	C4C-NC	6.86	1.44	1.36
2	B	500	HEM	C4C-NC	7.01	1.44	1.36

All (42) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	500	HEM	CBA-CAA-C2A	-6.68	100.55	112.53
2	A	500	HEM	CBA-CAA-C2A	-4.70	104.10	112.53
2	B	500	HEM	C3B-CAB-CBB	-3.69	118.79	124.46
3	A	600	H4B	N3-C2-N1	-3.61	119.62	125.53
4	A	800	HW1	C25-C26-N21	-3.49	119.03	122.96
3	B	600	H4B	N3-C2-N1	-3.36	120.02	125.53
2	A	500	HEM	C3B-CAB-CBB	-3.04	119.79	124.46
2	B	500	HEM	C3C-CAC-CBC	-3.04	119.79	124.46
4	A	800	HW1	C05-C06-N01	-3.01	119.56	122.96
4	B	800	HW1	C05-C06-N01	-2.96	119.61	122.96
2	A	500	HEM	C3B-C4B-CHC	-2.61	119.48	123.16
4	A	800	HW1	C04-C05-C06	-2.59	118.66	120.28
4	B	800	HW1	C25-C26-N21	-2.54	120.09	122.96
2	A	500	HEM	C3C-CAC-CBC	-2.38	120.80	124.46
2	B	500	HEM	CBD-CAD-C3D	-2.16	107.26	113.55
4	B	800	HW1	C04-C05-C06	-2.07	118.98	120.28
4	B	800	HW1	C24-C25-C26	-2.04	119.00	120.28
2	A	500	HEM	CBD-CAD-C3D	-2.03	107.66	113.55
3	A	600	H4B	C4A-C8A-N8	2.02	120.81	118.43
2	A	500	HEM	C3B-C4B-NB	2.21	115.86	111.63
3	A	600	H4B	N2-C2-N1	2.36	121.11	117.20

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	600	H4B	C2-N1-C8A	2.63	120.45	114.54
3	A	600	H4B	C2-N1-C8A	2.63	120.46	114.54
4	B	800	HW1	C14-C26-N21	2.82	119.85	115.69
2	B	500	HEM	CMD-C2D-C3D	3.42	129.50	114.35
2	A	500	HEM	CMD-C2D-C3D	3.48	129.76	114.35
3	B	600	H4B	C4-N3-C2	3.71	121.09	115.94
3	B	600	H4B	C4-C4A-C8A	4.21	118.37	114.56
3	A	600	H4B	C4-N3-C2	4.72	122.49	115.94
2	B	500	HEM	CAD-C3D-C4D	4.72	129.12	112.47
4	B	800	HW1	C22-N21-C26	4.81	121.65	118.23
2	B	500	HEM	CMC-C2C-C3C	4.81	128.54	116.53
2	A	500	HEM	CAD-C3D-C4D	4.85	129.57	112.47
3	A	600	H4B	C4-C4A-C8A	4.91	119.00	114.56
2	B	500	HEM	CAD-C3D-C2D	4.98	127.53	113.22
4	A	800	HW1	C22-N21-C26	5.00	121.78	118.23
2	A	500	HEM	CMC-C2C-C3C	5.06	129.15	116.53
2	A	500	HEM	CAD-C3D-C2D	5.06	127.77	113.22
2	B	500	HEM	CMB-C2B-C3B	5.48	130.21	116.53
2	A	500	HEM	CMB-C2B-C3B	5.78	130.96	116.53
4	A	800	HW1	C02-N01-C06	6.63	122.94	118.23
4	B	800	HW1	C02-N01-C06	6.75	123.03	118.23

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	500	HEM	2	0
3	A	600	H4B	2	0
4	A	800	HW1	3	0
2	B	500	HEM	6	0
3	B	600	H4B	1	0
4	B	800	HW1	3	0

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	403/443 (90%)	0.67	54 (13%) 4 7	34, 46, 77, 99	0
1	B	401/443 (90%)	0.53	41 (10%) 9 15	34, 49, 81, 110	0
All	All	804/886 (90%)	0.60	95 (11%) 6 10	34, 47, 80, 110	0

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	108	PRO	7.0
1	A	121	PRO	6.8
1	A	69	LYS	6.8
1	A	98	PRO	5.3
1	B	99	ARG	5.2
1	A	123	GLU	5.2
1	A	67	GLY	5.2
1	A	259	GLN	4.8
1	A	92	GLN	4.8
1	A	99	ARG	4.7
1	B	140	ILE	4.7
1	A	160	ALA	4.7
1	B	259	GLN	4.6
1	B	121	PRO	4.6
1	B	97	THR	4.5
1	A	93	ASP	4.4
1	A	91	GLN	4.3
1	B	146	GLN	4.3
1	A	239	GLY	4.2
1	A	238	PRO	4.2
1	A	94	GLY	4.1
1	A	95	PRO	4.1
1	A	122	ALA	4.0
1	A	261	GLY	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	93	ASP	3.7
1	A	454	ILE	3.7
1	B	98	PRO	3.7
1	A	108	PRO	3.6
1	A	124	GLN	3.6
1	A	126	LEU	3.6
1	B	261	GLY	3.6
1	A	451	VAL	3.5
1	B	143	SER	3.4
1	B	122	ALA	3.4
1	A	450	ILE	3.4
1	B	144	GLY	3.4
1	A	338	VAL	3.3
1	B	223	ARG	3.3
1	B	70	PHE	3.3
1	B	69	LYS	3.2
1	B	92	GLN	3.2
1	A	102	LEU	3.2
1	B	390	ARG	3.1
1	A	68	PRO	3.1
1	B	239	GLY	3.1
1	A	260	ASP	3.1
1	B	141	LYS	3.1
1	B	96	CYS	3.0
1	A	449	TRP	3.0
1	B	142	ARG	3.0
1	A	96	CYS	3.0
1	B	462	PHE	3.0
1	A	97	THR	3.0
1	B	147	ALA	2.9
1	A	156	GLU	2.9
1	B	100	ARG	2.9
1	A	337	ALA	2.9
1	B	260	ASP	2.9
1	A	336	PRO	2.8
1	A	362	THR	2.8
1	B	91	GLN	2.8
1	B	94	GLY	2.7
1	A	364	ILE	2.7
1	A	257	ARG	2.7
1	A	355	PHE	2.7
1	B	123	GLU	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	83	TYR	2.6
1	A	230	ILE	2.5
1	B	454	ILE	2.5
1	B	145	SER	2.5
1	A	414	LEU	2.5
1	A	130	ARG	2.5
1	B	150	GLU	2.4
1	A	106	VAL	2.4
1	B	107	LEU	2.4
1	B	258	GLN	2.3
1	B	102	LEU	2.3
1	A	452	PRO	2.2
1	A	365	GLY	2.2
1	A	231	THR	2.2
1	A	153	GLN	2.2
1	A	366	THR	2.2
1	B	353	ALA	2.1
1	B	456	GLY	2.1
1	A	187	VAL	2.1
1	B	95	PRO	2.1
1	A	100	ARG	2.1
1	A	223	ARG	2.1
1	A	101	CYS	2.1
1	A	390	ARG	2.0
1	A	456	GLY	2.0
1	A	159	VAL	2.0
1	B	232	VAL	2.0
1	A	232	VAL	2.0
1	B	451	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
1	CAS	B	384	9/10	0.96	0.10	-	54,55,70,76	0
1	CAS	A	384	9/10	0.97	0.09	-	45,46,61,64	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
5	ACT	B	860	4/4	0.99	0.28	20.17	56,60,62,64	0
2	HEM	A	500	43/43	0.98	0.23	1.11	30,32,42,51	0
5	ACT	A	861	4/4	0.97	0.20	1.10	41,42,43,45	0
4	HW1	B	800	28/28	0.92	0.19	0.75	36,47,60,63	0
2	HEM	B	500	43/43	0.99	0.15	0.69	33,37,44,50	0
4	HW1	A	800	28/28	0.92	0.23	0.54	34,47,55,56	0
3	H4B	A	600	17/17	0.98	0.19	0.13	39,42,46,50	0
3	H4B	B	600	17/17	0.97	0.15	-0.40	38,42,45,45	0
5	ACT	A	860	4/4	0.96	0.12	-0.63	52,59,59,61	0
5	ACT	B	861	4/4	0.96	0.10	-1.33	42,45,46,46	0
6	ZN	B	1483	1/1	0.99	0.02	-4.78	49,49,49,49	1

6.5 Other polymers [i](#)

There are no such residues in this entry.