



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 23, 2016 – 04:41 PM EDT

PDB ID : 5CXV  
Title : Structure of the human M1 muscarinic acetylcholine receptor bound to antagonist Tiotropium  
Authors : Sun, B.; Feng, D.; Li, X.; Kobilka, T.S.; Kobilka, B.K.  
Deposited on : 2015-07-29  
Resolution : 2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027107  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0122  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027107


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2103 (2.70-2.70)
Clashscore	102246	2422 (2.70-2.70)
Ramachandran outliers	100387	2382 (2.70-2.70)
Sidechain outliers	100360	2382 (2.70-2.70)
RSRZ outliers	91569	2107 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	515	<div> <div>5%</div> <div> <div></div> <div>69%</div> <div>17%</div> <div>14%</div> </div> </div>
2	C	7	<div> <div>43%</div> <div>43%</div> <div>14%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	EDO	A	505	-	-	-	X
5	EDO	A	507	-	-	-	X
6	PGE	A	504	-	-	-	X

## 2 Entry composition

There are 8 unique types of molecules in this entry. The entry contains 3599 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Muscarinic acetylcholine receptor M1,Endolysin,Muscarinic acetylcholine receptor M1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	444	Total	C	N	O	S	0	0	0
			3439	2248	572	597	22			

There are 39 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-24	MET	-	initiating methionine	UNP P11229
A	-23	LYS	-	expression tag	UNP P11229
A	-22	THR	-	expression tag	UNP P11229
A	-21	ILE	-	expression tag	UNP P11229
A	-20	ILE	-	expression tag	UNP P11229
A	-19	ALA	-	expression tag	UNP P11229
A	-18	LEU	-	expression tag	UNP P11229
A	-17	SER	-	expression tag	UNP P11229
A	-16	TYR	-	expression tag	UNP P11229
A	-15	ILE	-	expression tag	UNP P11229
A	-14	PHE	-	expression tag	UNP P11229
A	-13	CYS	-	expression tag	UNP P11229
A	-12	LEU	-	expression tag	UNP P11229
A	-11	VAL	-	expression tag	UNP P11229
A	-10	PHE	-	expression tag	UNP P11229
A	-9	ALA	-	expression tag	UNP P11229
A	-8	ASP	-	expression tag	UNP P11229
A	-7	TYR	-	expression tag	UNP P11229
A	-6	LYS	-	expression tag	UNP P11229
A	-5	ASP	-	expression tag	UNP P11229
A	-4	ASP	-	expression tag	UNP P11229
A	-3	ASP	-	expression tag	UNP P11229
A	-2	ASP	-	expression tag	UNP P11229
A	-1	ALA	-	expression tag	UNP P11229
A	0	ALA	-	expression tag	UNP P11229
A	1	ALA	-	expression tag	UNP P11229

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	2	GLN	ASN	engineered mutation	UNP P11229
A	12	GLN	ASN	engineered mutation	UNP P11229
A	110	GLN	ASN	engineered mutation	UNP P11229
A	1011	GLY	ARG	variant	UNP P00720
A	1053	THR	CYS	engineered mutation	UNP P00720
A	1096	ALA	CYS	engineered mutation	UNP P00720
A	1136	ARG	ILE	variant	UNP P00720
A	461	HIS	-	expression tag	UNP P11229
A	462	HIS	-	expression tag	UNP P11229
A	463	HIS	-	expression tag	UNP P11229
A	464	HIS	-	expression tag	UNP P11229
A	465	HIS	-	expression tag	UNP P11229
A	466	HIS	-	expression tag	UNP P11229


- Molecule 2 is a protein called FLAG peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	C	7	Total	C	N	O	0	0	0
			61	35	8	18			

- Molecule 3 is (1R,2R,4S,5S,7S)-7-{[hydroxy(dithiophen-2-yl)acetyl]oxy}-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.1.0 2,4 ]nonane (three-letter code: 0HK) (formula: C<sub>19</sub>H<sub>22</sub>NO<sub>4</sub>S<sub>2</sub>).




- Molecule 4 is CHOLESTEROL HEMISUCCINATE (three-letter code: Y01) (formula: C<sub>31</sub>H<sub>50</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			35	31	4		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

*Continued from previous page...*


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 7 is GLYCEROL (three-letter code: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	6	Total	O	0	0
			6	6		
8	C	4	Total	O	0	0
			4	4		




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Muscarinic acetylcholine receptor M1,Endolysin,Muscarinic acetylcholine receptor M1



- Molecule 2: FLAG peptide



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	58.05Å 72.19Å 175.69Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.00 – 2.70 28.94 – 2.70	Depositor EDS
% Data completeness (in resolution range)	96.0 (30.00-2.70) 96.2 (28.94-2.70)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 2.68Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, $R_{free}$	0.231 , 0.282 0.233 , 0.283	Depositor DCC
$R_{free}$ test set	1011 reflections (5.26%)	DCC
Wilson B-factor (Å <sup>2</sup> )	70.0	Xtriage
Anisotropy	0.285	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.33 , 61.3	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 20363 reflections (0.005%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3599	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	73.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.50% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 0HK, GOL, PGE, Y01, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.30	0/3520	0.47	1/4807 (0.0%)
2	C	0.21	0/61	0.43	0/81
All	All	0.30	0/3581	0.47	1/4888 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	376	LEU	CA-CB-CG	5.15	127.14	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3439	0	3402	53	0
2	C	61	0	44	3	0
3	A	26	0	22	1	0
4	A	35	0	49	0	0
5	A	12	0	18	0	0
6	A	10	0	14	0	0
7	A	6	0	8	0	0
8	A	6	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	C	4	0	0	0	0
All	All	3599	0	3557	55	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (55) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:173:VAL:HG23	1:A:179:TYR:CE2	2.09	0.87
1:A:1086:VAL:HG12	1:A:1121:GLN:HB2	1.66	0.76
2:C:2:TYR:O	2:C:4:ASP:N	2.21	0.73
1:A:1033:THR:HG22	1:A:1035:SER:H	1.53	0.73
1:A:218:ARG:NH2	1:A:360:GLU:OE2	2.25	0.70
1:A:173:VAL:HG23	1:A:179:TYR:CZ	2.31	0.65
1:A:91:TRP:N	1:A:176:GLY:O	2.19	0.64
1:A:423:LYS:HG2	1:A:424:ALA:H	1.65	0.62
1:A:418:TYR:O	1:A:422:ASN:HB3	2.00	0.61
1:A:25:VAL:HA	1:A:87:LEU:HD22	1.83	0.61
3:A:501:0HK:H20	3:A:501:0HK:O10	2.03	0.58
1:A:79:MET:SD	1:A:408:TYR:HB3	2.44	0.57
1:A:420:LEU:HD12	1:A:421:CYS:N	2.21	0.56
1:A:109:SER:HB3	1:A:378:TRP:HH2	1.72	0.55
1:A:137:ARG:NH2	2:C:1:ASP:HB3	2.22	0.55
1:A:72:LEU:O	1:A:76:THR:HG22	2.07	0.55
1:A:99:ASP:HB3	1:A:162:LEU:HD23	1.89	0.55
1:A:187:ILE:HD12	1:A:187:ILE:H	1.73	0.53
1:A:1065:LEU:O	1:A:1068:GLN:HG3	2.09	0.52
1:A:123:ARG:NH1	8:A:601:HOH:O	2.41	0.52
1:A:1010:GLU:OE2	1:A:1144:ARG:NH1	2.34	0.52
1:A:366:THR:HG21	1:A:418:TYR:HB3	1.92	0.52
1:A:416:MET:O	1:A:420:LEU:HG	2.10	0.51
1:A:434:LEU:HB3	1:A:436:ARG:HG2	1.94	0.50
1:A:1119:MET:HE3	1:A:1128:ALA:HA	1.94	0.50
1:A:115:ASN:N	1:A:115:ASN:HD22	2.10	0.49
1:A:385:VAL:O	1:A:389:THR:HG23	2.12	0.49
1:A:76:THR:HG23	1:A:77:PHE:CD2	2.47	0.49
1:A:360:GLU:HG3	2:C:3:LYS:H	1.77	0.49
1:A:372:LEU:O	1:A:376:LEU:HG	2.12	0.49
1:A:376:LEU:HD12	1:A:377:THR:HG23	1.95	0.48
1:A:207:LEU:O	1:A:210:ARG:N	2.47	0.48

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:ILE:O	1:A:379:THR:OG1	2.24	0.47
1:A:174:LEU:H	1:A:174:LEU:HD22	1.80	0.47
1:A:423:LYS:O	1:A:425:PHE:N	2.47	0.46
1:A:384:MET:HE1	1:A:400:TRP:HA	1.98	0.46
1:A:109:SER:HB3	1:A:378:TRP:CH2	2.51	0.44
1:A:28:ILE:O	1:A:32:THR:HG23	2.17	0.44
1:A:207:LEU:O	1:A:211:ILE:HD12	2.18	0.44
1:A:1116:SER:HA	1:A:1119:MET:HE2	1.99	0.44
1:A:1142:PRO:O	1:A:1146:LYS:HG3	2.18	0.44
1:A:418:TYR:O	1:A:422:ASN:CB	2.65	0.44
1:A:186:PRO:HA	1:A:189:THR:HG22	2.00	0.43
1:A:1125:TRP:HB3	1:A:1153:ARG:HA	2.00	0.43
1:A:188:ILE:O	1:A:192:THR:HG23	2.18	0.43
1:A:1082:LYS:HD2	1:A:1082:LYS:O	2.18	0.43
1:A:434:LEU:CB	1:A:436:ARG:HG2	2.49	0.42
1:A:415:PRO:O	1:A:419:ALA:HB3	2.19	0.42
1:A:33:GLY:O	1:A:37:LEU:HD12	2.19	0.42
1:A:399:LEU:HD23	1:A:399:LEU:HA	1.93	0.41
1:A:218:ARG:HH22	1:A:360:GLU:CD	2.24	0.41
1:A:396:PRO:HD2	1:A:399:LEU:HD12	2.02	0.41
1:A:82:TYR:HB2	1:A:101:TRP:HZ2	1.85	0.41
1:A:50:PHE:CE2	1:A:57:LYS:HA	2.55	0.41
1:A:86:LEU:HD11	1:A:401:GLU:HB3	2.03	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	442/515 (86%)	429 (97%)	13 (3%)	0	100	100
2	C	5/7 (71%)	3 (60%)	1 (20%)	1 (20%)	0	0

Continued on next page...

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
All	All	447/522 (86%)	432 (97%)	14 (3%)	1 (0%)	52 80

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	3	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	346/441 (78%)	334 (96%)	12 (4%)	43 74
2	C	7/7 (100%)	7 (100%)	0	100 100
All	All	353/448 (79%)	341 (97%)	12 (3%)	44 75

All (12) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	35	LEU
1	A	52	VAL
1	A	67	LEU
1	A	114	MET
1	A	115	ASN
1	A	171	ARG
1	A	1051	ARG
1	A	1060	ASP
1	A	1103	PHE
1	A	355	PHE
1	A	391	CYS
1	A	427	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1030	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	0HK	A	501	-	30,30,30	1.32	2 (6%)	32,48,48	2.31	7 (21%)
4	Y01	A	502	-	35,38,38	0.98	1 (2%)	54,57,57	1.19	3 (5%)
5	EDO	A	503	-	3,3,3	0.45	0	2,2,2	0.42	0
6	PGE	A	504	-	9,9,9	0.32	0	8,8,8	0.38	0
5	EDO	A	505	-	3,3,3	0.46	0	2,2,2	0.36	0
7	GOL	A	506	-	5,5,5	0.36	0	5,5,5	0.22	0
5	EDO	A	507	-	3,3,3	0.44	0	2,2,2	0.40	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	0HK	A	501	-	-	0/10/55/55	0/2/5/5
4	Y01	A	502	-	-	0/17/77/77	0/4/4/4
5	EDO	A	503	-	-	0/1/1/1	0/0/0/0
6	PGE	A	504	-	-	0/7/7/7	0/0/0/0
5	EDO	A	505	-	-	0/1/1/1	0/0/0/0
7	GOL	A	506	-	-	0/4/4/4	0/0/0/0
5	EDO	A	507	-	-	0/1/1/1	0/0/0/0

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	0HK	C3-N2	-5.47	1.42	1.53
3	A	501	0HK	C7-N2	3.61	1.61	1.53
4	A	502	Y01	OAW-CAY	4.41	1.47	1.34

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	0HK	C8-C7-N2	-4.43	100.64	103.56
3	A	501	0HK	O33-C30-C28	-2.35	104.26	107.80
3	A	501	0HK	C31-C30-C32	-2.30	106.49	110.62
4	A	502	Y01	CAU-CBI-CBE	-2.10	113.49	116.58
3	A	501	0HK	C9-C3-N2	2.22	105.02	103.56
3	A	501	0HK	C4-C5-C6	2.30	116.60	111.91
4	A	502	Y01	CAV-CAZ-CBH	2.76	120.41	116.41
4	A	502	Y01	OAW-CAY-CAM	3.58	119.07	111.53
3	A	501	0HK	C6-C7-N2	6.57	112.55	108.79
3	A	501	0HK	C4-C3-N2	8.31	113.54	108.79

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	0HK	1	0

## 5.7 Other polymers ⓘ

There are no such residues in this entry.



## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	444/515 (86%)	0.14	25 (5%) 28 26	50, 70, 108, 127	0
2	C	7/7 (100%)	-0.31	0 100 100	66, 77, 83, 83	0
All	All	451/522 (86%)	0.13	25 (5%) 29 27	50, 70, 108, 127	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	439	LYS	5.7
1	A	1055	GLY	5.0
1	A	1040	ALA	4.4
1	A	390	PHE	4.3
1	A	438	ASP	3.9
1	A	437	TRP	3.8
1	A	1036	PRO	3.8
1	A	1056	VAL	3.6
1	A	391	CYS	3.3
1	A	174	LEU	2.7
1	A	212	TYR	2.7
1	A	1037	SER	2.7
1	A	393	ASP	2.6
1	A	1039	ASN	2.6
1	A	1035	SER	2.5
1	A	1043	SER	2.4
1	A	1034	LYS	2.3
1	A	392	LYS	2.2
1	A	206	THR	2.2
1	A	398	THR	2.2
1	A	1050	GLY	2.1
1	A	436	ARG	2.1
1	A	1088	ASP	2.1
1	A	1024	TYR	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	186	PRO	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
5	EDO	A	505	4/4	0.92	0.32	4.21	69,72,78,86	0
6	PGE	A	504	10/10	0.77	0.35	4.11	63,81,87,88	0
5	EDO	A	507	4/4	0.91	0.22	2.60	70,74,78,80	0
4	Y01	A	502	35/35	0.90	0.20	0.69	68,71,91,92	0
3	0HK	A	501	26/26	0.94	0.23	0.50	55,65,76,76	0
7	GOL	A	506	6/6	0.82	0.24	-	82,87,89,91	0
5	EDO	A	503	4/4	0.70	0.21	-	86,90,94,95	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.