



Full wwPDB X-ray Structure Validation Report i

Feb 19, 2016 – 10:21 PM GMT

PDB ID : 5D11
Title : Kinase domain of cSrc in complex with RL235
Authors : Becker, C.; Gruetter, C.; Engel, J.; Rauh, D.
Deposited on : 2015-08-03
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026982
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20026982


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4253 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Proto-oncogene tyrosine-protein kinase Src.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
1	A	254	Total	C 2016	N 1291	O 336	S 371	18	0	3	0
1	B	255	Total	C 2063	N 1329	O 340	S 376	18	0	3	0

There are 10 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	248	GLY	-	expression tag	UNP P00523
A	249	HIS	-	expression tag	UNP P00523
A	250	MET	-	expression tag	UNP P00523
A	338	MET	THR	engineered mutation	UNP P00523
A	345	CYS	SER	engineered mutation	UNP P00523
B	248	GLY	-	expression tag	UNP P00523
B	249	HIS	-	expression tag	UNP P00523
B	250	MET	-	expression tag	UNP P00523
B	338	MET	THR	engineered mutation	UNP P00523
B	345	CYS	SER	engineered mutation	UNP P00523

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is N-[3-(4-(4-methylpiperazin-1-yl)-6-[(5-methyl-1H-pyrazol-3-yl)amino]pyrimidin-2-yl}oxy)phenyl]prop-2-enamide (three-letter code: 56G) (formula: C₂₂H₂₆N₈O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 32 22 8 2	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total C N O 32 22 8 2	0	0


- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	52	Total O 52 52	0	0
4	B	46	Total O 46 46	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



- Molecule 1: Proto-oncogene tyrosine-protein kinase Src



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	42.12 Å 63.43 Å 75.35 Å 78.65° 89.55° 90.29°	Depositor
Resolution (Å)	43.54 – 2.30 43.54 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.2 (43.54-2.30) 90.9 (43.54-2.30)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.96 (at 2.29 Å)	Xtriage
Refinement program	REFMAC	Depositor
R , R_{free}	0.214 , 0.269 0.220 , 0.269	Depositor DCC
R_{free} test set	991 reflections (3.09%)	DCC
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.006	Xtriage
Bulk solvent $k_{sol}(e/\text{\AA}^3)$, $B_{sol}(\text{\AA}^2)$	0.33 , 36.3	EDS
Estimated twinning fraction	0.036 for h,-k,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Outliers	0 of 33021 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4253	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 56G

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.92	1/2072 (0.0%)	0.94	1/2808 (0.0%)
1	B	0.87	0/2121	0.97	7/2870 (0.2%)
All	All	0.90	1/4193 (0.0%)	0.96	8/5678 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	345	CYS	CB-SG	-7.57	1.69	1.82

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	506	ARG	NE-CZ-NH1	8.81	124.71	120.30
1	B	518	ASP	CB-CG-OD1	7.49	125.04	118.30
1	B	506	ARG	NE-CZ-NH2	-6.70	116.95	120.30
1	B	518	ASP	CB-CG-OD2	-6.58	112.37	118.30
1	B	379	ARG	NE-CZ-NH1	6.29	123.45	120.30
1	A	379	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	B	359	ARG	NE-CZ-NH1	5.70	123.15	120.30
1	B	438	ARG	NE-CZ-NH1	5.17	122.89	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbit. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2016	0	1966	12	3
1	B	2063	0	2040	6	0
2	A	6	0	8	0	0
2	B	6	0	8	2	0
3	A	32	0	25	0	0
3	B	32	0	25	0	0
4	A	52	0	0	1	0
4	B	46	0	0	0	0
All	All	4253	0	4072	18	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (18) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:362:GLN:HG3	2:B:601:GOL:H32	1.62	0.79
1:A:343:LYS:HE3	1:A:396[B]:GLU:OE1	2.09	0.53
1:A:377:VAL:HG21	1:A:405:PHE:HZ	1.75	0.52
1:B:263:PRO:O	1:B:266:SER:OG	2.29	0.51
1:A:354:MET:HA	1:A:354:MET:HE3	1.94	0.50
1:A:263:PRO:O	1:A:266:SER:OG	2.31	0.48
1:A:396[B]:GLU:OE1	1:A:396[B]:GLU:HA	2.13	0.48
1:B:359:ARG:H	1:B:362:GLN:HE21	1.62	0.47
1:A:500:ARG:NH1	1:A:505:GLU:O	2.48	0.44
1:B:362:GLN:HG3	2:B:601:GOL:C3	2.42	0.43
1:A:279:GLY:HA3	1:A:296:THR:O	2.19	0.43
1:A:315:LYS:HG3	1:A:325:LEU:HD23	2.01	0.42
1:B:261:GLU:OE1	1:B:330:SER:OG	2.36	0.42
1:A:354:MET:CE	1:A:354:MET:HA	2.50	0.42
1:B:310:GLU:O	1:B:314:MET:HB2	2.20	0.41
1:A:277:CYS:O	1:A:298:LYS:NZ	2.52	0.40
1:A:520:PHE:HB3	4:A:745:HOH:O	2.20	0.40
1:A:451:LEU:HD12	1:A:451:LEU:O	2.21	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:396[B]:GLU:OE2	1:A:436:TYR:C[1_455]	1.77	0.43
1:A:396[B]:GLU:OE2	1:A:436:TYR:O[1_455]	1.98	0.22
1:A:396[B]:GLU:OE2	1:A:437:GLY:N[1_455]	2.14	0.06

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	251/286 (88%)	236 (94%)	15 (6%)	0	100 100
1	B	252/286 (88%)	241 (96%)	11 (4%)	0	100 100
All	All	503/572 (88%)	477 (95%)	26 (5%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	212/245 (86%)	194 (92%)	18 (8%)	13 16
1	B	221/245 (90%)	211 (96%)	10 (4%)	34 46
All	All	433/490 (88%)	405 (94%)	28 (6%)	22 27

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	269	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	277	CYS
1	A	291	ARG
1	A	296	THR
1	A	298	LYS
1	A	301	THR
1	A	330	SER
1	A	345	CYS
1	A	354	MET
1	A	407	LEU
1	A	426	ILE
1	A	432	GLU
1	A	451	LEU
1	A	469	ARG
1	A	474	GLN
1	A	493[A]	ASP
1	A	493[B]	ASP
1	A	524	GLU
1	B	273	LEU
1	B	310	GLU
1	B	316	LYS
1	B	330	SER
1	B	354	MET
1	B	385	ARG
1	B	388	ARG
1	B	426	ILE
1	B	504	GLU
1	B	524	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	397	ASN

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no carbohydrates in this entry.

5.6 Ligand geometry (i)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	GOL	A	601	-	5,5,5	0.45	0	5,5,5	0.55	0
3	56G	A	602	1	34,35,35	2.02	8 (23%)	41,48,48	3.15	16 (39%)
2	GOL	B	601	-	5,5,5	0.33	0	5,5,5	1.02	1 (20%)
3	56G	B	602	1	34,35,35	1.62	7 (20%)	41,48,48	3.58	16 (39%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	A	601	-	-	0/4/4/4	0/0/0/0
3	56G	A	602	1	-	0/16/28/28	0/4/4/4
2	GOL	B	601	-	-	0/4/4/4	0/0/0/0
3	56G	B	602	1	-	0/16/28/28	0/4/4/4

All (15) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	56G	NAU-NAP	-5.45	1.26	1.37
3	A	602	56G	CAN-CAL	-3.80	1.37	1.51
3	A	602	56G	CAY-NAS	-3.08	1.35	1.41
3	B	602	56G	CAY-NAS	-3.07	1.35	1.41
3	B	602	56G	NAU-NAP	-2.85	1.31	1.37
3	B	602	56G	CAN-CAL	-2.53	1.41	1.51
3	A	602	56G	CAI-CAX	-2.53	1.34	1.39

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	56G	CAX-NAU	-2.46	1.31	1.34
3	A	602	56G	CAX-NAU	-2.39	1.31	1.34
3	B	602	56G	CAN-NBF	2.01	1.49	1.46
3	A	602	56G	CAH-CAY	2.08	1.43	1.39
3	B	602	56G	CAK-CAW	2.56	1.52	1.48
3	A	602	56G	CAO-NBF	4.23	1.53	1.46
3	A	602	56G	CAA-CAK	4.36	1.51	1.30
3	B	602	56G	CAA-CAK	4.60	1.52	1.30

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	56G	CAA-CAK-CAW	-16.19	108.00	122.45
3	A	602	56G	CAA-CAK-CAW	-12.08	111.67	122.45
3	A	602	56G	N1-C2-N3	-6.25	118.67	127.80
3	B	602	56G	N1-C2-N3	-5.52	119.74	127.80
3	A	602	56G	CAI-CBA-NAP	-3.66	104.86	110.65
3	B	602	56G	C5-C4-N3	-3.48	116.33	122.28
3	A	602	56G	C5-C6-N1	-3.45	115.81	123.48
3	A	602	56G	C2-OAV-CAZ	-3.23	110.91	118.99
3	A	602	56G	CAY-NAS-CAW	-3.08	123.35	128.22
3	B	602	56G	CAI-CBA-NAP	-3.05	105.83	110.65
3	B	602	56G	C5-C6-N1	-3.05	116.71	123.48
3	A	602	56G	C5-C4-N3	-2.78	117.53	122.28
3	A	602	56G	CAF-CAY-NAS	-2.44	112.25	120.40
3	B	602	56G	CAY-NAS-CAW	-2.34	124.52	128.22
3	B	602	56G	CAL-CAN-NBF	2.03	114.39	110.59
3	B	602	56G	CAO-NBF-C4	2.03	125.16	120.45
3	B	602	56G	C5-C4-NBF	2.11	124.94	122.27
2	B	601	GOL	O3-C3-C2	2.18	121.05	109.97
3	A	602	56G	N3-C4-NBF	2.26	119.40	116.69
3	B	602	56G	OAV-C2-N3	2.48	123.88	115.98
3	A	602	56G	CAH-CAY-NAS	2.76	129.06	120.14
3	A	602	56G	CAM-CAO-NBF	2.80	115.82	110.59
3	A	602	56G	CAN-NBF-CAO	2.88	117.51	111.54
3	B	602	56G	CAL-NBE-CAM	3.23	113.89	109.50
3	A	602	56G	OAV-C2-N3	3.33	126.58	115.98
3	B	602	56G	C6-C5-C4	3.37	120.82	116.67
3	A	602	56G	C6-C5-C4	3.83	121.39	116.67
3	B	602	56G	CAO-CAM-NBE	4.14	115.27	110.76
3	A	602	56G	CAL-NBE-CAM	5.04	116.35	109.50
3	B	602	56G	CAN-CAL-NBE	5.42	116.65	110.76

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	56G	CAC-NBE-CAL	6.30	120.74	110.68
3	B	602	56G	C2-N1-C6	6.40	123.62	115.31
3	A	602	56G	C2-N1-C6	8.06	125.78	115.31

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	601	GOL	2	0

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	254/286 (88%)	0.21	12 (4%) 35 44	18, 37, 79, 98	0
1	B	255/286 (89%)	0.26	18 (7%) 19 26	18, 35, 80, 98	0
All	All	509/572 (88%)	0.24	30 (5%) 26 34	18, 36, 80, 98	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	259	ALA	5.1
1	B	407	LEU	4.2
1	A	258	ASP	3.9
1	B	269	LEU	3.6
1	B	272	LYS	3.4
1	B	277	CYS	3.3
1	A	405	PHE	3.3
1	B	278	PHE	3.2
1	A	311	ALA	3.2
1	B	328	VAL	3.0
1	B	408	ALA	3.0
1	A	300	GLY	2.9
1	B	424	PHE	2.8
1	B	290	THR	2.8
1	B	331	GLU	2.7
1	A	277	CYS	2.7
1	B	271	VAL	2.7
1	A	260	TRP	2.6
1	A	424	PHE	2.5
1	B	332	GLU	2.5
1	A	312	GLN	2.5
1	B	335	TYR	2.5
1	A	301	THR	2.4
1	B	267	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	273	LEU	2.2
1	B	308	LEU	2.2
1	B	311	ALA	2.1
1	B	333	PRO	2.1
1	A	299	PRO	2.1
1	A	404	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	56G	A	602	32/32	0.96	0.14	0.20	25,32,46,52	0
3	56G	B	602	32/32	0.96	0.10	-1.13	24,34,45,46	0
2	GOL	A	601	6/6	0.89	0.14	-	50,52,56,59	0
2	GOL	B	601	6/6	0.82	0.18	-	50,53,59,60	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.