



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:03 AM GMT

PDB ID : 3D5E
Title : Crystal structure of human plasma platelet activating factor acetylhydrolase covalently inhibited by paraoxon
Authors : Samanta, U.; Bahnson, B.J.
Deposited on : 2008-05-16
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4460 (2.10-2.10)
Ramachandran outliers	100387	4413 (2.10-2.10)
Sidechain outliers	100360	4414 (2.10-2.10)
RSRZ outliers	91569	3948 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	383	<div> <div>6%</div> <div>77%</div> <div>17%</div> <div>...</div> </div>
1	B	383	<div> <div>6%</div> <div>77%</div> <div>18%</div> <div>..</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	DEP	A	473	-	-	-	X
2	DEP	B	473	-	-	-	X

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	FMT	A	4	-	-	-	X
3	FMT	A	5	-	-	-	X
3	FMT	B	10	-	-	-	X
3	FMT	B	15	-	-	-	X
3	FMT	B	9	-	-	X	X

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6365 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Platelet-activating factor acetylhydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	373	Total	C	N	O	S	0	6	0
			3028	1938	522	555	13			
1	B	370	Total	C	N	O	S	0	3	0
			2985	1909	514	549	13			

- Molecule 2 is DIETHYL PHOSPHONATE (three-letter code: DEP) (formula: $C_4H_{11}O_3P$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	O	P	0	0
			8	4	3	1		
2	B	1	Total	C	O	P	0	0
			8	4	3	1		

- Molecule 3 is FORMIC ACID (three-letter code: FMT) (formula: CH_2O_2).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	A	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			3	1	2		
3	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	148	Total	O	0	0
			148	148		
4	B	140	Total	O	0	0
			140	140		


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Platelet-activating factor acetylhydrolase



- Molecule 1: Platelet-activating factor acetylhydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	117.21Å 78.70Å 97.34Å 90.00° 101.61° 90.00°	Depositor
Resolution (Å)	50.00 – 2.10 33.51 – 1.95	Depositor EDS
% Data completeness (in resolution range)	94.9 (50.00-2.10) 82.9 (33.51-1.95)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.87 (at 1.95Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.204 , 0.261 0.203 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	29.7	Xtriage
Anisotropy	0.212	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 62.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 52608 reflections	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6365	wwPDB-VP
Average B, all atoms (Å ²)	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.61% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: DEP, FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.08	4/3124 (0.1%)	0.94	9/4222 (0.2%)
1	B	1.01	0/3071	0.90	5/4150 (0.1%)
All	All	1.04	4/6195 (0.1%)	0.92	14/8372 (0.2%)

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	340	GLU	CB-CG	5.36	1.62	1.52
1	A	90	ASN	CB-CG	5.21	1.63	1.51
1	A	285	GLU	CD-OE2	5.18	1.31	1.25
1	A	339	LYS	CE-NZ	5.18	1.61	1.49

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	83	LEU	CA-CB-CG	5.87	128.81	115.30
1	B	406	ASP	CB-CG-OD1	5.80	123.52	118.30
1	A	264	ARG	NE-CZ-NH1	-5.75	117.42	120.30
1	A	204	LEU	CA-CB-CG	5.75	128.53	115.30
1	A	341	ARG	NE-CZ-NH2	-5.65	117.47	120.30
1	A	290	ARG	NE-CZ-NH1	5.61	123.11	120.30
1	A	157	ARG	NE-CZ-NH1	5.60	123.10	120.30
1	B	82	ARG	NE-CZ-NH2	-5.46	117.57	120.30
1	A	214	GLU	OE1-CD-OE2	-5.38	116.84	123.30
1	B	139	ARG	NE-CZ-NH1	5.34	122.97	120.30
1	A	296	ASP	CB-CG-OD1	5.23	123.01	118.30
1	B	290	ARG	NE-CZ-NH2	-5.23	117.69	120.30
1	B	240	ASP	CB-CG-OD2	5.13	122.92	118.30
1	A	304	ASP	CB-CG-OD2	5.06	122.85	118.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3028	0	2979	52	0
1	B	2985	0	2935	46	0
2	A	8	0	10	1	0
2	B	8	0	10	1	0
3	A	21	0	7	1	0
3	B	27	0	9	7	0
4	A	148	0	0	2	0
4	B	140	0	0	7	0
All	All	6365	0	5950	97	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (97) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:89:ASP:OD1	1:A:91:ASP:HB2	1.32	1.27
1:B:330:LYS:HE3	4:B:487:HOH:O	1.59	1.01
3:B:13:FMT:H	4:B:606:HOH:O	1.72	0.87
1:B:299:MET:H	1:B:327:ASN:HD21	1.25	0.83
1:A:299:MET:H	1:A:327:ASN:HD21	1.33	0.74
1:B:422:ILE:HD11	3:B:9:FMT:H	1.67	0.74
1:A:114[A]:HIS:HD2	1:A:116:LEU:HB2	1.57	0.68
1:A:114[A]:HIS:CD2	1:A:116:LEU:HB2	2.30	0.66
1:A:266[A]:LYS:HD3	1:A:396:LEU:O	1.96	0.66
1:A:213:GLU:HB3	1:A:217:ILE:HD12	1.78	0.66
1:A:307:TYR:CG	1:A:333:LYS:HG2	2.33	0.64
1:A:401:ASP:O	1:A:404:GLN:HG2	1.97	0.64
1:B:323:GLN:O	3:B:9:FMT:C	2.47	0.63
1:A:210:LYS:HB2	1:A:213:GLU:HG3	1.83	0.61
1:B:401:ASP:O	1:B:404:GLN:HG2	2.00	0.61

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:265:GLU:OE2	1:B:290:ARG:NH2	2.32	0.61
1:A:340:GLU:HG2	1:A:395:HIS:CE1	2.36	0.60
1:A:233:LEU:HD11	1:A:267:ILE:HD13	1.82	0.60
1:B:157:ARG:HB3	1:B:178:GLU:HB2	1.83	0.60
1:A:263:ASP:OD2	1:A:266[A]:LYS:NZ	2.31	0.59
1:A:93:LEU:HB3	1:A:131:PRO:HA	1.83	0.59
1:A:114[A]:HIS:CD2	1:A:117:MET:H	2.21	0.59
1:A:213:GLU:HB3	1:A:217:ILE:CD1	2.33	0.58
1:A:114[A]:HIS:HD2	1:A:116:LEU:CB	2.16	0.58
1:A:325:PRO:O	1:A:329:ILE:HG23	2.03	0.58
1:A:96:LEU:HD23	1:A:99:PRO:HA	1.86	0.58
1:B:333:LYS:NZ	4:B:609:HOH:O	2.37	0.57
1:A:330:LYS:HE3	3:A:5:FMT:O1	2.03	0.57
1:B:88:GLN:NE2	1:B:140:PRO:HD2	2.20	0.57
1:A:55:LYS:HB2	1:A:359:PHE:O	2.05	0.56
1:A:305:GLU:OE2	1:A:306:VAL:HG23	2.06	0.56
1:B:376:ASP:OD2	1:B:378:ASN:HB2	2.07	0.55
1:A:329:ILE:HG22	1:A:424:THR:HG21	1.88	0.54
1:B:88:GLN:HE21	1:B:140:PRO:HD2	1.73	0.53
1:B:323:GLN:O	3:B:9:FMT:H	2.08	0.53
1:A:424:THR:HG22	1:A:425:THR:H	1.73	0.53
1:B:273:SER:HB2	1:B:351:HIS:CE1	2.43	0.53
1:B:117:MET:CE	1:B:117:MET:HA	2.39	0.52
1:A:114[B]:HIS:HB3	1:A:116:LEU:H	1.74	0.52
1:A:216:HIS:O	1:A:220[A]:GLU:HG3	2.09	0.52
1:B:266:LYS:HD3	1:B:398:LEU:HD23	1.91	0.52
1:B:214:GLU:OE1	1:B:218:ARG:HG2	2.10	0.51
1:B:92:ARG:O	1:B:133:ASN:OD1	2.29	0.51
1:B:299:MET:H	1:B:327:ASN:ND2	2.02	0.50
1:B:182:ARG:HG2	1:B:204:LEU:HD11	1.91	0.50
1:A:101:LYS:NZ	4:A:585:HOH:O	2.45	0.50
1:B:193:GLN:O	1:B:197:GLU:HG2	2.11	0.50
1:B:89:ASP:HB3	1:B:91:ASP:OD1	2.12	0.50
1:A:214:GLU:HG3	1:A:218:ARG:HG2	1.94	0.50
1:A:90:ASN:ND2	1:A:93:LEU:HD21	2.27	0.49
1:B:93:LEU:HB3	1:B:131:PRO:HA	1.92	0.49
1:A:316:PHE:CE1	1:A:331:MET:HG2	2.47	0.49
1:B:214:GLU:OE1	1:B:218:ARG:NE	2.46	0.48
1:A:266[B]:LYS:HD3	1:A:398:LEU:HD23	1.96	0.48
1:B:345:THR:HB	1:B:417:ILE:HB	1.94	0.48
1:A:299:MET:H	1:A:327:ASN:ND2	2.08	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:330:LYS:HE2	4:B:609:HOH:O	2.13	0.47
1:A:90:ASN:HA	1:A:90:ASN:HD22	1.49	0.47
1:A:424:THR:HG22	1:A:425:THR:N	2.30	0.47
1:A:54:THR:HG21	4:A:485:HOH:O	2.14	0.46
1:A:333:LYS:HE2	1:A:333:LYS:HB2	1.66	0.46
1:A:270:ILE:HG12	1:A:293:ILE:HB	1.98	0.45
1:B:192:ASP:OD1	1:B:194:SER:N	2.49	0.45
1:A:121:LEU:HA	1:A:121:LEU:HD23	1.82	0.45
1:A:307:TYR:CB	1:A:333:LYS:HG2	2.46	0.45
1:B:383:LEU:HD11	1:B:415:ASN:HB2	1.98	0.45
1:A:173:ILE:HD12	1:A:262:ILE:HD13	1.98	0.45
1:B:422:ILE:HD11	3:B:9:FMT:C	2.43	0.45
1:B:100:ASN:ND2	4:B:504:HOH:O	2.45	0.45
1:A:294:ALA:HB1	1:A:297:ALA:HB2	1.99	0.44
1:B:294:ALA:HB1	1:B:297:ALA:HB2	1.97	0.44
1:B:100:ASN:HB2	1:B:103:TYR:CD2	2.53	0.44
1:A:259[B]:LYS:HE3	1:B:309:ARG:NH2	2.33	0.44
1:A:139:ARG:HA	1:A:140:PRO:HD2	1.82	0.43
1:B:244:PRO:O	4:B:612:HOH:O	2.20	0.43
1:B:234:SER:OG	3:B:8:FMT:H	2.19	0.43
1:B:274:PHE:HB2	2:B:473:DEP:O3	2.19	0.42
1:B:72:PHE:HA	1:B:73:ASP:HA	1.82	0.42
1:B:102:GLU:HG3	1:B:105:TRP:CZ3	2.55	0.42
1:A:88:GLN:O	1:A:89:ASP:HB3	2.20	0.42
1:A:335:TYR:HA	1:A:341:ARG:HD3	2.01	0.42
1:B:347[A]:ARG:HB3	1:B:347[A]:ARG:HE	1.63	0.42
1:A:263:ASP:CG	1:A:266[A]:LYS:HG3	2.40	0.42
1:A:113:THR:HB	1:A:114[B]:HIS:H	1.58	0.42
1:B:193:GLN:NE2	4:B:597:HOH:O	2.53	0.42
1:B:102:GLU:HG3	1:B:105:TRP:CE3	2.55	0.41
1:B:117:MET:HA	1:B:117:MET:HE2	2.00	0.41
1:A:168:ALA:HA	1:A:172:PHE:O	2.20	0.41
1:A:274:PHE:HB2	2:A:473:DEP:O3	2.20	0.41
1:B:205:TYR:H	3:B:7:FMT:C	2.33	0.41
1:B:340:GLU:C	1:B:341:ARG:HD2	2.40	0.41
1:A:290:ARG:O	1:A:313:PRO:HD2	2.21	0.41
1:B:287:GLN:HE22	1:B:311:PRO:HG2	1.84	0.41
1:B:209:LEU:HD12	1:B:214:GLU:OE1	2.21	0.41
1:B:300:PHE:N	1:B:301:PRO:CD	2.84	0.41
1:A:114[A]:HIS:CD2	1:A:117:MET:N	2.88	0.40
1:A:381:ILE:HA	1:A:381:ILE:HD12	1.82	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	377/383 (98%)	363 (96%)	13 (3%)	1 (0%)	46	45
1	B	371/383 (97%)	358 (96%)	12 (3%)	1 (0%)	46	45
All	All	748/766 (98%)	721 (96%)	25 (3%)	2 (0%)	46	45

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	91	ASP
1	B	91	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	329/329 (100%)	308 (94%)	21 (6%)	22	18
1	B	323/329 (98%)	307 (95%)	16 (5%)	30	27
All	All	652/658 (99%)	615 (94%)	37 (6%)	27	22

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	83	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	90	ASN
1	A	114[A]	HIS
1	A	114[B]	HIS
1	A	133	ASN
1	A	212	GLU
1	A	214	GLU
1	A	215	THR
1	A	222	VAL
1	A	246	LYS
1	A	252	LYS
1	A	266[A]	LYS
1	A	266[B]	LYS
1	A	274	PHE
1	A	296	ASP
1	A	304	ASP
1	A	305	GLU
1	A	329	ILE
1	A	341	ARG
1	A	370	LYS
1	A	424	THR
1	B	54	THR
1	B	55	LYS
1	B	90	ASN
1	B	91	ASP
1	B	117	MET
1	B	123	LEU
1	B	142	GLU
1	B	212	GLU
1	B	214	GLU
1	B	227	LYS
1	B	241	HIS
1	B	246	LYS
1	B	254	ASP
1	B	274	PHE
1	B	296	ASP
1	B	333	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	60	ASN
1	A	90	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	133	ASN
1	A	135	ASN
1	A	327	ASN
1	B	60	ASN
1	B	88	GLN
1	B	90	ASN
1	B	100	ASN
1	B	135	ASN
1	B	287	GLN
1	B	327	ASN
1	B	423	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	FMT	A	11	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	12	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	16	-	0,2,2	0.00	-	0,1,1	0.00	-

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	FMT	A	2	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	3	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	A	4	-	0,2,2	0.00	-	0,1,1	0.00	-
2	DEP	A	473	1	2,7,7	0.75	0	2,7,7	0.71	0
3	FMT	A	5	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	1	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	10	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	13	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	14	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	15	-	0,2,2	0.00	-	0,1,1	0.00	-
2	DEP	B	473	1	2,7,7	0.69	0	2,7,7	0.25	0
3	FMT	B	6	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	7	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	8	-	0,2,2	0.00	-	0,1,1	0.00	-
3	FMT	B	9	-	0,2,2	0.00	-	0,1,1	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	FMT	A	11	-	-	0/0/0/0	0/0/0/0
3	FMT	A	12	-	-	0/0/0/0	0/0/0/0
3	FMT	A	16	-	-	0/0/0/0	0/0/0/0
3	FMT	A	2	-	-	0/0/0/0	0/0/0/0
3	FMT	A	3	-	-	0/0/0/0	0/0/0/0
3	FMT	A	4	-	-	0/0/0/0	0/0/0/0
2	DEP	A	473	1	-	0/2/6/6	0/0/0/0
3	FMT	A	5	-	-	0/0/0/0	0/0/0/0
3	FMT	B	1	-	-	0/0/0/0	0/0/0/0
3	FMT	B	10	-	-	0/0/0/0	0/0/0/0
3	FMT	B	13	-	-	0/0/0/0	0/0/0/0
3	FMT	B	14	-	-	0/0/0/0	0/0/0/0
3	FMT	B	15	-	-	0/0/0/0	0/0/0/0
2	DEP	B	473	1	-	0/2/6/6	0/0/0/0
3	FMT	B	6	-	-	0/0/0/0	0/0/0/0
3	FMT	B	7	-	-	0/0/0/0	0/0/0/0
3	FMT	B	8	-	-	0/0/0/0	0/0/0/0
3	FMT	B	9	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

7 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	473	DEP	1	0
3	A	5	FMT	1	0
3	B	13	FMT	1	0
2	B	473	DEP	1	0
3	B	7	FMT	1	0
3	B	8	FMT	1	0
3	B	9	FMT	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	373/383 (97%)	0.12	22 (5%) 26 34	13, 26, 45, 62	0
1	B	370/383 (96%)	0.27	23 (6%) 24 32	14, 28, 49, 66	0
All	All	743/766 (96%)	0.19	45 (6%) 25 33	13, 27, 48, 66	0

All (45) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	423	ASN	8.4
1	A	115	TRP	6.1
1	B	115	TRP	5.1
1	A	54	THR	4.8
1	A	211	GLN	4.5
1	B	91	ASP	4.5
1	B	209	LEU	4.4
1	A	426	ASN	3.8
1	B	123	LEU	3.8
1	A	212	GLU	3.4
1	A	364	ILE	3.4
1	B	213	GLU	3.3
1	B	210	LYS	3.2
1	B	116	LEU	3.2
1	B	216	HIS	3.2
1	B	114	HIS	3.1
1	A	425	THR	3.0
1	A	363	LYS	2.9
1	B	212	GLU	2.9
1	A	295	LEU	2.8
1	A	368	MET	2.8
1	B	368	MET	2.7
1	B	127	SER	2.7
1	B	117	MET	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	256	GLU	2.7
1	B	149	PHE	2.6
1	A	116	LEU	2.5
1	A	123	LEU	2.5
1	A	114[A]	HIS	2.5
1	A	304	ASP	2.5
1	A	90	ASN	2.5
1	B	112	GLY	2.5
1	A	149	PHE	2.4
1	A	250	ASP	2.4
1	B	174	VAL	2.4
1	B	92	ARG	2.3
1	A	163	ILE	2.3
1	A	176	ALA	2.2
1	B	270	ILE	2.2
1	A	148	VAL	2.1
1	A	424	THR	2.1
1	B	89	ASP	2.1
1	B	364	ILE	2.1
1	B	54	THR	2.0
1	A	150	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
3	FMT	B	9	3/3	0.95	0.61	30.16	41,41,42,44	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	FMT	B	15	3/3	0.82	0.23	9.25	35,35,40,43	0
2	DEP	B	473	8/8	0.62	0.31	6.72	35,45,47,48	0
3	FMT	A	5	3/3	0.94	0.15	6.64	38,38,40,41	0
2	DEP	A	473	8/8	0.72	0.27	3.81	31,42,48,48	0
3	FMT	A	4	3/3	0.84	0.32	3.04	57,57,57,58	0
3	FMT	B	10	3/3	0.79	0.22	2.36	33,33,37,41	0
3	FMT	B	13	3/3	0.98	0.11	1.24	19,19,20,25	0
3	FMT	A	3	3/3	0.93	0.16	0.49	45,45,46,46	0
3	FMT	B	7	3/3	0.95	0.23	-	42,42,43,45	0
3	FMT	A	2	3/3	0.96	0.16	-	40,40,41,42	0
3	FMT	A	12	3/3	0.87	0.25	-	48,48,49,49	0
3	FMT	A	11	3/3	0.85	0.20	-	46,46,48,48	0
3	FMT	B	8	3/3	0.96	0.13	-	33,33,35,35	0
3	FMT	A	16	3/3	0.93	0.09	-	47,47,49,50	0
3	FMT	B	6	3/3	0.98	0.11	-	16,16,17,22	0
3	FMT	B	14	3/3	0.85	0.20	-	61,61,62,62	0
3	FMT	B	1	3/3	0.95	0.10	-	35,35,37,38	0

6.5 Other polymers [i](#)

There are no such residues in this entry.