



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 08:05 AM GMT

PDB ID : 3D8E  
Title : Crystal structure of the human Fe65-PTB1 domain (trigonal crystal form)  
Authors : Radzimanowski, J.; Ravaud, S.; Sinning, I.; Wild, K.  
Deposited on : 2008-05-23  
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	148	<div> <div>8%</div> <div>64%</div> <div>17%</div> <div>••</div> <div>17%</div> </div>
1	B	148	<div> <div>3%</div> <div>61%</div> <div>17%</div> <div>•</div> <div>19%</div> </div>
1	C	148	<div> <div>3%</div> <div>59%</div> <div>22%</div> <div>••</div> <div>18%</div> </div>
1	D	148	<div> <div>4%</div> <div>66%</div> <div>15%</div> <div>•</div> <div>18%</div> </div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 3835 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Amyloid beta A4 precursor protein-binding family B member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	123	Total	C	N	O	S	0	0	0
			970	610	171	181	8			
1	B	120	Total	C	N	O	S	0	0	0
			944	591	170	175	8			
1	C	122	Total	C	N	O	S	0	0	0
			961	604	169	180	8			
1	D	122	Total	C	N	O	S	0	0	0
			960	603	172	177	8			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	506	LEU	-	EXPRESSION TAG	UNP O00213
A	507	GLU	-	EXPRESSION TAG	UNP O00213
A	508	HIS	-	EXPRESSION TAG	UNP O00213
A	509	HIS	-	EXPRESSION TAG	UNP O00213
A	510	HIS	-	EXPRESSION TAG	UNP O00213
A	511	HIS	-	EXPRESSION TAG	UNP O00213
A	512	HIS	-	EXPRESSION TAG	UNP O00213
A	513	HIS	-	EXPRESSION TAG	UNP O00213
B	506	LEU	-	EXPRESSION TAG	UNP O00213
B	507	GLU	-	EXPRESSION TAG	UNP O00213
B	508	HIS	-	EXPRESSION TAG	UNP O00213
B	509	HIS	-	EXPRESSION TAG	UNP O00213
B	510	HIS	-	EXPRESSION TAG	UNP O00213
B	511	HIS	-	EXPRESSION TAG	UNP O00213
B	512	HIS	-	EXPRESSION TAG	UNP O00213
B	513	HIS	-	EXPRESSION TAG	UNP O00213
C	506	LEU	-	EXPRESSION TAG	UNP O00213
C	507	GLU	-	EXPRESSION TAG	UNP O00213
C	508	HIS	-	EXPRESSION TAG	UNP O00213
C	509	HIS	-	EXPRESSION TAG	UNP O00213

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	510	HIS	-	EXPRESSION TAG	UNP O00213
C	511	HIS	-	EXPRESSION TAG	UNP O00213
C	512	HIS	-	EXPRESSION TAG	UNP O00213
C	513	HIS	-	EXPRESSION TAG	UNP O00213
D	506	LEU	-	EXPRESSION TAG	UNP O00213
D	507	GLU	-	EXPRESSION TAG	UNP O00213
D	508	HIS	-	EXPRESSION TAG	UNP O00213
D	509	HIS	-	EXPRESSION TAG	UNP O00213
D	510	HIS	-	EXPRESSION TAG	UNP O00213
D	511	HIS	-	EXPRESSION TAG	UNP O00213
D	512	HIS	-	EXPRESSION TAG	UNP O00213
D	513	HIS	-	EXPRESSION TAG	UNP O00213


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1




- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1



- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1



- Molecule 1: Amyloid beta A4 precursor protein-binding family B member 1





## 4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	145.89 Å   145.89 Å   79.05 Å 90.00°   90.00°   120.00°	Depositor
Resolution (Å)	25.00 – 2.80 24.78 – 2.80	Depositor EDS
% Data completeness (in resolution range)	100.0 (25.00-2.80) 99.8 (24.78-2.80)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.34 (at 2.80 Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.240 , 0.304 0.232 , 0.289	Depositor DCC
$R_{free}$ test set	1535 reflections (11.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	66.4	Xtriage
Anisotropy	0.160	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 44.1	EDS
Estimated twinning fraction	0.031 for h,-h-k,-l	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 15409 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	3835	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	59.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 76.33 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 1.0631e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.79	0/983	0.81	0/1323
1	B	0.79	0/955	0.84	0/1283
1	C	0.83	1/974 (0.1%)	0.86	1/1312 (0.1%)
1	D	0.78	0/971	0.87	0/1305
All	All	0.80	1/3883 (0.0%)	0.84	1/5223 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	397	CYS	CB-SG	-5.06	1.73	1.81

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	422	LEU	CA-CB-CG	6.14	129.41	115.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	970	0	982	23	0
1	B	944	0	954	20	0
1	C	961	0	969	30	0
1	D	960	0	976	22	0
All	All	3835	0	3881	84	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:375:LEU:HD22	1:B:485:GLU:HG2	1.20	1.15
1:C:385:LEU:HD12	1:C:477:LEU:HD13	1.57	0.84
1:A:375:LEU:CD2	1:B:485:GLU:HG2	2.07	0.84
1:B:428:ASP:O	1:B:430:THR:HG22	1.79	0.82
1:C:375:LEU:HD22	1:D:485:GLU:HG2	1.63	0.80
1:D:506:LEU:O	1:D:507:GLU:HG2	1.81	0.79
1:B:507:GLU:H	1:B:507:GLU:CD	1.84	0.79
1:C:385:LEU:CD1	1:C:477:LEU:HD13	2.12	0.79
1:D:428:ASP:O	1:D:430:THR:HG22	1.83	0.78
1:C:402:SER:O	1:C:403:TYR:CB	2.36	0.74
1:C:375:LEU:HD22	1:D:485:GLU:CG	2.19	0.73
1:C:403:TYR:H	1:C:461:ARG:NH2	1.87	0.72
1:C:403:TYR:H	1:C:461:ARG:HH22	1.37	0.71
1:B:504:ALA:O	1:B:506:LEU:N	2.25	0.68
1:A:451:ARG:O	1:A:452:VAL:HG13	1.93	0.68
1:B:435:GLU:HG2	1:B:437:GLN:OE1	1.96	0.66
1:A:463:ARG:HG3	1:A:463:ARG:HH11	1.61	0.66
1:B:428:ASP:O	1:B:430:THR:CG2	2.44	0.66
1:B:505:GLU:O	1:B:506:LEU:HD23	1.96	0.65
1:C:474:THR:HG23	1:C:476:MET:H	1.63	0.64
1:B:437:GLN:HG2	1:B:438:SER:N	2.13	0.63
1:C:400:GLN:O	1:D:490:ASN:ND2	2.32	0.63
1:C:402:SER:O	1:C:403:TYR:HB3	1.99	0.61
1:A:420:LYS:HB3	1:A:422:LEU:HD21	1.83	0.59
1:C:385:LEU:HD12	1:C:477:LEU:CD1	2.32	0.58
1:C:471:ASP:HB3	1:C:474:THR:HG22	1.85	0.57
1:C:378:VAL:HG12	1:C:401:LEU:HD11	1.86	0.57
1:A:471:ASP:CG	1:A:474:THR:HG22	2.24	0.57
1:A:375:LEU:HD22	1:B:485:GLU:CG	2.13	0.56
1:D:505:GLU:O	1:D:506:LEU:HD23	2.06	0.56
1:D:428:ASP:O	1:D:430:THR:CG2	2.53	0.56
1:D:487:PRO:O	1:D:490:ASN:HB2	2.07	0.54
1:A:435:GLU:HG3	1:A:438:SER:OG	2.08	0.54
1:B:507:GLU:N	1:B:507:GLU:CD	2.60	0.53
1:C:506:LEU:O	1:C:507:GLU:HG3	2.09	0.51
1:D:380:MET:HE2	1:D:385:LEU:HD21	1.92	0.51
1:C:403:TYR:HB2	1:D:487:PRO:HD3	1.93	0.51

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:403:TYR:CD2	1:D:463:ARG:HD3	2.45	0.51
1:C:429:GLU:CG	1:C:502:ILE:HD11	2.42	0.50
1:A:393:ALA:HA	1:A:396:ASN:HD22	1.76	0.50
1:B:439:GLN:HA	1:B:439:GLN:NE2	2.27	0.49
1:A:370:PHE:CD1	1:A:370:PHE:N	2.78	0.49
1:B:471:ASP:CB	1:B:474:THR:HG22	2.43	0.49
1:A:451:ARG:O	1:A:452:VAL:CG1	2.61	0.49
1:A:429:GLU:HG3	1:A:502:ILE:HD11	1.95	0.48
1:C:375:LEU:CD2	1:D:485:GLU:HG2	2.39	0.48
1:C:402:SER:HB3	1:C:461:ARG:HH21	1.79	0.47
1:A:486:ALA:HB1	1:A:487:PRO:HD2	1.96	0.47
1:D:474:THR:O	1:D:475:GLN:HB2	2.14	0.47
1:C:438:SER:O	1:C:439:GLN:HB2	2.15	0.47
1:C:451:ARG:O	1:C:452:VAL:HG13	2.15	0.47
1:A:436:PRO:O	1:A:439:GLN:NE2	2.48	0.47
1:B:437:GLN:CG	1:B:438:SER:N	2.76	0.46
1:C:429:GLU:HG3	1:C:502:ILE:HD11	1.97	0.46
1:D:380:MET:HE2	1:D:385:LEU:CD2	2.45	0.46
1:C:401:LEU:HD12	1:C:481:VAL:CG2	2.46	0.45
1:C:375:LEU:HD22	1:D:485:GLU:HG3	1.96	0.45
1:C:422:LEU:HA	1:C:436:PRO:HD3	1.98	0.45
1:D:424:LEU:HD23	1:D:424:LEU:O	2.17	0.45
1:A:463:ARG:CG	1:A:463:ARG:HH11	2.27	0.45
1:A:472:LYS:HG2	1:A:473:LEU:N	2.32	0.45
1:D:457:ARG:HG2	1:D:457:ARG:O	2.18	0.44
1:D:380:MET:CE	1:D:385:LEU:CD2	2.96	0.44
1:A:392:VAL:HG12	1:A:396:ASN:HD21	1.83	0.44
1:A:429:GLU:CG	1:A:502:ILE:HD11	2.48	0.43
1:C:379:GLU:HG3	1:C:476:MET:HG3	1.99	0.43
1:D:380:MET:CE	1:D:385:LEU:HD23	2.49	0.43
1:B:504:ALA:C	1:B:506:LEU:H	2.15	0.43
1:C:402:SER:O	1:C:403:TYR:HB2	2.17	0.43
1:D:455:VAL:HG11	1:D:489:LYS:HA	2.01	0.43
1:B:506:LEU:HD23	1:B:506:LEU:HA	1.77	0.42
1:A:438:SER:O	1:A:439:GLN:HB2	2.20	0.42
1:B:373:ARG:HG3	1:B:419:GLY:O	2.19	0.42
1:C:502:ILE:O	1:C:506:LEU:HG	2.20	0.42
1:A:400:GLN:OE1	1:B:368:LYS:NZ	2.41	0.42
1:A:505:GLU:HG2	1:A:505:GLU:O	2.21	0.41
1:C:431:LEU:HD23	1:C:431:LEU:C	2.41	0.41
1:C:385:LEU:HD21	1:C:468:VAL:HG11	2.02	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:435:GLU:HA	1:B:436:PRO:HD3	1.92	0.41
1:D:438:SER:O	1:D:439:GLN:HB2	2.21	0.41
1:A:463:ARG:NH1	1:A:463:ARG:CG	2.84	0.40
1:D:489:LYS:HE2	1:D:489:LYS:HB3	1.95	0.40
1:A:426:LEU:HA	1:A:426:LEU:HD23	1.94	0.40
1:B:504:ALA:C	1:B:506:LEU:N	2.74	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	117/148 (79%)	108 (92%)	8 (7%)	1 (1%)	21	55
1	B	112/148 (76%)	102 (91%)	8 (7%)	2 (2%)	11	34
1	C	116/148 (78%)	106 (91%)	8 (7%)	2 (2%)	11	36
1	D	114/148 (77%)	108 (95%)	6 (5%)	0	100	100
All	All	459/592 (78%)	424 (92%)	30 (6%)	5 (1%)	17	50

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	505	GLU
1	A	473	LEU
1	C	505	GLU
1	B	504	ALA
1	C	498	ILE

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	107/127 (84%)	99 (92%)	8 (8%)	17	43
1	B	103/127 (81%)	89 (86%)	14 (14%)	5	14
1	C	106/127 (84%)	97 (92%)	9 (8%)	13	36
1	D	105/127 (83%)	96 (91%)	9 (9%)	13	36
All	All	421/508 (83%)	381 (90%)	40 (10%)	11	30

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	378	VAL
1	A	380	MET
1	A	395	ASN
1	A	420	LYS
1	A	422	LEU
1	A	472	LYS
1	A	473	LEU
1	A	484	CYS
1	B	378	VAL
1	B	380	MET
1	B	383	GLU
1	B	389	ARG
1	B	394	VAL
1	B	422	LEU
1	B	430	THR
1	B	442	LEU
1	B	457	ARG
1	B	458	ASP
1	B	462	GLU
1	B	474	THR
1	B	489	LYS
1	B	503	MET
1	C	368	LYS
1	C	378	VAL

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	C	395	ASN
1	C	422	LEU
1	C	430	THR
1	C	462	GLU
1	C	470	ARG
1	C	474	THR
1	C	501	LYS
1	D	378	VAL
1	D	383	GLU
1	D	391	SER
1	D	395	ASN
1	D	430	THR
1	D	435	GLU
1	D	457	ARG
1	D	474	THR
1	D	501	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	396	ASN
1	B	425	GLN
1	B	439	GLN
1	B	496	HIS
1	C	425	GLN
1	D	396	ASN
1	D	425	GLN
1	D	480	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 5.6 Ligand geometry

There are no ligands in this entry.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	123/148 (83%)	0.50	12 (9%) 10 5	41, 59, 85, 95	0
1	B	120/148 (81%)	0.23	5 (4%) 40 28	41, 59, 81, 93	0
1	C	122/148 (82%)	0.36	4 (3%) 50 38	33, 54, 77, 85	0
1	D	122/148 (82%)	0.24	6 (4%) 33 22	37, 58, 82, 94	0
All	All	487/592 (82%)	0.33	27 (5%) 29 18	33, 58, 82, 95	0

All (27) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	473	LEU	6.8
1	A	460	GLY	4.7
1	A	420	LYS	4.7
1	D	474	THR	4.1
1	D	457	ARG	3.5
1	D	458	ASP	3.3
1	A	403	TYR	3.2
1	B	457	ARG	2.9
1	B	474	THR	2.8
1	A	472	LYS	2.8
1	C	376	GLY	2.6
1	B	461	ARG	2.6
1	A	387	PRO	2.5
1	C	473	LEU	2.5
1	D	507	GLU	2.4
1	A	462	GLU	2.4
1	C	383	GLU	2.4
1	A	383	GLU	2.4
1	A	473	LEU	2.3
1	A	466	ALA	2.3
1	A	459	SER	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	376	GLY	2.2
1	B	503	MET	2.2
1	D	387	PRO	2.1
1	C	462	GLU	2.1
1	A	501	LYS	2.1
1	B	458	ASP	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.