



# wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 3, 2016 – 10:19 AM EDT

PDB ID : 5DDS  
Title : Crystal structure of aminotransferase CrmG from Actinoalloteichus sp. WH1-2216-6 in complex with PLP  
Authors : Xu, J.; Feng, Z.; Liu, J.  
Deposited on : 2015-08-25  
Resolution : 2.60 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	523	81% 15% ..
1	B	523	83% 14% ..
1	C	523	82% 14% ..
1	D	523	84% 13% ..

## 2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 16012 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CrmG.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	514	Total 3966	C 2480	N 718	O 757	S 11	0	0	0
1	B	511	Total 3954	C 2474	N 715	O 754	S 11	0	0	0
1	C	510	Total 3946	C 2470	N 714	O 751	S 11	0	0	0
1	D	512	Total 3958	C 2476	N 716	O 755	S 11	0	0	0

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C<sub>8</sub>H<sub>10</sub>NO<sub>6</sub>P).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	Total 15	C 8	N 1	O 5	P 1	0	0
2	B	1	Total 15	C 8	N 1	O 5	P 1	0	0

*Continued on next page...*

Continued from previous page...


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			15	8	1	5	1		
2	D	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is ACETIC ACID (three-letter code: ACY) (formula: C<sub>2</sub>H<sub>4</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	D	1	Total C O 4 2 2	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	33	Total O 33 33	0	0
5	B	32	Total O 32 32	0	0
5	C	20	Total O 20 20	0	0
5	D	27	Total O 27 27	0	0



- Molecule 1: CrmG



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	84.08Å 84.21Å 88.95Å 106.71° 109.16° 95.03°	Depositor
Resolution (Å)	62.02 – 2.60 62.02 – 2.60	Depositor EDS
% Data completeness (in resolution range)	92.8 (62.02-2.60) 82.0 (62.02-2.60)	Depositor EDS
$R_{merge}$	0.14	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.61Å)	Xtriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.215 , 0.262 0.218 , 0.264	Depositor DCC
$R_{free}$ test set	3146 reflections (5.37%)	DCC
Wilson B-factor (Å <sup>2</sup> )	32.4	Xtriage
Anisotropy	0.219	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 12.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.136 for -k,-h,-l	Xtriage
Reported twinning fraction	0.925 for H, K, L 0.075 for -K, -H, -L	Depositor
Outliers	0 of 61690 reflections	Xtriage
$F_o, F_c$ correlation	0.90	EDS
Total number of atoms	16012	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	34.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 15.54% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACY, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.77	0/4033	0.42	0/5463
1	B	0.76	0/4021	0.41	0/5448
1	C	0.76	0/4013	0.41	0/5437
1	D	0.75	0/4025	0.40	0/5453
All	All	0.76	0/16092	0.41	0/21801

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	1
1	D	0	2
All	All	0	7

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 7 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	160	GLY	Peptide
1	A	325	GLY	Peptide
1	B	160	GLY	Peptide
1	B	325	GLY	Peptide
1	C	325	GLY	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3966	0	3938	48	0
1	B	3954	0	3931	32	0
1	C	3946	0	3927	38	0
1	D	3958	0	3934	32	0
2	A	15	0	7	0	0
2	B	15	0	7	0	0
2	C	15	0	7	0	0
2	D	15	0	7	0	0
3	A	6	0	8	0	0
3	B	6	0	8	0	0
4	D	4	0	3	0	0
5	A	33	0	0	3	0
5	B	32	0	0	1	0
5	C	20	0	0	0	0
5	D	27	0	0	2	0
All	All	16012	0	15777	142	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

The worst 5 of 142 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:481:ARG:HG3	1:C:481:ARG:HH11	1.30	0.97
1:A:218:THR:O	1:A:224:ARG:NH2	2.03	0.91
1:A:224:ARG:HH21	1:A:224:ARG:HG3	1.41	0.82
1:A:177:GLY:O	1:A:178:ASP:HB3	1.80	0.82
1:A:178:ASP:O	1:A:182:VAL:HG23	1.83	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	510/523 (98%)	484 (95%)	21 (4%)	5 (1%)	19	39
1	B	507/523 (97%)	484 (96%)	19 (4%)	4 (1%)	24	46
1	C	506/523 (97%)	485 (96%)	17 (3%)	4 (1%)	24	46
1	D	508/523 (97%)	485 (96%)	19 (4%)	4 (1%)	24	46
All	All	2031/2092 (97%)	1938 (95%)	76 (4%)	17 (1%)	24	46

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	ASP
1	A	263	LYS
1	B	263	LYS
1	C	263	LYS
1	D	263	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	416/422 (99%)	388 (93%)	28 (7%)	20	40
1	B	416/422 (99%)	393 (94%)	23 (6%)	27	51
1	C	415/422 (98%)	387 (93%)	28 (7%)	20	40
1	D	416/422 (99%)	395 (95%)	21 (5%)	30	56
All	All	1663/1688 (98%)	1563 (94%)	100 (6%)	24	47

5 of 100 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	416	SER
1	C	194	ARG
1	D	398	ARG
1	B	424	VAL
1	B	522	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 18 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	448	GLN
1	C	34	ASN
1	D	113	ASN
1	B	360	GLN
1	B	401	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PLP	A	601	1	15,15,16	1.03	0	21,22,23	0.84	0
3	GOL	A	602	-	5,5,5	0.65	0	5,5,5	0.25	0
2	PLP	B	601	1	15,15,16	0.91	0	21,22,23	0.94	0
3	GOL	B	602	-	5,5,5	0.58	0	5,5,5	0.26	0
2	PLP	C	601	1	15,15,16	1.28	2 (13%)	21,22,23	0.85	0
2	PLP	D	601	1	15,15,16	0.78	0	21,22,23	0.99	2 (9%)
4	ACY	D	602	-	0,3,3	0.00	-	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	601	1	-	0/6/6/8	0/1/1/1
3	GOL	A	602	-	-	0/4/4/4	0/0/0/0
2	PLP	B	601	1	-	0/6/6/8	0/1/1/1
3	GOL	B	602	-	-	0/4/4/4	0/0/0/0
2	PLP	C	601	1	-	0/6/6/8	0/1/1/1
2	PLP	D	601	1	-	0/6/6/8	0/1/1/1
4	ACY	D	602	-	-	0/0/0/0	0/0/0/0

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	601	PLP	C3-C2	-3.70	1.38	1.40
2	C	601	PLP	C2-N1	2.07	1.37	1.33

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	601	PLP	C6-C5-C4	2.22	120.01	118.12
2	D	601	PLP	O3-C3-C2	2.36	120.99	117.53

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	514/523 (98%)	-0.47	0 <b>100</b>   <b>100</b>	13, 30, 62, 84	0
1	B	511/523 (97%)	-0.48	1 (0%) <b>95</b>   <b>95</b>	13, 31, 64, 87	0
1	C	510/523 (97%)	-0.49	1 (0%) <b>95</b>   <b>95</b>	13, 31, 61, 100	0
1	D	512/523 (97%)	-0.43	3 (0%) <b>90</b>   <b>88</b>	13, 32, 65, 89	0
All	All	2047/2092 (97%)	-0.47	5 (0%) <b>95</b>   <b>95</b>	13, 31, 63, 100	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	263	LYS	3.9
1	D	267	ARG	3.1
1	C	179	VAL	2.3
1	D	159	THR	2.2
1	D	243	GLU	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.

The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
2	PLP	D	601	15/16	0.95	0.14	0.09	32,35,39,40	0
3	GOL	A	602	6/6	0.86	0.12	-0.17	32,35,36,38	0
2	PLP	C	601	15/16	0.96	0.12	-0.56	28,32,40,41	0
2	PLP	B	601	15/16	0.96	0.11	-0.84	29,33,35,36	0
3	GOL	B	602	6/6	0.94	0.11	-1.28	28,29,31,31	0
2	PLP	A	601	15/16	0.96	0.09	-1.28	25,30,33,33	0
4	ACY	D	602	4/4	0.97	0.11	-1.48	29,30,31,38	0

## 6.5 Other polymers

There are no such residues in this entry.