



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:32 AM GMT

PDB ID : 2DGE  
Title : Crystal structure of oxidized cytochrome C6A from Arabidopsis thaliana  
Authors : Chida, H.; Yokoyama, T.; Kawai, F.; Nakazawa, A.; Akazaki, H.; Takayama, Y.; Hirano, T.; Suruga, K.; Satoh, T.; Yamada, S.; Kawachi, R.; Unzai, S.; Nishio, T.; Park, S.-Y.; Oku, T.  
Deposited on : 2006-03-11  
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2072 (1.50-1.50)
Clashscore	102246	2274 (1.50-1.50)
Ramachandran outliers	100387	2218 (1.50-1.50)
Sidechain outliers	100360	2216 (1.50-1.50)
RSRZ outliers	91569	2075 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	105	<div> <div>8%</div> <div>83%14%•</div> </div>
1	B	105	<div> <div>5%</div> <div>90%6%••</div> </div>
1	C	105	<div> <div>8%</div> <div>85%11%••</div> </div>
1	D	105	<div> <div>9%</div> <div>85%9%•5%</div> </div>

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 3608 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cytochrome c6.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	102	Total	C	N	O	S	0	0	0
			800	502	141	152	5			
1	B	102	Total	C	N	O	S	0	0	0
			800	502	141	152	5			
1	C	102	Total	C	N	O	S	0	0	0
			800	502	141	152	5			
1	D	100	Total	C	N	O	S	0	0	0
			786	493	139	149	5			

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	B	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	C	1	Total 43	C 34	Fe 1	N 4	O 4	0	0
3	D	1	Total 43	C 34	Fe 1	N 4	O 4	0	0



- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	61	Total	O	0	0
			61	61		
4	B	66	Total	O	0	0
			66	66		
4	C	73	Total	O	0	0
			73	73		
4	D	48	Total	O	0	0
			48	48		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



#### • Molecule 1: Cytochrome c6





#### • Molecule 1: Cytochrome c6



#### • Molecule 1: Cytochrome c6



#### • Molecule 1: Cytochrome c6



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	50.44Å 51.84Å 53.69Å 82.51° 62.08° 63.26°	Depositor
Resolution (Å)	31.47 – 1.50 31.40 – 1.50	Depositor EDS
% Data completeness (in resolution range)	100.0 (31.47-1.50) 85.7 (31.40-1.50)	Depositor EDS
$R_{merge}$	0.04	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.76 (at 1.50Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, $R_{free}$	0.186 , 0.215 0.188 , 0.215	Depositor DCC
$R_{free}$ test set	3029 reflections (5.32%)	DCC
Wilson B-factor (Å <sup>2</sup> )	15.0	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.45 , 53.0	EDS
Estimated twinning fraction	0.021 for h,h-k,h-l 0.012 for -h,-l,-k 0.018 for -h,-h+l,-h+k	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 59952 reflections	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3608	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.05% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: HEM, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.43	0/816	0.74	3/1100 (0.3%)
1	B	0.47	0/816	0.74	4/1100 (0.4%)
1	C	0.44	0/816	0.74	3/1100 (0.3%)
1	D	0.44	0/802	0.71	3/1080 (0.3%)
All	All	0.45	0/3250	0.73	13/4380 (0.3%)

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	166	ASP	CB-CG-OD2	7.67	125.20	118.30
1	C	166	ASP	CB-CG-OD2	6.85	124.47	118.30
1	B	166	ASP	CB-CG-OD2	6.74	124.37	118.30
1	A	151	ASP	CB-CG-OD2	6.49	124.14	118.30
1	A	166	ASP	CB-CG-OD2	6.40	124.06	118.30
1	B	107	ASP	CB-CG-OD2	5.95	123.66	118.30
1	B	151	ASP	CB-CG-OD2	5.74	123.47	118.30
1	C	114	ASP	CB-CG-OD2	5.62	123.36	118.30
1	C	151	ASP	CB-CG-OD2	5.53	123.27	118.30
1	B	114	ASP	CB-CG-OD2	5.42	123.18	118.30
1	A	107	ASP	CB-CG-OD2	5.36	123.12	118.30
1	D	114	ASP	CB-CG-OD2	5.12	122.90	118.30
1	D	107	ASP	CB-CG-OD2	5.05	122.84	118.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	800	0	772	5	0
1	B	800	0	772	1	0
1	C	800	0	772	3	0
1	D	786	0	756	6	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	43	0	30	1	0
3	B	43	0	30	0	0
3	C	43	0	30	1	0
3	D	43	0	30	1	0
4	A	61	0	0	0	0
4	B	66	0	0	0	0
4	C	73	0	0	0	0
4	D	48	0	0	1	0
All	All	3608	0	3192	14	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (14) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:139:PRO:HD2	1:C:142:GLN:HG3	1.69	0.73
1:C:130:MET:HB2	3:C:200:HEM:C1D	2.44	0.53
1:D:71:GLN:CG	1:D:76:GLN:HG3	2.40	0.51
1:A:81:LEU:HD21	1:D:150:GLN:HE22	1.77	0.49
1:A:130:MET:HB2	3:A:200:HEM:C1D	2.47	0.49
1:A:71:GLN:HE22	1:A:172:VAL:HG22	1.77	0.49
1:D:130:MET:HB2	3:D:200:HEM:C4D	2.47	0.48
1:A:71:GLN:NE2	1:A:172:VAL:HG13	2.27	0.48
1:C:137:CYS:O	1:C:148:ARG:NH2	2.47	0.47
1:D:106:LYS:HD2	1:D:106:LYS:HA	1.74	0.46
1:D:71:GLN:HG3	1:D:76:GLN:HG3	2.01	0.43
1:D:150:GLN:NE2	4:D:230:HOH:O	2.52	0.42
1:B:71:GLN:HE21	1:B:172:VAL:HG12	1.85	0.41

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:142:GLN:O	1:A:142:GLN:HG3	2.20	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	100/105 (95%)	98 (98%)	2 (2%)	0	100	100
1	B	100/105 (95%)	96 (96%)	4 (4%)	0	100	100
1	C	100/105 (95%)	99 (99%)	1 (1%)	0	100	100
1	D	98/105 (93%)	96 (98%)	2 (2%)	0	100	100
All	All	398/420 (95%)	389 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	84/87 (97%)	77 (92%)	7 (8%)	14	1
1	B	84/87 (97%)	80 (95%)	4 (5%)	31	5
1	C	84/87 (97%)	78 (93%)	6 (7%)	18	1
1	D	82/87 (94%)	77 (94%)	5 (6%)	23	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	334/348 (96%)	312 (93%)	22 (7%)	21 2

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	87	ILE
1	A	92	THR
1	A	106	LYS
1	A	140	ARG
1	A	152	GLU
1	A	155	LYS
1	A	156	LEU
1	B	71	GLN
1	B	152	GLU
1	B	155	LYS
1	B	172	VAL
1	C	71	GLN
1	C	77	ARG
1	C	121	ARG
1	C	148	ARG
1	C	171	THR
1	C	172	VAL
1	D	71	GLN
1	D	114	ASP
1	D	121	ARG
1	D	138	THR
1	D	140	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	71	GLN
1	A	83	ASN
1	A	111	ASN
1	B	71	GLN
1	B	83	ASN
1	B	111	ASN
1	C	71	GLN
1	C	76	GLN
1	C	83	ASN
1	C	111	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	D	83	ASN
1	D	111	ASN
1	D	150	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 2 are monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	HEM	A	200	1	30,50,50	2.17	8 (26%)	24,82,82	2.87	12 (50%)
3	HEM	B	200	1	30,50,50	2.24	8 (26%)	24,82,82	3.00	10 (41%)
3	HEM	C	200	1	30,50,50	2.24	10 (33%)	24,82,82	2.76	10 (41%)
3	HEM	D	200	1	30,50,50	2.46	8 (26%)	24,82,82	2.81	9 (37%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	HEM	A	200	1	-	0/10/54/54	0/0/8/8
3	HEM	B	200	1	-	0/10/54/54	0/0/8/8
3	HEM	C	200	1	-	0/10/54/54	0/0/8/8
3	HEM	D	200	1	-	0/10/54/54	0/0/8/8

All (34) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	D	200	HEM	C3B-C4B	-8.43	1.44	1.51
3	B	200	HEM	C3B-C4B	-6.97	1.45	1.51
3	C	200	HEM	C3B-C4B	-6.42	1.46	1.51
3	A	200	HEM	C3B-C4B	-5.89	1.46	1.51
3	D	200	HEM	C3D-C4D	-5.60	1.44	1.51
3	C	200	HEM	C3D-C4D	-5.49	1.44	1.51
3	B	200	HEM	C3D-C4D	-4.99	1.45	1.51
3	A	200	HEM	C3D-C4D	-4.93	1.45	1.51
3	A	200	HEM	C2C-C1C	-4.32	1.44	1.52
3	C	200	HEM	C2C-C1C	-4.31	1.44	1.52
3	B	200	HEM	C2C-C1C	-3.96	1.45	1.52
3	D	200	HEM	C2C-C1C	-3.86	1.45	1.52
3	B	200	HEM	C2D-C1D	-2.38	1.44	1.51
3	C	200	HEM	C2D-C1D	-2.34	1.44	1.51
3	A	200	HEM	C2D-C1D	-2.31	1.44	1.51
3	B	200	HEM	C2B-C1B	-2.30	1.44	1.51
3	A	200	HEM	C2B-C1B	-2.20	1.44	1.51
3	D	200	HEM	C2D-C1D	-2.20	1.44	1.51
3	D	200	HEM	C2B-C1B	-2.17	1.44	1.51
3	C	200	HEM	C2B-C1B	-2.13	1.44	1.51
3	C	200	HEM	C1C-NC	2.09	1.38	1.36
3	D	200	HEM	FE-NC	2.28	2.04	1.95
3	A	200	HEM	C3B-CAB	2.36	1.55	1.51
3	C	200	HEM	FE-NC	2.41	2.05	1.95
3	C	200	HEM	C3B-CAB	2.41	1.55	1.51
3	A	200	HEM	C3C-CAC	2.57	1.56	1.51
3	C	200	HEM	C3C-CAC	2.72	1.56	1.51
3	B	200	HEM	C3C-CAC	2.77	1.56	1.51
3	C	200	HEM	C4C-NC	2.81	1.39	1.36
3	B	200	HEM	C4C-NC	2.94	1.39	1.36
3	D	200	HEM	C3C-CAC	2.98	1.56	1.51
3	B	200	HEM	FE-NC	2.98	2.07	1.95
3	D	200	HEM	C4C-NC	3.54	1.40	1.36
3	A	200	HEM	C4C-NC	4.29	1.41	1.36

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	200	HEM	C3B-CAB-CBB	-7.61	112.79	124.46
3	D	200	HEM	C3B-CAB-CBB	-7.01	113.70	124.46
3	C	200	HEM	C3B-CAB-CBB	-6.46	114.55	124.46
3	A	200	HEM	C3B-CAB-CBB	-6.15	115.03	124.46
3	B	200	HEM	C3C-CAC-CBC	-5.52	115.98	124.46
3	C	200	HEM	C3C-CAC-CBC	-4.64	117.33	124.46
3	D	200	HEM	C3C-CAC-CBC	-3.93	118.42	124.46
3	A	200	HEM	CBA-CAA-C2A	-3.54	106.19	112.53
3	A	200	HEM	C3C-CAC-CBC	-3.31	119.38	124.46
3	D	200	HEM	CBA-CAA-C2A	-2.86	107.40	112.53
3	B	200	HEM	CBA-CAA-C2A	-2.85	107.42	112.53
3	A	200	HEM	CBD-CAD-C3D	-2.62	105.92	113.55
3	B	200	HEM	CAA-C2A-C1A	-2.61	124.17	127.01
3	A	200	HEM	CMA-C3A-C4A	-2.58	124.10	128.36
3	C	200	HEM	CBA-CAA-C2A	-2.45	108.13	112.53
3	B	200	HEM	CMA-C3A-C4A	-2.31	124.54	128.36
3	A	200	HEM	CAA-C2A-C1A	-2.17	124.65	127.01
3	C	200	HEM	C3B-C4B-CHC	2.17	126.21	123.16
3	D	200	HEM	C2D-C3D-C4D	2.23	105.28	101.50
3	D	200	HEM	CMD-C2D-C3D	2.48	125.31	114.35
3	C	200	HEM	C2D-C3D-C4D	2.49	105.72	101.50
3	A	200	HEM	C3B-C4B-CHC	2.57	126.78	123.16
3	B	200	HEM	CMD-C2D-C3D	2.59	125.81	114.35
3	A	200	HEM	CMD-C2D-C3D	2.76	126.55	114.35
3	C	200	HEM	CMD-C2D-C3D	2.78	126.62	114.35
3	C	200	HEM	CAD-C3D-C2D	3.95	124.58	113.22
3	D	200	HEM	CAD-C3D-C4D	4.10	126.94	112.47
3	D	200	HEM	CMC-C2C-C3C	4.26	127.18	116.53
3	C	200	HEM	CMC-C2C-C3C	4.35	127.40	116.53
3	B	200	HEM	CMC-C2C-C3C	4.36	127.41	116.53
3	B	200	HEM	CAD-C3D-C4D	4.38	127.91	112.47
3	C	200	HEM	CMB-C2B-C3B	4.48	127.72	116.53
3	A	200	HEM	CMC-C2C-C3C	4.48	127.72	116.53
3	B	200	HEM	CMB-C2B-C3B	4.58	127.97	116.53
3	D	200	HEM	CMB-C2B-C3B	4.62	128.07	116.53
3	A	200	HEM	CAD-C3D-C2D	4.64	126.55	113.22
3	C	200	HEM	CAD-C3D-C4D	4.88	129.70	112.47
3	A	200	HEM	CAD-C3D-C4D	4.91	129.80	112.47
3	B	200	HEM	CAD-C3D-C2D	5.00	127.59	113.22
3	A	200	HEM	CMB-C2B-C3B	5.03	129.09	116.53
3	D	200	HEM	CAD-C3D-C2D	5.07	127.78	113.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	200	HEM	1	0
3	C	200	HEM	1	0
3	D	200	HEM	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	102/105 (97%)	0.47	8 (7%) 16 15	13, 19, 28, 33	0
1	B	102/105 (97%)	0.31	5 (4%) 33 34	11, 19, 28, 33	0
1	C	102/105 (97%)	0.40	8 (7%) 16 15	10, 17, 30, 40	0
1	D	100/105 (95%)	0.50	9 (9%) 12 11	11, 19, 32, 40	0
All	All	406/420 (96%)	0.42	30 (7%) 17 17	10, 19, 30, 40	0

All (30) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	145	PHE	8.2
1	C	145	PHE	6.7
1	D	140	ARG	6.0
1	C	140	ARG	5.5
1	A	171	THR	5.4
1	C	172	VAL	5.2
1	A	136	LYS	4.5
1	D	141	GLY	4.5
1	D	138	THR	3.7
1	A	172	VAL	3.4
1	A	135	GLU	3.4
1	C	171	THR	3.4
1	C	71	GLN	3.1
1	A	140	ARG	3.1
1	B	171	THR	2.9
1	B	172	VAL	2.7
1	C	139	PRO	2.7
1	C	106	LYS	2.6
1	D	114	ASP	2.6
1	D	71	GLN	2.4
1	A	71	GLN	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	138	THR	2.3
1	A	138	THR	2.3
1	D	110	ARG	2.3
1	D	106	LYS	2.2
1	B	136	LYS	2.1
1	A	87	ILE	2.1
1	B	114	ASP	2.1
1	C	84	ARG	2.1
1	D	84	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
3	HEM	A	200	43/43	0.98	0.10	0.27	12,14,19,22	0
3	HEM	C	200	43/43	0.99	0.09	0.01	8,11,14,19	0
3	HEM	B	200	43/43	0.98	0.09	-0.01	10,13,19,22	0
3	HEM	D	200	43/43	0.98	0.09	-0.28	9,11,17,18	0
2	ZN	B	1002	1/1	1.00	0.07	-	12,12,12,12	0
2	ZN	A	1001	1/1	1.00	0.06	-	13,13,13,13	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.