



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:10 AM GMT

PDB ID : 3DLS
Title : Crystal structure of human PAS kinase bound to ADP
Authors : Antonysamy, S.; Bonanno, J.B.; Romero, R.; Russell, M.; Iizuka, M.; Gheyi, T.; Wasserman, S.R.; Rutter, J.; Sauder, J.M.; Burley, S.K.; New York SGX Research Center for Structural Genomics (NYSGXRC)
Deposited on : 2008-06-29
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	335	 73% 11% • 15%
1	B	335	 68% 14% • 16%
1	C	335	 69% 13% • 16%
1	D	335	 67% 16% • 15%
1	E	335	 68% 15% • 16%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	335	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	MG	A	20	-	-	-	X
2	MG	A	25	-	-	-	X
2	MG	C	21	-	-	-	X

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 13724 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called PAS domain-containing serine/threonine-protein kinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	285	2236	1465	350	411	10	0	0	0
1	B	280	2200	1435	353	403	9	0	0	0
1	C	281	2217	1447	355	407	8	0	0	0
1	D	284	2220	1453	351	406	10	0	0	0
1	E	282	2216	1449	353	405	9	0	0	0
1	F	281	2174	1420	345	400	9	0	0	0

There are 66 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	974	MET	-	expression tag	UNP Q96RG2
A	975	ALA	-	expression tag	UNP Q96RG2
A	976	LEU	-	expression tag	UNP Q96RG2
A	1301	GLU	-	expression tag	UNP Q96RG2
A	1302	GLY	-	expression tag	UNP Q96RG2
A	1303	HIS	-	expression tag	UNP Q96RG2
A	1304	HIS	-	expression tag	UNP Q96RG2
A	1305	HIS	-	expression tag	UNP Q96RG2
A	1306	HIS	-	expression tag	UNP Q96RG2
A	1307	HIS	-	expression tag	UNP Q96RG2
A	1308	HIS	-	expression tag	UNP Q96RG2
B	974	MET	-	expression tag	UNP Q96RG2
B	975	ALA	-	expression tag	UNP Q96RG2
B	976	LEU	-	expression tag	UNP Q96RG2
B	1301	GLU	-	expression tag	UNP Q96RG2
B	1302	GLY	-	expression tag	UNP Q96RG2
B	1303	HIS	-	expression tag	UNP Q96RG2

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	1304	HIS	-	expression tag	UNP Q96RG2
B	1305	HIS	-	expression tag	UNP Q96RG2
B	1306	HIS	-	expression tag	UNP Q96RG2
B	1307	HIS	-	expression tag	UNP Q96RG2
B	1308	HIS	-	expression tag	UNP Q96RG2
C	974	MET	-	expression tag	UNP Q96RG2
C	975	ALA	-	expression tag	UNP Q96RG2
C	976	LEU	-	expression tag	UNP Q96RG2
C	1301	GLU	-	expression tag	UNP Q96RG2
C	1302	GLY	-	expression tag	UNP Q96RG2
C	1303	HIS	-	expression tag	UNP Q96RG2
C	1304	HIS	-	expression tag	UNP Q96RG2
C	1305	HIS	-	expression tag	UNP Q96RG2
C	1306	HIS	-	expression tag	UNP Q96RG2
C	1307	HIS	-	expression tag	UNP Q96RG2
C	1308	HIS	-	expression tag	UNP Q96RG2
D	974	MET	-	expression tag	UNP Q96RG2
D	975	ALA	-	expression tag	UNP Q96RG2
D	976	LEU	-	expression tag	UNP Q96RG2
D	1301	GLU	-	expression tag	UNP Q96RG2
D	1302	GLY	-	expression tag	UNP Q96RG2
D	1303	HIS	-	expression tag	UNP Q96RG2
D	1304	HIS	-	expression tag	UNP Q96RG2
D	1305	HIS	-	expression tag	UNP Q96RG2
D	1306	HIS	-	expression tag	UNP Q96RG2
D	1307	HIS	-	expression tag	UNP Q96RG2
D	1308	HIS	-	expression tag	UNP Q96RG2
E	974	MET	-	expression tag	UNP Q96RG2
E	975	ALA	-	expression tag	UNP Q96RG2
E	976	LEU	-	expression tag	UNP Q96RG2
E	1301	GLU	-	expression tag	UNP Q96RG2
E	1302	GLY	-	expression tag	UNP Q96RG2
E	1303	HIS	-	expression tag	UNP Q96RG2
E	1304	HIS	-	expression tag	UNP Q96RG2
E	1305	HIS	-	expression tag	UNP Q96RG2
E	1306	HIS	-	expression tag	UNP Q96RG2
E	1307	HIS	-	expression tag	UNP Q96RG2
E	1308	HIS	-	expression tag	UNP Q96RG2
F	974	MET	-	expression tag	UNP Q96RG2
F	975	ALA	-	expression tag	UNP Q96RG2
F	976	LEU	-	expression tag	UNP Q96RG2
F	1301	GLU	-	expression tag	UNP Q96RG2

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	1302	GLY	-	expression tag	UNP Q96RG2
F	1303	HIS	-	expression tag	UNP Q96RG2
F	1304	HIS	-	expression tag	UNP Q96RG2
F	1305	HIS	-	expression tag	UNP Q96RG2
F	1306	HIS	-	expression tag	UNP Q96RG2
F	1307	HIS	-	expression tag	UNP Q96RG2
F	1308	HIS	-	expression tag	UNP Q96RG2

- Molecule 2 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	D	3	Total Mg 3 3	0	0
2	E	2	Total Mg 2 2	0	0
2	B	4	Total Mg 4 4	0	0
2	C	4	Total Mg 4 4	0	0
2	A	4	Total Mg 4 4	0	0
2	F	2	Total Mg 2 2	0	0

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	B	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	C	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	D	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	E	1	Total	C	N	O	P	0	0
			27	10	5	10	2		
3	F	1	Total	C	N	O	P	0	0
			27	10	5	10	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	55	Total	O	0	0
			55	55		
4	B	57	Total	O	0	0
			57	57		
4	C	47	Total	O	0	0
			47	47		
4	D	48	Total	O	0	0
			48	48		
4	E	39	Total	O	0	0
			39	39		
4	F	34	Total	O	0	0
			34	34		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: PAS domain-containing serine/threonine-protein kinase



- Molecule 1: PAS domain-containing serine/threonine-protein kinase



- Molecule 1: PAS domain-containing serine/threonine-protein kinase



MET
GLY
ASN
ARG
SER
LEU
SER
SER
ASP
VAL
ALA
GLN
ALA
GLN
GLU
LEU
CYS
GLY
GLY
GLY
GLY
HIS
HIS
HIS
HIS
HIS

• Molecule 1: PAS domain-containing serine/threonine-protein kinase



MET
ALA
LEU
GLU
GLU
PRO
PRO
LYS
A982
I985
F1010
M1022
I1030
K1031
K1032
E1033
K1034
W1035
G1181
L1036
D1038
G1039
W1040
I1041
L1046
F1197
F1198
E1199
K1048
V1049
T1050
L1051
E1052
I1053
I1065
I1070
F1071
E1072
M1073
Q1074
F1077
Q1078
L1079
I1094
D1095
R1096
H1097
P1098
R1099
L1100
I1108

G1117
K1122
I1125
H1126
R1127
I1142
G1148
S1149
L1153
K1157
I1166
P1171
R1180
G1181
P1182
M1186
L1196
V1197
F1198
E1199
F1203
L1206
T1209
V1210
A1212
L1219
L1231
T1240
T1241
L1242
E1243
K1244
L1245
A1258
F1266
ARG
VAL
ASN
LYS
PRO

GLU
SER
GLY
VAL
LEU
SER
ALA
ALA
SER
LEU
GLU
MET
GLY
ASN
ARG
SER
LEU
SER
ASP
VAL
ALA
GLN
ALA
GLN
GLU
LEU
CYS
GLY
GLY
GLY
GLY
HIS
HIS
HIS

• Molecule 1: PAS domain-containing serine/threonine-protein kinase



MET
ALA
LEU
GLU
GLU
PRO
PRO
LYS
A982
I985
G1006
S1007
G1008
A1009
D1018
M1022
I1030
K1031
K1032
V1035
LEU
LEU
ASP
CYS
GLY
GLY
GLY
GLY
HIS
HIS
HIS
HIS

I1121
K1122
D1123
I1124
H1125
H1126
R1127
I1142
K1143
L1144
S1149
L1153
E1154
R1155
G1156
K1157
I1166
L1174
P1182
M1186
L1196
V1197
F1198
E1199
E1200
T1209
I1214
H1215
P1216
P1217
L1218
L1219
E1237
T1241
L1242
E1243
K1244
L1245
V1246
T1247
D1248
P1249
F1266
R1267
VAL
ASN
LYS

PRO
GLU
SER
GLY
VAL
LEU
SER
MET
GLY
ASN
ARG
SER
LEU
SER
SER
ASP
VAL
ALA
GLN
ALA
ALA
GLN
GLN
GLU
LEU
CYS
GLY
GLY
GLY
GLY
HIS
HIS
HIS
HIS

• Molecule 1: PAS domain-containing serine/threonine-protein kinase



MET
ALA
LEU
GLU
GLU
PRO
PRO
LYS
A982
I985
I988
A989
C991
Y995
A1009
V1017
I1030
K1031
K1032
V1035
LEU
LEU
GLU
GLU
ASP
CYS
W1040
I1041
E1042
K1048
L1051
I1055
I1055
D1069
I1070
F1071
E1072
F1077
Q1078
L1079
I1094
L1100
L1121
K1122
D1123
I1124
I1125

H1126
R1127
S1149
L1153
I1166
P1171
M1175
P1182
M1186
L1196
F1203
C1204
E1205
L1206
E1207
T1209
V1210
A1212
L1219
V1235
T1241
K1244
L1245
L1257
Y1260
F1266
ARG
VAL
ASN
LYS
PRO
GLU
SER
GLY
VAL
LEU
SER
SER
ALA
ALA
SER
LEU
LEU
GLU

MET
GLY
ASN
ARG
SER
LEU
SER
SER
ASP
VAL
ALA
GLN
ALA
GLN
GLU
LEU
CYS
GLY
GLY
GLY
GLY
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	85.78Å 85.84Å 94.15Å 77.28° 77.50° 60.09°	Depositor
Resolution (Å)	(Not available) – 2.30 31.20 – 2.30	Depositor EDS
% Data completeness (in resolution range)	(Not available) ((Not available)-2.30) 91.7 (31.20-2.30)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.42 (at 2.29Å)	Xtriage
Refinement program	REFMAC5	Depositor
R, R_{free}	0.241 , 0.297 0.240 , 0.295	Depositor DCC
R_{free} test set	4846 reflections (5.27%)	DCC
Wilson B-factor (Å ²)	40.0	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	(Not available) , (Not available)	EDS
Estimated twinning fraction	0.216 for -k,-h,-l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.40$, $\langle L^2 \rangle = 0.22$	Xtriage
Outliers	0 of 96810 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	13724	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.46% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2291	0.61	0/3118
1	B	0.46	0/2252	0.61	2/3062 (0.1%)
1	C	0.46	0/2269	0.58	0/3083
1	D	0.43	0/2274	0.59	0/3094
1	E	0.42	0/2270	0.57	0/3088
1	F	0.42	0/2226	0.55	0/3033
All	All	0.44	0/13582	0.59	2/18478 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	1005	LEU	O-C-N	6.14	133.64	123.20
1	B	1005	LEU	CA-C-N	-5.30	105.60	116.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2236	0	2198	21	0
1	B	2200	0	2168	27	0
1	C	2217	0	2190	29	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	2220	0	2178	31	0
1	E	2216	0	2178	28	0
1	F	2174	0	2109	18	0
2	A	4	0	0	0	0
2	B	4	0	0	0	0
2	C	4	0	0	0	0
2	D	3	0	0	0	0
2	E	2	0	0	0	0
2	F	2	0	0	0	0
3	A	27	0	12	1	0
3	B	27	0	12	0	0
3	C	27	0	12	0	0
3	D	27	0	12	1	0
3	E	27	0	12	2	0
3	F	27	0	12	0	0
4	A	55	0	0	1	0
4	B	57	0	0	0	0
4	C	47	0	0	2	0
4	D	48	0	0	0	0
4	E	39	0	0	2	0
4	F	34	0	0	0	0
All	All	13724	0	13093	153	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:1125:ILE:HG23	1:C:1127:ARG:HG3	1.67	0.74
1:D:1032:LYS:HZ1	1:D:1074:GLN:HA	1.51	0.74
1:A:1240:THR:HG23	1:A:1244:LYS:HD2	1.70	0.73
1:A:1125:ILE:HG23	1:A:1127:ARG:HG3	1.72	0.71
1:E:1094:ILE:HD11	1:E:1196:LEU:HA	1.71	0.71

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	283/335 (84%)	272 (96%)	11 (4%)	0	100	100
1	B	276/335 (82%)	264 (96%)	12 (4%)	0	100	100
1	C	277/335 (83%)	260 (94%)	17 (6%)	0	100	100
1	D	280/335 (84%)	271 (97%)	9 (3%)	0	100	100
1	E	278/335 (83%)	268 (96%)	9 (3%)	1 (0%)	39	48
1	F	277/335 (83%)	260 (94%)	17 (6%)	0	100	100
All	All	1671/2010 (83%)	1595 (96%)	75 (4%)	1 (0%)	56	68

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	1009	ALA

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/291 (81%)	218 (92%)	18 (8%)	16	20
1	B	233/291 (80%)	223 (96%)	10 (4%)	35	47
1	C	235/291 (81%)	224 (95%)	11 (5%)	32	43
1	D	234/291 (80%)	218 (93%)	16 (7%)	20	25
1	E	234/291 (80%)	217 (93%)	17 (7%)	17	22
1	F	226/291 (78%)	211 (93%)	15 (7%)	21	27

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1398/1746 (80%)	1311 (94%)	87 (6%)	23 30

5 of 87 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	985	LEU
1	D	1125	ILE
1	F	1100	LEU
1	D	1022	ASN
1	D	1078	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 9 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	997	GLN
1	F	1253	GLN
1	D	1078	GLN
1	B	997	GLN
1	C	1078	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 25 ligands modelled in this entry, 19 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link

column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	ADP	A	1	2	22,29,29	1.07	2 (9%)	27,45,45	1.84	3 (11%)
3	ADP	B	2	2	22,29,29	1.10	2 (9%)	27,45,45	1.83	4 (14%)
3	ADP	C	3	2	22,29,29	1.07	2 (9%)	27,45,45	1.77	2 (7%)
3	ADP	D	4	2	22,29,29	1.10	2 (9%)	27,45,45	1.63	2 (7%)
3	ADP	E	5	2	22,29,29	1.13	2 (9%)	27,45,45	1.74	3 (11%)
3	ADP	F	6	2	22,29,29	1.13	2 (9%)	27,45,45	1.62	2 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	ADP	A	1	2	-	0/12/32/32	0/3/3/3
3	ADP	B	2	2	-	0/12/32/32	0/3/3/3
3	ADP	C	3	2	-	0/12/32/32	0/3/3/3
3	ADP	D	4	2	-	0/12/32/32	0/3/3/3
3	ADP	E	5	2	-	0/12/32/32	0/3/3/3
3	ADP	F	6	2	-	0/12/32/32	0/3/3/3

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	6	ADP	O4'-C1'	2.08	1.43	1.41
3	E	5	ADP	O4'-C1'	2.13	1.43	1.41
3	D	4	ADP	O4'-C1'	2.40	1.44	1.41
3	A	1	ADP	O4'-C1'	2.40	1.44	1.41
3	B	2	ADP	O4'-C1'	2.51	1.44	1.41

The worst 5 of 16 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	3	ADP	N3-C2-N1	-7.80	122.92	128.89
3	A	1	ADP	N3-C2-N1	-7.65	123.03	128.89
3	B	2	ADP	N3-C2-N1	-7.54	123.12	128.89
3	E	5	ADP	N3-C2-N1	-7.17	123.41	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	D	4	ADP	N3-C2-N1	-6.71	123.76	128.89

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1	ADP	1	0
3	D	4	ADP	1	0
3	E	5	ADP	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	285/335 (85%)	-0.09	1 (0%) 93 95	14, 29, 46, 59	0
1	B	280/335 (83%)	0.07	7 (2%) 61 70	17, 33, 53, 72	0
1	C	281/335 (83%)	0.00	6 (2%) 67 74	17, 32, 54, 73	0
1	D	284/335 (84%)	0.05	3 (1%) 82 86	22, 38, 56, 70	0
1	E	282/335 (84%)	0.18	6 (2%) 67 74	22, 42, 58, 73	0
1	F	281/335 (83%)	0.29	11 (3%) 43 52	26, 46, 67, 76	0
All	All	1693/2010 (84%)	0.08	34 (2%) 68 75	14, 37, 59, 76	0

The worst 5 of 34 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1204	CYS	5.0
1	C	1046	LEU	4.7
1	F	1121	LEU	4.2
1	F	1041	ILE	3.7
1	B	1040	TRP	3.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
2	MG	C	21	1/1	0.97	0.27	4.20	30,30,30,30	0
2	MG	A	20	1/1	0.89	0.18	3.11	24,24,24,24	0
2	MG	A	25	1/1	0.97	0.23	2.38	22,22,22,22	0
2	MG	B	19	1/1	0.90	0.20	1.07	33,33,33,33	0
2	MG	B	24	1/1	0.98	0.21	0.72	26,26,26,26	0
3	ADP	D	4	27/27	0.96	0.13	-0.27	22,32,37,43	0
3	ADP	B	2	27/27	0.97	0.13	-0.28	12,23,34,41	0
3	ADP	E	5	27/27	0.94	0.14	-0.36	27,36,52,53	0
3	ADP	C	3	27/27	0.98	0.12	-0.37	17,21,30,36	0
3	ADP	A	1	27/27	0.98	0.11	-0.52	13,20,33,36	0
2	MG	D	23	1/1	0.96	0.12	-0.89	31,31,31,31	0
2	MG	C	22	1/1	0.95	0.09	-1.18	35,35,35,35	0
3	ADP	F	6	27/27	0.96	0.10	-1.35	34,40,45,46	0
2	MG	B	10	1/1	0.96	0.18	-	23,23,23,23	0
2	MG	D	13	1/1	0.99	0.20	-	14,14,14,14	0
2	MG	A	7	1/1	0.96	0.14	-	10,10,10,10	0
2	MG	F	17	1/1	0.96	0.24	-	25,25,25,25	0
2	MG	C	12	1/1	0.93	0.30	-	35,35,35,35	0
2	MG	E	16	1/1	0.94	0.25	-	36,36,36,36	0
2	MG	C	11	1/1	0.99	0.20	-	13,13,13,13	0
2	MG	A	8	1/1	0.93	0.27	-	28,28,28,28	0
2	MG	D	14	1/1	0.95	0.29	-	33,33,33,33	0
2	MG	B	9	1/1	0.98	0.16	-	15,15,15,15	0
2	MG	E	15	1/1	0.95	0.21	-	24,24,24,24	0
2	MG	F	18	1/1	0.94	0.24	-	31,31,31,31	0

6.5 Other polymers [i](#)

There are no such residues in this entry.