



wwPDB X-ray Structure Validation Summary Report ⓘ

Jan 31, 2016 – 07:00 PM GMT

PDB ID : 1DNK
Title : THE X-RAY STRUCTURE OF THE DNASE I-D(GGTATACC)2 COM-
PLEX AT 2.3 ANGSTROMS RESOLUTION
Authors : Weston, S.A.; Lahm, A.; Suck, D.
Deposited on : 1992-08-10
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	B	7	<div> <div style="width: 71%; background-color: yellow;"></div> <div style="width: 29%; background-color: orange;"></div> </div> <div>71% 29%</div>
2	C	8	<div> <div style="width: 50%; background-color: yellow;"></div> <div style="width: 50%; background-color: orange;"></div> </div> <div>50% 50%</div>
3	A	260	<div> <div style="width: 62%; background-color: green;"></div> <div style="width: 28%; background-color: yellow;"></div> <div style="width: 5%; background-color: red;"></div> <div style="width: 5%; background-color: grey;"></div> </div> <div>62% 28% 5% . .</div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NAG	A	261	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 2390 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*GP*GP*TP*AP*TP*AP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	B	7	Total	C	N	O	P	0	0	0
			142	69	27	40	6			

- Molecule 2 is a DNA chain called DNA (5'-D(*GP*GP*TP*AP*TP*AP*CP*C)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	8	Total	C	N	O	P	0	0	0
			161	78	30	46	7			

- Molecule 3 is a protein called PROTEIN (DEOXYRIBONUCLEASE I (DNASE I) (E.C.3.1.21.1)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	A	250	Total	C	N	O	S	0	0	0
			1982	1264	330	382	6			

- Molecule 4 is a polymer of unknown type called SUGAR (2-MER).

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	2	Total	C	N	O	0	0
			28	16	2	10		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	70	Total	O	0	0
			70	70		
5	B	3	Total	O	0	0
			3	3		
5	C	4	Total	O	0	0
			4	4		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: DNA (5'-D(*GP*GP*TP*AP*TP*AP*C)-3')

Chain B: 

G301
G302
T303
A304
T305
A306
C307

- Molecule 2: DNA (5'-D(*GP*GP*TP*AP*TP*AP*CP*C)-3')

Chain C: 

G309
G310
T311
A312
T313
A314
C315
C316

- Molecule 3: PROTEIN (DEOXYRIBONUCLEASE I (DNASE I) (E.C.3.1.21.1))

Chain A: 

L1
K2
F6
N7
I8
R9
T10
F11
T14
S17
N18
A19
T20
L21
V26
R30
R31
Y32
D33
R41
D42
S43
H44
L45
V48
L52
N56
Q57
D58
D59
P60
Y63
H64
Y65
E69
R73
N74
S75
T76
K77
E78
R79
L83
F84
R85
P86
N87

K88
Y89
S90
Y91
L92
L92
Y97
ASP
ASP
GLY
GLY
GLU
SER
CYS
GLY
ASN
ASP
S108
F109
S110
R111
E112
K117
F118
S119
K124
V125
S138
D139
A140
I144
Y148
Y151
L152
W158
H159
L160
D168
F169
N170
A171
D172
C173
S174
R185
L186
R187
Q193
T207

V216
L220
L221
Q222
S223
S224
G228
S229
A230
F233
Q236
Y239
G240
L241
M245
E252
Y253
P254
V255
E256
V257
T258
L259
T260

4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section will therefore be incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	51.10 Å 108.40 Å 62.10 Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	6.00 – 2.30	Depositor
% Data completeness (in resolution range)	(Not available) (6.00-2.30)	Depositor
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	TNT, PROLSQ	Depositor
R, R_{free}	0.188 , (Not available)	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	2390	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	B	2.21	7/159 (4.4%)	3.33	29/244 (11.9%)
2	C	2.54	13/180 (7.2%)	3.71	51/276 (18.5%)
3	A	3.83	1/2027 (0.0%)	1.84	31/2760 (1.1%)
All	All	3.66	21/2366 (0.9%)	2.20	111/3280 (3.4%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
4	A	1	0

The worst 5 of 21 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	260	THR	C-OXT	166.98	4.40	1.23
2	C	310	DG	P-O5'	9.03	1.68	1.59
1	B	307	DC	C2'-C1'	8.16	1.60	1.52
2	C	310	DG	C2'-C1'	7.51	1.59	1.52
1	B	301	DG	C3'-C2'	7.28	1.60	1.52

The worst 5 of 111 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	185	ARG	NE-CZ-NH1	17.64	129.12	120.30
3	A	30	ARG	NE-CZ-NH1	17.26	128.93	120.30
3	A	85	ARG	NE-CZ-NH2	15.28	127.94	120.30
3	A	111	ARG	NE-CZ-NH1	14.45	127.52	120.30
3	A	79	ARG	NE-CZ-NH1	13.07	126.83	120.30

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	261	NAG	C2

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	142	0	81	2	0
2	C	161	0	92	5	0
3	A	1982	0	1937	47	1
4	A	28	0	22	0	1
5	A	70	0	0	4	0
5	B	3	0	0	0	0
5	C	4	0	0	0	0
All	All	2390	0	2132	52	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 52 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:125:VAL:HA	3:A:220:LEU:HD13	1.52	0.91
3:A:84:PHE:O	3:A:86:PRO:HD3	1.79	0.81
3:A:252:HIS:HB3	5:A:468:HOH:O	1.81	0.78
3:A:63:TYR:OH	3:A:85:ARG:NH1	2.20	0.73
3:A:10:THR:HG23	3:A:41:ARG:HD3	1.76	0.68

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:228:GLY:CA	4:A:262:NAG:O7[4_455]	2.18	0.02

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	A	246/260 (95%)	229 (93%)	17 (7%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	A	221/229 (96%)	201 (91%)	20 (9%)	12	14

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	A	110	SER
3	A	119	SER
3	A	220	LEU
3	A	89	VAL
3	A	90	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
3	A	87	ASN
3	A	155	GLN
3	A	161	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	A	236	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

2 carbohydrates are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	NAG	A	261	3,4	14,14,15	1.49	2 (14%)	15,19,21	9.78	10 (66%)
4	NAG	A	262	4	14,14,15	2.21	4 (28%)	15,19,21	4.90	11 (73%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NAG	A	261	3,4	1/1/5/7	0/6/23/26	0/1/1/1
4	NAG	A	262	4	-	0/6/23/26	0/1/1/1

The worst 5 of 6 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	262	NAG	C2-N2	-3.06	1.40	1.46

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	261	NAG	C4-C5	-2.13	1.48	1.53
4	A	262	NAG	O5-C1	3.85	1.50	1.43
4	A	262	NAG	C8-C7	4.05	1.58	1.50
4	A	262	NAG	O7-C7	4.28	1.33	1.23

The worst 5 of 21 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	261	NAG	O7-C7-C8	-12.65	98.86	122.06
4	A	262	NAG	O3-C3-C2	-7.00	95.24	109.11
4	A	262	NAG	C8-C7-N2	-6.04	104.55	116.11
4	A	261	NAG	O3-C3-C4	-3.83	101.71	110.34
4	A	261	NAG	C1-O5-C5	-2.89	108.58	112.25

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
4	A	261	NAG	C2

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	262	NAG	0	1

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section will therefore be empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section will therefore be empty.

6.4 Ligands [i](#)

EDS was not executed - this section will therefore be empty.

6.5 Other polymers [i](#)

EDS was not executed - this section will therefore be empty.