



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 03:53 PM GMT

PDB ID : 4DQL
Title : Crystal structure of the FAD binding domain of cytochrome P450 BM3 in complex with NADP+
Authors : joyce, M.G.; leys, D.
Deposited on : 2012-02-16
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	393	<div> <div> <div></div> <div>87%</div> <div>9%</div> <div>.</div> </div> </div>
1	B	393	<div> <div> <div>2%</div> <div>88%</div> <div>8%</div> <div>.</div> <div>.</div> </div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	1PE	A	1104	-	-	-	X
6	PG4	A	1105	-	-	-	X
6	PG4	B	1104	-	-	-	X
6	PG4	B	1105	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 6955 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional P-450/NADPH-P450 reductase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	378	Total	C	N	O	S	0	6	0
			3007	1896	514	586	11			
1	B	379	Total	C	N	O	S	0	8	0
			2998	1895	512	579	12			

- Molecule 2 is FLAVIN-ADENINE DINUCLEOTIDE (three-letter code: FAD) (formula: $C_{27}H_{33}N_9O_{15}P_2$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			53	27	9	15	2		
2	B	1	Total	C	N	O	P	0	0
			53	27	9	15	2		

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			37	14	5	15	3		
3	B	1	Total	C	N	O	P	0	0
			31	10	5	13	3		

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 6 is TETRAETHYLENE GLYCOL (three-letter code: PG4) (formula: $C_8H_{18}O_5$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	C	O	0	0
			13	8	5		
6	B	1	Total	C	O	0	0
			13	8	5		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			13	8	5		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	362	Total	O	0	0
			362	362		
7	B	355	Total	O	0	0
			355	355		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



- Molecule 1: Bifunctional P-450/NADPH-P450 reductase



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	190.66 Å 190.66 Å 74.33 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	19.92 – 2.15 19.92 – 2.15	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.92-2.15) 99.4 (19.92-2.15)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.66 (at 2.15 Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.191 , 0.226 0.186 , 0.220	Depositor DCC
R_{free} test set	4176 reflections (5.25%)	DCC
Wilson B-factor (Å ²)	32.8	Xtriage
Anisotropy	0.039	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 38.4	EDS
Estimated twinning fraction	0.022 for -h,-k,l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	0 of 83698 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6955	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, NAP, PG4, FAD, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.33	14/3068 (0.5%)	0.95	4/4156 (0.1%)
1	B	1.23	7/3060 (0.2%)	1.01	12/4149 (0.3%)
All	All	1.28	21/6128 (0.3%)	0.98	16/8305 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	871	GLU	CG-CD	8.91	1.65	1.51
1	A	749	GLU	CG-CD	8.19	1.64	1.51
1	A	755	GLU	CD-OE1	7.98	1.34	1.25
1	B	826	ARG	CD-NE	-6.89	1.34	1.46
1	B	989	GLU	CG-CD	6.88	1.62	1.51
1	A	817	GLU	CD-OE1	6.85	1.33	1.25
1	A	1021	VAL	CB-CG1	6.60	1.66	1.52
1	B	755	GLU	CG-CD	6.46	1.61	1.51
1	A	820	ALA	CA-CB	6.43	1.66	1.52
1	B	803	GLU	CG-CD	6.38	1.61	1.51
1	A	840	GLU	CG-CD	6.33	1.61	1.51
1	A	982	GLN	CG-CD	6.01	1.64	1.51
1	A	661	GLY	N-CA	5.88	1.54	1.46
1	A	748	GLU	CG-CD	5.80	1.60	1.51
1	B	715	VAL	CB-CG2	5.58	1.64	1.52
1	B	669	ALA	CA-CB	5.58	1.64	1.52

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	841	LYS	CD-CE	5.49	1.65	1.51
1	A	790	TYR	CE1-CZ	-5.27	1.31	1.38
1	A	851	SER	CB-OG	-5.17	1.35	1.42
1	B	789	ALA	CA-CB	5.16	1.63	1.52
1	A	803	GLU	CG-CD	5.10	1.59	1.51

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	826	ARG	NE-CZ-NH2	-13.15	113.72	120.30
1	B	826	ARG	CG-CD-NE	-8.37	94.22	111.80
1	A	766	ARG	NE-CZ-NH2	-7.84	116.38	120.30
1	B	912	ARG	NE-CZ-NH2	-6.87	116.87	120.30
1	B	826	ARG	NE-CZ-NH1	6.82	123.71	120.30
1	B	912	ARG	NE-CZ-NH1	6.80	123.70	120.30
1	A	990	LEU	CB-CG-CD1	6.61	122.24	111.00
1	B	938	ARG	NE-CZ-NH1	6.58	123.59	120.30
1	B	1015	MET	CG-SD-CE	6.13	110.01	100.20
1	B	795	LEU	CB-CG-CD1	5.90	121.03	111.00
1	B	1020	ASP	CB-CG-OD1	5.75	123.48	118.30
1	B	826	ARG	CB-CA-C	-5.67	99.07	110.40
1	B	750	LEU	CA-CB-CG	5.62	128.22	115.30
1	B	839	ASP	CB-CG-OD1	-5.29	113.54	118.30
1	A	771	LYS	CD-CE-NZ	-5.17	99.80	111.70
1	A	961	LEU	CA-CB-CG	5.10	127.02	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	826	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3007	0	2931	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2998	0	2895	14	0
2	A	53	0	31	0	0
2	B	53	0	31	0	0
3	A	37	0	16	0	0
3	B	31	0	11	1	0
4	A	5	0	0	0	0
4	B	5	0	0	1	0
5	A	10	0	12	0	0
6	A	13	0	18	1	0
6	B	26	0	36	2	0
7	A	362	0	0	5	0
7	B	355	0	0	8	1
All	All	6955	0	5981	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (30) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:837:ARG:NE	7:B:1426:HOH:O	2.21	0.73
1:B:797[A]:LYS:O	7:B:1541:HOH:O	2.11	0.68
1:B:797[B]:LYS:O	7:B:1541:HOH:O	2.12	0.67
1:A:973:LYS:HD2	7:A:1556:HOH:O	1.95	0.67
6:B:1105:PG4:H42	7:B:1431:HOH:O	1.97	0.63
1:B:868:TYR:OH	7:B:1505:HOH:O	2.14	0.61
1:A:1037[A]:GLU:HG2	1:A:1042:TYR:HD2	1.66	0.58
1:A:699:ASP:OD1	1:A:884:THR:HG22	2.04	0.58
1:A:993:GLN:NE2	7:A:1373:HOH:O	2.40	0.54
1:B:1011:GLU:HG3	1:B:1033:LEU:CD2	2.42	0.49
1:B:890:THR:HB	7:B:1461:HOH:O	2.11	0.49
1:A:930:GLU:OE1	1:A:986:LYS:CE	2.60	0.49
1:A:1030[A]:ARG:HH11	1:A:1030[A]:ARG:HG3	1.79	0.47
6:B:1105:PG4:C4	7:B:1431:HOH:O	2.59	0.47
1:A:930:GLU:OE1	1:A:986:LYS:HE2	2.16	0.46
1:B:775:PRO:HB2	1:B:776:PRO:HD3	1.98	0.45
1:A:671:LYS:HZ1	6:A:1105:PG4:H42	1.82	0.44
1:B:988:ILE:HD13	1:B:988:ILE:HA	1.72	0.44
1:B:776:PRO:O	1:B:779:VAL:HG22	2.17	0.44
1:A:1037[A]:GLU:CG	1:A:1042:TYR:HD2	2.28	0.43
1:A:1026:GLU:HG2	7:A:1333:HOH:O	2.18	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:672:GLU:OE1	1:A:681:SER:OG	2.29	0.43
1:A:1030[A]:ARG:NH1	1:A:1030[A]:ARG:HG3	2.33	0.43
1:B:877[A]:THR:HG22	7:B:1468:HOH:O	2.19	0.43
1:B:977:GLN:OE1	3:B:1102:NAP:H2A	2.20	0.42
1:B:991:LEU:HD22	1:B:1036:LEU:HD21	2.02	0.42
1:A:973:LYS:HE2	7:A:1555:HOH:O	2.20	0.42
1:B:919:LYS:HB2	1:B:958:ILE:HD13	2.02	0.42
1:A:1028:ASP:OD1	7:A:1473:HOH:O	2.21	0.41
1:B:762:ARG:HD2	4:B:1103:SO4:O4	2.20	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:B:1395:HOH:O	7:B:1533:HOH:O[4_557]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	380/393 (97%)	370 (97%)	10 (3%)	0	100	100
1	B	383/393 (98%)	368 (96%)	15 (4%)	0	100	100
All	All	763/786 (97%)	738 (97%)	25 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	319/332 (96%)	315 (99%)	4 (1%)	76	82
1	B	312/332 (94%)	305 (98%)	7 (2%)	60	63
All	All	631/664 (95%)	620 (98%)	11 (2%)	70	74

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	744	THR
1	A	824	SER
1	A	944	TYR
1	A	990	LEU
1	B	749	GLU
1	B	750	LEU
1	B	795	LEU
1	B	814[A]	LYS
1	B	814[B]	LYS
1	B	944	TYR
1	B	985	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	788	GLN
1	A	924	GLN
1	A	971	GLN
1	A	993	GLN
1	B	707	ASN
1	B	842	GLN
1	B	924	GLN
1	B	1034	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

10 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FAD	A	1101	-	48,58,58	1.64	8 (16%)	54,89,89	2.40	10 (18%)
3	NAP	A	1102	-	31,39,52	1.62	6 (19%)	36,60,80	2.42	8 (22%)
4	SO4	A	1103	-	4,4,4	0.46	0	6,6,6	0.77	0
5	1PE	A	1104	-	9,9,15	0.67	0	8,8,14	0.84	0
6	PG4	A	1105	-	12,12,12	0.45	0	11,11,11	0.66	0
2	FAD	B	1101	-	48,58,58	1.47	6 (12%)	54,89,89	2.25	14 (25%)
3	NAP	B	1102	-	27,33,52	1.62	7 (25%)	34,52,80	2.13	6 (17%)
4	SO4	B	1103	-	4,4,4	0.24	0	6,6,6	0.43	0
6	PG4	B	1104	-	12,12,12	0.54	0	11,11,11	0.36	0
6	PG4	B	1105	-	12,12,12	0.75	0	11,11,11	1.21	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FAD	A	1101	-	-	0/30/50/50	0/6/6/6
3	NAP	A	1102	-	-	0/27/47/67	0/3/3/5
4	SO4	A	1103	-	-	0/0/0/0	0/0/0/0
5	1PE	A	1104	-	-	0/7/7/13	0/0/0/0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	PG4	A	1105	-	-	0/10/10/10	0/0/0/0
2	FAD	B	1101	-	-	0/30/50/50	0/6/6/6
3	NAP	B	1102	-	-	0/17/37/67	0/3/3/5
4	SO4	B	1103	-	-	0/0/0/0	0/0/0/0
6	PG4	B	1104	-	-	0/10/10/10	0/0/0/0
6	PG4	B	1105	-	-	0/10/10/10	0/0/0/0

All (27) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1101	FAD	C6-C5X	-3.09	1.37	1.41
3	A	1102	NAP	O4B-C1B	-2.80	1.37	1.41
3	B	1102	NAP	O4B-C1B	-2.71	1.37	1.41
3	B	1102	NAP	P2B-O3X	-2.65	1.45	1.54
3	A	1102	NAP	PA-O2A	-2.53	1.44	1.54
2	A	1101	FAD	C5A-N7A	-2.37	1.31	1.39
3	A	1102	NAP	C4A-N3A	-2.30	1.32	1.35
3	A	1102	NAP	PA-O1A	-2.10	1.43	1.51
3	B	1102	NAP	PA-O2A	-2.04	1.46	1.54
3	B	1102	NAP	P2B-O2B	2.04	1.66	1.60
3	B	1102	NAP	C2A-N1A	2.18	1.38	1.33
2	B	1101	FAD	O4B-C1B	2.27	1.44	1.41
2	A	1101	FAD	C5X-N5	2.42	1.39	1.35
2	B	1101	FAD	C4-N3	2.71	1.38	1.33
2	A	1101	FAD	C8M-C8	2.74	1.56	1.51
3	B	1102	NAP	C2A-N3A	2.74	1.37	1.32
2	A	1101	FAD	C10-N1	3.10	1.40	1.35
2	B	1101	FAD	C2A-N1A	3.26	1.40	1.33
2	A	1101	FAD	C2A-N3A	3.30	1.38	1.32
3	A	1102	NAP	P2B-O2B	3.33	1.70	1.60
2	A	1101	FAD	C4-N3	3.42	1.39	1.33
3	A	1102	NAP	C5D-C4D	3.67	1.57	1.51
3	B	1102	NAP	PN-O1N	3.98	1.62	1.50
2	B	1101	FAD	C1'-N10	4.47	1.53	1.48
2	B	1101	FAD	C4X-N5	4.63	1.40	1.33
2	A	1101	FAD	C1'-N10	4.75	1.53	1.48
2	A	1101	FAD	C4X-N5	5.47	1.41	1.33

All (40) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	1102	NAP	N3A-C2A-N1A	-9.55	121.58	128.89

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1102	NAP	N3A-C2A-N1A	-9.51	121.61	128.89
2	B	1101	FAD	N3A-C2A-N1A	-8.35	122.50	128.89
2	A	1101	FAD	N3A-C2A-N1A	-8.17	122.64	128.89
2	A	1101	FAD	C4X-C4-N3	-4.31	117.70	123.59
2	A	1101	FAD	C4A-C5A-N7A	-4.16	105.65	109.48
2	A	1101	FAD	P-O3P-PA	-3.65	122.47	132.73
3	A	1102	NAP	O3X-P2B-O1X	-3.19	100.31	110.58
2	B	1101	FAD	C4-C4X-C10	-3.15	117.92	119.94
2	B	1101	FAD	P-O3P-PA	-2.91	124.55	132.73
2	B	1101	FAD	C4X-C4-N3	-2.85	119.69	123.59
3	B	1102	NAP	O3-PN-O1N	-2.53	99.79	107.70
2	B	1101	FAD	C4A-C5A-N7A	-2.47	107.20	109.48
3	A	1102	NAP	PN-O3-PA	-2.43	125.91	132.73
2	A	1101	FAD	C4-C4X-C10	-2.30	118.47	119.94
3	A	1102	NAP	O2B-P2B-O1X	-2.27	101.44	107.11
3	B	1102	NAP	C1B-N9A-C4A	-2.09	123.78	126.94
2	B	1101	FAD	O3B-C3B-C4B	-2.03	104.96	111.05
6	B	1105	PG4	O4-C7-C8	2.07	119.97	110.43
3	B	1102	NAP	O5D-PN-O3	2.07	114.18	104.65
3	A	1102	NAP	O2A-PA-O3	2.09	114.56	105.09
2	B	1101	FAD	C1'-C2'-C3'	2.09	115.80	109.82
2	A	1101	FAD	O2'-C2'-C1'	2.10	115.11	109.94
2	A	1101	FAD	C1'-C2'-C3'	2.15	115.98	109.82
3	B	1102	NAP	C2A-N1A-C6A	2.23	122.76	118.77
2	B	1101	FAD	O2'-C2'-C1'	2.30	115.59	109.94
2	B	1101	FAD	C4X-N5-C5X	2.35	119.47	116.76
2	B	1101	FAD	O3P-P-O5'	2.55	109.70	102.94
3	A	1102	NAP	O3-PA-O5B	2.63	109.92	102.94
2	B	1101	FAD	C5X-C9A-N10	2.76	119.72	117.62
6	B	1105	PG4	C7-O4-C6	2.87	125.66	113.31
2	A	1101	FAD	C1'-N10-C9A	2.93	122.15	118.86
2	B	1101	FAD	C1'-N10-C9A	3.69	123.00	118.86
2	B	1101	FAD	C4-C4X-N5	4.01	123.59	118.72
3	A	1102	NAP	O4B-C1B-N9A	4.25	116.99	108.10
3	B	1102	NAP	O4B-C1B-N9A	4.35	117.21	108.10
2	A	1101	FAD	C5X-C9A-N10	5.62	121.89	117.62
3	A	1102	NAP	O3X-P2B-O2X	5.92	129.94	107.38
2	B	1101	FAD	C4-N3-C2	8.48	122.58	115.25
2	A	1101	FAD	C4-N3-C2	9.80	123.72	115.25

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

4 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	A	1105	PG4	1	0
3	B	1102	NAP	1	0
4	B	1103	SO4	1	0
6	B	1105	PG4	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	378/393 (96%)	-0.29	5 (1%) 79 84	16, 27, 47, 64	0
1	B	379/393 (96%)	-0.24	9 (2%) 62 71	16, 28, 50, 67	0
All	All	757/786 (96%)	-0.26	14 (1%) 71 79	16, 28, 48, 67	0

All (14) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	742	ALA	7.5
1	B	1027	ALA	4.8
1	B	925	GLY	4.6
1	B	658	LYS	4.0
1	B	970	ASN	3.2
1	A	743	LYS	3.0
1	B	1026	GLU	3.0
1	B	972	PRO	2.8
1	A	730	GLU	2.5
1	A	756	LEU	2.4
1	B	971	GLN	2.3
1	B	1019	ALA	2.2
1	A	679	ALA	2.0
1	B	659	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
6	PG4	A	1105	13/13	0.86	0.16	4.93	52,59,71,73	0
6	PG4	B	1105	13/13	0.74	0.25	3.92	57,60,66,67	0
6	PG4	B	1104	13/13	0.90	0.13	3.88	38,46,57,59	0
5	1PE	A	1104	10/16	0.91	0.16	2.06	32,39,47,51	0
4	SO4	A	1103	5/5	0.96	0.11	0.23	57,59,60,62	0
2	FAD	B	1101	53/53	0.98	0.06	-0.77	12,17,21,24	0
3	NAP	A	1102	37/48	0.97	0.08	-0.96	19,25,52,54	0
2	FAD	A	1101	53/53	0.98	0.06	-1.04	16,21,27,33	0
3	NAP	B	1102	31/48	0.98	0.07	-1.14	21,26,53,53	0
4	SO4	B	1103	5/5	0.91	0.29	-	70,71,75,77	0

6.5 Other polymers [i](#)

There are no such residues in this entry.