wwPDB NMR Structure Validation Summary Report (i)

Apr 26, 2016 – 04:13 PM BST

PDB ID	:	1DT7
Title	:	SOLUTION STRUCTURE OF THE C-TERMINAL NEGATIVE REGULATORY DOMAIN OF P53 IN A COMPLEX WITH CA2+-BOUND S100B(BB)
Authors	:	Rustandi, R.R.; Baldisseri, D.M.; Weber, D.J.
Deposited on	:	2000-01-11

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
http://wwpdb.org/validation/2016/NMRValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (i)) were used in the production of this report:

Cyrange	:	Kirchner and Güntert (2011)
NmrClust	:	Kelley et al. (1996)
MolProbity	:	4.02b-467
Mogul	:	unknown
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	rb-20027457
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	rb-20027457

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length	Quality of chain
1	A	92	65% 32% .
1	B	92	65% 32% .
2	X	22	23% 18% 5% 55%
2	Y	22	18% 27% 55%

2 Ensemble composition and analysis (i)

This entry contains 40 models. The atoms present in the NMR models are not consistent. Some calculations may have failed as a result. All residues are included in the validation scores. The authors have identified model 22 as representative, based on the following criterion: *lowest energy*. No medoid model was calculated.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	**Residue range (total)**	**Backbone RMSD (Å)**	**Medoid model**
1	A:0-A:88, B:0-B:88, X:104-X:113, Y:104-Y:113 (198)	Not calculated	Not calculated

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

NmrClust was unable to cluster the ensemble.

Error message: Inconsistent models in file

3 Entry composition (i)

There are 3 unique types of molecules in this entry. The entry contains 3684 atoms, of which 1828 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called S100 CALCIUM-BINDING PROTEIN.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	92	1462	470	714	118	153	7	0
1	B	92	1462	470	714	118	153	7	0

- Molecule 2 is a protein called CELLULAR TUMOR ANTIGEN P53.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
2	X	22	378	111	200	36	30	1	0
2	Y	22	378	111	200	36	30	1	0

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	
			Total	Ca
3	B	2	2	2
3	A	2	2	2

4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: S100 CALCIUM-BINDING PROTEIN

Chain A:



- Molecule 1: S100 CALCIUM-BINDING PROTEIN

Chain B:



- Molecule 2: CELLULAR TUMOR ANTIGEN P53

Chain X:



- Molecule 2: CELLULAR TUMOR ANTIGEN P53

Chain Y:

4.2 Residue scores for the representative (author defined) model from the NMR ensemble

The representative model is number 22. Colouring as in section 4.1 above.

- Molecule 1: S100 CALCIUM-BINDING PROTEIN

Chain A:



- Molecule 1: S100 CALCIUM-BINDING PROTEIN

Chain B:



- Molecule 2: CELLULAR TUMOR ANTIGEN P53

Chain X:

- Molecule 2: CELLULAR TUMOR ANTIGEN P53

Chain Y:

5 Refinement protocol and experimental data overview (i)

Of the 40 calculated structures, 40 were deposited, based on the following criterion: *structure with the lowest energy and the least restraint violations.*

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.851
X-PLOR	refinement	3.851

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	1.14±0.00	0±0/734 (0.0±0.0%)	0.86±0.01	0±0/982 (0.0±0.0%)
1	B	1.14±0.00	0±0/734 (0.0±0.0%)	0.86±0.01	0±0/982 (0.0±0.0%)
2	X	1.06±0.02	0±0/90 (0.0±0.0%)	0.86±0.02	0±0/115 (0.0±0.0%)
2	Y	1.30±1.51	0±0/90 (0.0±0.2%)	0.96±0.63	0±0/115 (0.0±0.1%)
All	All	1.20	1/65921 (0.0%)	0.88	1/87762 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	Y	0.0±0.2	0.0±0.0
All	All	1	0

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	Y	113	GLU	CA-C	101.72	4.17	1.52	29	1

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(^{o})	Ideal(^{o})	Models	
								Worst	Total
2	Y	113	GLU	CB-CA-C	-52.35	5.70	110.40	29	1

All unique chiral outliers are listed below.

Mol	Chain	Res	Type	Atoms	Models (Total)
2	Y	113	GLU	CA	1

There are no planarity outliers.

6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	722	694	690	26±6
1	B	722	694	690	25±6
2	X	89	102	100	7±2
2	Y	89	102	100	7±2
All	All	65041	63680	63184	2454

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

5 of 733 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:B:39:GLU:O	1:B:40:LEU:HD23	0.79	1.78	23	8
1:A:39:GLU:O	1:A:40:LEU:HD23	0.78	1.78	23	8
2:Y:108:LEU:HD11	2:Y:111:LYS:NZ	0.73	1.97	26	2
2:X:108:LEU:HD11	2:X:111:LYS:NZ	0.73	1.98	26	2
2:X:108:LEU:HD11	2:X:111:LYS:HZ1	0.72	1.44	26	3

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	88/92 (96%)	82±1 (94±2%)	5±1 (5±2%)	1±1 (1±1%)	23 69

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	B	88/92 (96%)	82±1 (94±1%)	5±1 (5±2%)	1±1 (1±1%)	23 69
2	X	9/22 (41%)	9±0 (97±5%)	0±0 (3±5%)	0±0 (0±0%)	100 100
2	Y	9/22 (41%)	9±0 (97±5%)	0±0 (3±5%)	0±0 (0±0%)	100 100
All	All	7760/9120 (85%)	7287 (94%)	395 (5%)	78 (1%)	24 71

5 of 13 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	29	LYS	13
1	B	22	GLY	12
1	A	22	GLY	12
1	B	29	LYS	12
1	B	45	GLU	7

6.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	81/84 (96%)	75±2 (93±2%)	6±2 (7±2%)	22 67
1	B	81/84 (96%)	75±2 (92±2%)	6±2 (8±2%)	21 66
2	X	10/21 (48%)	8±1 (81±10%)	2±1 (19±10%)	5 37
2	Y	10/21 (48%)	8±1 (81±10%)	2±1 (19±10%)	5 38
All	All	7280/8400 (87%)	6641 (91%)	639 (9%)	17 62

5 of 94 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	13	VAL	38
1	B	13	VAL	37
1	B	77	VAL	29
1	A	77	VAL	27
2	X	109	MET	23

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no carbohydrates in this entry.

6.6 Ligand geometry (i)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

7 Chemical shift validation (i)

No chemical shift data were provided