



Full wwPDB X-ray Structure Validation Report i

Jan 31, 2016 – 07:03 PM GMT

PDB ID : 1DUZ
Title : HUMAN CLASS I HISTOCOMPATIBILITY ANTIGEN (HLA-A 0201) IN COMPLEX WITH A NONAMERIC PEPTIDE FROM HTLV-1 TAX PROTEIN
Authors : Khan, A.R.; Baker, B.M.; Ghosh, P.; Biddison, W.E.; Wiley, D.C.
Deposited on : 2000-01-19
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	102246	5383 (1.80-1.80)
Ramachandran outliers	100387	5320 (1.80-1.80)
Sidechain outliers	100360	5319 (1.80-1.80)
RSRZ outliers	91569	4547 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 6930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA-A*0201.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			

- Molecule 2 is a protein called BETA-2 MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

2	E	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	ALA	CONFLICT	UNP P61769
E	0	MET	ALA	CONFLICT	UNP P61769

- Molecule 3 is a protein called HTLV-1 OCTAMERIC TAX PEPTIDE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	9	Total	C	N	O		0	0	0
			77	56	9	12				

3	F	9	Total	C	N	O		0	0	0
			77	56	9	12				

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	203	Total	O	0	0
			203	203		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	86	Total O 86 86	0	0
4	C	5	Total O 5 5	0	0
4	D	207	Total O 207 207	0	0
4	E	100	Total O 100 100	0	0
4	F	7	Total O 7 7	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA-A*0201




- Molecule 1: HLA-A*0201



- Molecule 2: BETA-2 MICROGLOBULIN



- Molecule 2: BETA-2 MICROGLOBULIN




- Molecule 3: HTLV-1 OCTAMERIC TAX PEPTIDE

Chain C:  67% 33%



- Molecule 3: HTLV-1 OCTAMERIC TAX PEPTIDE

Chain F:  11% 89% 11%



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.56 Å 63.79 Å 75.08 Å 81.58° 75.66° 77.38°	Depositor
Resolution (Å)	50.00 – 1.80 28.93 – 1.80	Depositor EDS
% Data completeness (in resolution range)	85.3 (50.00-1.80) 83.7 (28.93-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	3.49 (at 1.80 Å)	Xtriage
Refinement program	CNS 0.9	Depositor
R , R_{free}	0.198 , 0.250 0.207 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	DCC
Wilson B-factor (Å ²)	20.6	Xtriage
Anisotropy	0.307	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 43.8	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 72184 reflections	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6930	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.41% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.48	21/2312 (0.9%)	1.29	19/3137 (0.6%)
1	D	1.47	18/2312 (0.8%)	1.35	20/3137 (0.6%)
2	B	1.60	8/860 (0.9%)	1.21	3/1162 (0.3%)
2	E	1.60	9/860 (1.0%)	1.28	4/1162 (0.3%)
3	C	1.45	1/80 (1.2%)	1.10	0/108
3	F	1.54	0/80	1.05	0/108
All	All	1.51	57/6504 (0.9%)	1.29	46/8814 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (57) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	26	TYR	CD1-CE1	7.27	1.50	1.39
2	E	9	VAL	CB-CG2	7.01	1.67	1.52
1	A	159	TYR	CD1-CE1	6.90	1.49	1.39
1	A	147	TRP	CB-CG	6.89	1.62	1.50
1	D	1	GLY	N-CA	6.70	1.56	1.46
2	B	37	VAL	CB-CG1	6.35	1.66	1.52
2	B	44	GLU	CG-CD	6.31	1.61	1.51
1	D	241	PHE	CD2-CE2	6.16	1.51	1.39
1	A	173	GLU	CG-CD	6.07	1.61	1.51
3	C	8	TYR	CD1-CE1	6.04	1.48	1.39
1	D	190	THR	CB-CG2	-6.03	1.32	1.52
1	D	261	VAL	CB-CG1	5.98	1.65	1.52
1	D	171	TYR	CD1-CE1	5.86	1.48	1.39
2	E	56	PHE	CG-CD1	5.86	1.47	1.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	264	GLU	CB-CG	-5.73	1.41	1.52
1	A	244	TRP	CE3-CZ3	5.72	1.48	1.38
1	A	58	GLU	CD-OE1	5.69	1.31	1.25
2	E	30	PHE	CB-CG	5.68	1.61	1.51
1	A	12	VAL	CB-CG2	5.65	1.64	1.52
1	D	209	TYR	CE1-CZ	5.57	1.45	1.38
2	E	55	SER	CB-OG	5.57	1.49	1.42
2	B	70	PHE	CD1-CE1	5.56	1.50	1.39
2	B	74	GLU	CG-CD	5.56	1.60	1.51
1	D	159	TYR	CE1-CZ	5.50	1.45	1.38
2	E	47	GLU	CB-CG	-5.47	1.41	1.52
1	D	211	ALA	CA-CB	-5.46	1.41	1.52
1	A	209	TYR	CE1-CZ	5.46	1.45	1.38
1	D	159	TYR	CD1-CE1	5.44	1.47	1.39
2	B	22	PHE	CD1-CE1	5.42	1.50	1.39
1	D	171	TYR	CD2-CE2	5.41	1.47	1.39
1	A	173	GLU	CB-CG	5.41	1.62	1.52
1	A	7	TYR	CD2-CE2	5.40	1.47	1.39
2	E	48	LYS	C-O	5.38	1.33	1.23
1	D	204	TRP	CB-CG	-5.37	1.40	1.50
1	A	7	TYR	CD1-CE1	5.37	1.47	1.39
1	D	36	PHE	CD2-CE2	5.37	1.50	1.39
1	A	273	ARG	CG-CD	5.36	1.65	1.51
2	E	58	LYS	CD-CE	5.36	1.64	1.51
1	A	99	TYR	CD1-CE1	5.36	1.47	1.39
1	D	208	PHE	CD2-CE2	5.35	1.50	1.39
2	E	26	TYR	CD1-CE1	5.34	1.47	1.39
1	A	5	MET	CG-SD	5.33	1.95	1.81
1	D	173	GLU	CD-OE1	5.32	1.31	1.25
1	A	204	TRP	CE3-CZ3	5.29	1.47	1.38
1	A	223	ASP	CB-CG	5.27	1.62	1.51
1	A	241	PHE	CD1-CE1	5.23	1.49	1.39
1	A	109	PHE	CD2-CE2	5.22	1.49	1.39
1	A	99	TYR	CE2-CZ	5.20	1.45	1.38
1	D	217	TRP	CZ3-CH2	5.18	1.48	1.40
2	B	69	GLU	CG-CD	5.18	1.59	1.51
2	E	66	TYR	CE1-CZ	5.18	1.45	1.38
1	D	22	PHE	CD1-CE1	5.16	1.49	1.39
1	A	71	SER	CB-OG	-5.15	1.35	1.42
1	D	207	SER	CB-OG	-5.08	1.35	1.42
1	A	58	GLU	CG-CD	-5.06	1.44	1.51
1	D	147	TRP	CG-CD1	5.02	1.43	1.36

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	62	PHE	CD2-CE2	5.02	1.49	1.39

All (46) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	D	169	ARG	NE-CZ-NH1	13.62	127.11	120.30
1	D	157	ARG	NE-CZ-NH2	-12.84	113.88	120.30
1	D	169	ARG	NE-CZ-NH2	-11.97	114.31	120.30
1	A	5	MET	CG-SD-CE	-11.53	81.76	100.20
2	E	45	ARG	NE-CZ-NH1	-10.95	114.82	120.30
1	D	108	ARG	NE-CZ-NH2	-9.67	115.46	120.30
1	D	108	ARG	NE-CZ-NH1	9.62	125.11	120.30
2	E	48	LYS	CD-CE-NZ	-9.27	90.39	111.70
1	D	30	ASP	CB-CG-OD1	-8.74	110.43	118.30
1	A	61	ASP	CB-CG-OD1	8.67	126.10	118.30
1	A	111	ARG	NE-CZ-NH1	8.41	124.50	120.30
1	A	220	ASP	CB-CG-OD1	8.34	125.80	118.30
1	A	108	ARG	NE-CZ-NH1	-8.29	116.16	120.30
1	D	111	ARG	NE-CZ-NH2	-8.29	116.16	120.30
1	D	157	ARG	CG-CD-NE	-8.15	94.68	111.80
1	A	14	ARG	NE-CZ-NH1	-7.81	116.40	120.30
2	B	97	ARG	NE-CZ-NH2	-7.78	116.41	120.30
1	A	5	MET	CA-CB-CG	7.71	126.40	113.30
2	E	45	ARG	NE-CZ-NH2	7.65	124.13	120.30
1	D	111	ARG	NE-CZ-NH1	7.58	124.09	120.30
1	D	144	LYS	CD-CE-NZ	7.38	128.67	111.70
1	A	256	ARG	NE-CZ-NH1	6.99	123.80	120.30
1	A	256	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	A	273	ARG	NE-CZ-NH1	-6.86	116.87	120.30
2	E	34	ASP	CB-CG-OD2	-6.64	112.32	118.30
1	D	39	ASP	CB-CG-OD1	6.51	124.16	118.30
1	D	219	ARG	NE-CZ-NH1	6.45	123.52	120.30
1	D	256	ARG	NE-CZ-NH1	6.20	123.40	120.30
1	D	273	ARG	NE-CZ-NH1	5.95	123.28	120.30
2	B	98	ASP	N-CA-C	5.95	127.05	111.00
1	D	44	ARG	NE-CZ-NH2	-5.86	117.37	120.30
1	A	223	ASP	CB-CG-OD2	5.70	123.43	118.30
1	A	61	ASP	CB-CG-OD2	-5.62	113.24	118.30
1	A	129	ASP	CB-CG-OD1	5.61	123.35	118.30
1	D	65	ARG	NE-CZ-NH2	-5.52	117.54	120.30
1	A	273	ARG	NE-CZ-NH2	5.49	123.04	120.30
1	A	28	VAL	CG1-CB-CG2	-5.45	102.17	110.90

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	65	ARG	NE-CZ-NH1	5.43	123.01	120.30
1	D	29	ASP	CB-CG-OD2	5.42	123.18	118.30
1	A	181	ARG	NE-CZ-NH1	5.37	122.99	120.30
1	D	227	ASP	CB-CG-OD1	5.29	123.06	118.30
1	A	14	ARG	NE-CZ-NH2	5.24	122.92	120.30
1	D	273	ARG	NE-CZ-NH2	-5.21	117.70	120.30
2	B	23	LEU	CB-CG-CD2	-5.14	102.27	111.00
1	D	27	TYR	CB-CA-C	-5.07	100.25	110.40
1	A	179	LEU	N-CA-C	5.03	124.57	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	123	TYR	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2096	30	0
1	D	2247	0	2096	37	0
2	B	837	0	803	15	0
2	E	837	0	803	6	0
3	C	77	0	79	1	0
3	F	77	0	79	2	0
4	A	203	0	0	7	0
4	B	86	0	0	4	0
4	C	5	0	0	0	0
4	D	207	0	0	6	0
4	E	100	0	0	1	0
4	F	7	0	0	0	0
All	All	6930	0	5956	85	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (85) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:223:ASP:HB2	4:A:729:HOH:O	1.45	1.16
1:D:97:ARG:HH21	1:D:114:HIS:HE1	1.18	0.92
2:B:96:ASP:O	2:B:98:ASP:N	2.12	0.82
1:D:190:THR:HG21	2:E:98:ASP:OD2	1.80	0.81
1:A:5:MET:HG3	1:A:168:LEU:HB2	1.62	0.79
1:A:263:HIS:CD2	1:A:265:GLY:H	2.03	0.77
1:D:97:ARG:HH21	1:D:114:HIS:CE1	2.04	0.76
1:A:263:HIS:HD2	1:A:265:GLY:H	1.31	0.75
1:D:263:HIS:CD2	1:D:265:GLY:H	2.06	0.73
1:D:263:HIS:HD2	1:D:265:GLY:H	1.37	0.72
1:D:21:ARG:HG2	4:D:789:HOH:O	1.90	0.72
1:D:232:GLU:OE2	2:E:6:LYS:NZ	2.22	0.71
2:B:89:GLN:HG3	4:B:578:HOH:O	1.92	0.70
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.28	0.68
1:D:182:THR:HB	4:D:861:HOH:O	1.94	0.68
1:A:111:ARG:HD2	1:A:113:TYR:OH	1.94	0.67
1:D:19:GLU:OE1	1:D:75:ARG:HD2	1.95	0.67
1:A:97:ARG:HH21	1:A:114:HIS:HE1	1.42	0.67
1:A:93:HIS:HD2	1:A:119:ASP:OD2	1.80	0.64
1:D:19:GLU:OE1	1:D:75:ARG:NH1	2.32	0.63
1:A:82:ARG:HD2	1:A:88:SER:O	1.99	0.62
2:B:85:VAL:HG13	4:B:314:HOH:O	2.00	0.62
1:D:82:ARG:HB3	1:D:82:ARG:HH11	1.65	0.61
1:D:155:GLN:NE2	3:F:5:TYR:CD2	2.68	0.61
1:D:97:ARG:NH2	1:D:114:HIS:HE1	1.95	0.60
1:A:165:VAL:HG12	1:A:169:ARG:NH1	2.17	0.60
1:D:93:HIS:HE1	4:D:594:HOH:O	1.84	0.60
1:D:155:GLN:NE2	3:F:5:TYR:CG	2.70	0.59
1:A:165:VAL:CG1	1:A:169:ARG:NH1	2.67	0.57
2:B:96:ASP:C	2:B:98:ASP:H	2.07	0.57
1:A:273:ARG:NH2	4:A:784:HOH:O	2.27	0.56
1:A:145:HIS:HD2	4:A:645:HOH:O	1.88	0.56
1:D:111:ARG:HD2	1:D:113:TYR:OH	2.04	0.56
1:A:97:ARG:HH21	1:A:114:HIS:CE1	2.22	0.56
1:A:222:GLU:HA	1:A:222:GLU:OE1	2.08	0.54
2:B:96:ASP:C	2:B:98:ASP:N	2.61	0.54
1:D:82:ARG:HH11	1:D:82:ARG:CG	2.21	0.53
1:A:5:MET:HE3	1:A:5:MET:CA	2.26	0.53
3:C:3:PHE:CZ	3:C:5:TYR:HB2	2.44	0.53
2:B:36:GLU:HB3	2:B:83:ASN:HB2	1.90	0.53
1:A:131:ARG:HD2	1:A:157:ARG:HH12	1.73	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:68:LYS:O	1:D:71:SER:HB3	2.09	0.52
2:E:25:CYS:HB2	2:E:39:LEU:HD21	1.93	0.51
1:A:93:HIS:HE1	4:A:453:HOH:O	1.94	0.50
1:A:97:ARG:NH2	1:A:114:HIS:HE1	2.08	0.50
2:E:85:VAL:HG13	4:E:302:HOH:O	2.12	0.49
1:A:262:GLN:NE2	4:A:432:HOH:O	2.23	0.49
1:A:65:ARG:NH2	4:A:742:HOH:O	2.45	0.49
2:B:85:VAL:HG13	4:B:669:HOH:O	2.12	0.49
1:A:182:THR:CG2	1:A:265:GLY:HA2	2.42	0.49
1:D:82:ARG:HH11	1:D:82:ARG:CB	2.26	0.48
1:D:194:VAL:O	1:D:194:VAL:CG1	2.61	0.48
2:B:85:VAL:CG1	4:B:669:HOH:O	2.62	0.47
1:A:97:ARG:HE	1:A:114:HIS:CE1	2.32	0.47
2:B:22:PHE:CE2	2:B:69:GLU:HG3	2.49	0.46
1:D:165:VAL:O	1:D:169:ARG:HG3	2.16	0.46
1:D:267:PRO:HG2	1:D:268:LYS:H	1.81	0.45
1:D:197:HIS:HA	1:D:251:SER:OG	2.17	0.45
1:A:155:GLN:HG2	4:A:904:HOH:O	2.17	0.45
1:A:197:HIS:HA	1:A:251:SER:OG	2.17	0.45
1:D:89:GLU:HG3	1:D:89:GLU:O	2.16	0.45
1:D:173:GLU:O	1:D:176:LYS:HB2	2.17	0.44
1:A:5:MET:HB3	1:A:5:MET:HE2	1.32	0.44
2:B:96:ASP:O	2:B:97:ARG:C	2.55	0.44
1:A:210:PRO:O	1:A:263:HIS:HE1	2.01	0.43
2:B:44:GLU:HG2	2:B:45:ARG:N	2.33	0.43
2:E:7:ILE:HG12	2:E:27:VAL:HG12	2.01	0.42
1:D:82:ARG:NH1	1:D:82:ARG:CG	2.83	0.42
1:D:253:GLN:NE2	1:D:256:ARG:HH11	2.17	0.42
1:D:207:SER:HB3	4:D:617:HOH:O	2.18	0.42
1:A:253:GLN:NE2	1:A:256:ARG:HH11	2.18	0.42
1:D:82:ARG:NH1	1:D:82:ARG:HG2	2.35	0.42
1:D:13:SER:HA	1:D:20:PRO:HB3	2.01	0.42
1:D:66:LYS:HE2	4:D:638:HOH:O	2.19	0.41
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.55	0.41
1:D:190:THR:HG22	4:D:370:HOH:O	2.21	0.41
1:D:78:LEU:HD23	1:D:95:VAL:HG23	2.02	0.41
1:A:182:THR:HG21	1:A:265:GLY:HA2	2.02	0.41
1:D:266:LEU:HA	1:D:267:PRO:HD2	1.69	0.41
2:B:83:ASN:ND2	2:B:90:PRO:HB3	2.36	0.41
2:B:33:SER:HB3	2:B:62:PHE:CE2	2.56	0.41
1:D:82:ARG:HH11	1:D:82:ARG:HG2	1.86	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:HIS:HD2	1:D:119:ASP:OD2	2.04	0.40
1:A:182:THR:HG23	1:A:265:GLY:HA2	2.03	0.40
2:B:38:ASP:OD1	2:B:45:ARG:HD2	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	273/275 (99%)	265 (97%)	7 (3%)	1 (0%)	39 23
1	D	273/275 (99%)	266 (97%)	7 (3%)	0	100 100
2	B	98/100 (98%)	95 (97%)	2 (2%)	1 (1%)	19 5
2	E	98/100 (98%)	96 (98%)	2 (2%)	0	100 100
3	C	7/9 (78%)	7 (100%)	0	0	100 100
3	F	7/9 (78%)	7 (100%)	0	0	100 100
All	All	756/768 (98%)	736 (97%)	18 (2%)	2 (0%)	46 29

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	97	ARG
1	A	17	ARG

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	231/231 (100%)	223 (96%)	8 (4%)	43 25
1	D	231/231 (100%)	225 (97%)	6 (3%)	54 37
2	B	95/95 (100%)	92 (97%)	3 (3%)	46 29
2	E	95/95 (100%)	91 (96%)	4 (4%)	36 18
3	C	8/8 (100%)	8 (100%)	0	100 100
3	F	8/8 (100%)	8 (100%)	0	100 100
All	All	668/668 (100%)	647 (97%)	21 (3%)	47 30

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	MET
1	A	15	PRO
1	A	35	ARG
1	A	58	GLU
1	A	67	VAL
1	A	182	THR
1	A	223	ASP
1	A	268	LYS
2	B	9	VAL
2	B	34	ASP
2	B	70	PHE
1	D	35	ARG
1	D	67	VAL
1	D	82	ARG
1	D	110	LEU
1	D	182	THR
1	D	227	ASP
2	E	34	ASP
2	E	70	PHE
2	E	75	LYS
2	E	91	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	93	HIS
1	A	114	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	145	HIS
1	A	174	ASN
1	A	253	GLN
1	A	263	HIS
2	B	2	GLN
2	B	89	GLN
1	D	93	HIS
1	D	114	HIS
1	D	145	HIS
1	D	174	ASN
1	D	253	GLN
1	D	263	HIS

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

There are no ligands in this entry.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	-0.16	6 (2%) 65 60	13, 22, 36, 46	0
1	D	275/275 (100%)	-0.11	6 (2%) 65 60	13, 22, 37, 45	0
2	B	100/100 (100%)	-0.07	2 (2%) 68 64	14, 22, 34, 42	0
2	E	100/100 (100%)	-0.39	0 100 100	13, 19, 29, 34	0
3	C	9/9 (100%)	0.10	0 100 100	17, 21, 30, 35	0
3	F	9/9 (100%)	0.09	1 (11%) 7 5	18, 25, 30, 36	0
All	All	768/768 (100%)	-0.16	15 (1%) 68 64	13, 22, 36, 46	0

All (15) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	VAL	4.0
1	D	194	VAL	4.0
1	D	251	SER	3.4
1	D	196	ASP	3.1
1	D	16	GLY	2.9
1	A	196	ASP	2.8
1	D	195	SER	2.7
3	F	5	TYR	2.5
1	D	18	GLY	2.5
1	A	223	ASP	2.4
2	B	48	LYS	2.3
1	A	192	HIS	2.3
1	A	195	SER	2.2
1	A	16	GLY	2.1
2	B	99	MET	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

There are no ligands in this entry.

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.