



# wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 08:15 AM GMT

PDB ID : 3DXB  
Title : Structure of the UHM domain of Puf60 fused to thioredoxin  
Authors : Corsini, L.; Hothorn, M.; Scheffzek, K.; Stier, G.; Sattler, M.  
Deposited on : 2008-07-24  
Resolution : 2.20 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3774 (2.20-2.20)
Clashscore	102246	4477 (2.20-2.20)
Ramachandran outliers	100387	4404 (2.20-2.20)
Sidechain outliers	100360	4405 (2.20-2.20)
RSRZ outliers	91569	3781 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	222	<div> <div>3%</div> <div>85%</div> <div>10%</div> <div>• •</div> </div>
1	B	222	<div> <div>3%</div> <div>81%</div> <div>11%</div> <div>• 5%</div> </div>
1	C	222	<div> <div>%</div> <div>81%</div> <div>13%</div> <div>• 5%</div> </div>
1	D	222	<div> <div>5%</div> <div>81%</div> <div>13%</div> <div>• •</div> </div>
1	E	222	<div> <div>3%</div> <div>84%</div> <div>9%</div> <div>• 5%</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	222	 82% 13% 5%
1	G	222	 85% 9% 5%
1	H	222	 83% 11% 5%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	E	557	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 14337 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called thioredoxin N-terminally fused to Puf60(UHM).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1655	1047	270	330	8			
1	B	211	Total	C	N	O	S	0	1	0
			1624	1030	267	320	7			
1	C	211	Total	C	N	O	S	0	4	0
			1657	1046	270	334	7			
1	D	216	Total	C	N	O	S	0	2	0
			1684	1064	276	336	8			
1	E	210	Total	C	N	O	S	0	0	0
			1622	1027	265	323	7			
1	F	211	Total	C	N	O	S	0	1	0
			1636	1035	267	327	7			
1	G	211	Total	C	N	O	S	0	0	0
			1630	1031	266	326	7			
1	H	211	Total	C	N	O	S	0	0	0
			1630	1031	266	326	7			

There are 104 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	MET	-	EXPRESSION TAG	UNP P0AA27
A	336	LYS	-	EXPRESSION TAG	UNP P0AA27
A	337	HIS	-	EXPRESSION TAG	UNP P0AA27
A	338	HIS	-	EXPRESSION TAG	UNP P0AA27
A	339	HIS	-	EXPRESSION TAG	UNP P0AA27
A	340	HIS	-	EXPRESSION TAG	UNP P0AA27
A	341	HIS	-	EXPRESSION TAG	UNP P0AA27
A	342	HIS	-	EXPRESSION TAG	UNP P0AA27
A	343	PRO	-	EXPRESSION TAG	UNP P0AA27
A	453	GLY	-	LINKER	UNP Q9UHX1
A	454	SER	-	LINKER	UNP Q9UHX1
A	455	ALA	-	LINKER	UNP Q9UHX1
A	456	MET	-	LINKER	UNP Q9UHX1

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
B	335	MET	-	EXPRESSION TAG	UNP P0AA27
B	336	LYS	-	EXPRESSION TAG	UNP P0AA27
B	337	HIS	-	EXPRESSION TAG	UNP P0AA27
B	338	HIS	-	EXPRESSION TAG	UNP P0AA27
B	339	HIS	-	EXPRESSION TAG	UNP P0AA27
B	340	HIS	-	EXPRESSION TAG	UNP P0AA27
B	341	HIS	-	EXPRESSION TAG	UNP P0AA27
B	342	HIS	-	EXPRESSION TAG	UNP P0AA27
B	343	PRO	-	EXPRESSION TAG	UNP P0AA27
B	453	GLY	-	LINKER	UNP Q9UHX1
B	454	SER	-	LINKER	UNP Q9UHX1
B	455	ALA	-	LINKER	UNP Q9UHX1
B	456	MET	-	LINKER	UNP Q9UHX1
C	335	MET	-	EXPRESSION TAG	UNP P0AA27
C	336	LYS	-	EXPRESSION TAG	UNP P0AA27
C	337	HIS	-	EXPRESSION TAG	UNP P0AA27
C	338	HIS	-	EXPRESSION TAG	UNP P0AA27
C	339	HIS	-	EXPRESSION TAG	UNP P0AA27
C	340	HIS	-	EXPRESSION TAG	UNP P0AA27
C	341	HIS	-	EXPRESSION TAG	UNP P0AA27
C	342	HIS	-	EXPRESSION TAG	UNP P0AA27
C	343	PRO	-	EXPRESSION TAG	UNP P0AA27
C	453	GLY	-	LINKER	UNP Q9UHX1
C	454	SER	-	LINKER	UNP Q9UHX1
C	455	ALA	-	LINKER	UNP Q9UHX1
C	456	MET	-	LINKER	UNP Q9UHX1
D	335	MET	-	EXPRESSION TAG	UNP P0AA27
D	336	LYS	-	EXPRESSION TAG	UNP P0AA27
D	337	HIS	-	EXPRESSION TAG	UNP P0AA27
D	338	HIS	-	EXPRESSION TAG	UNP P0AA27
D	339	HIS	-	EXPRESSION TAG	UNP P0AA27
D	340	HIS	-	EXPRESSION TAG	UNP P0AA27
D	341	HIS	-	EXPRESSION TAG	UNP P0AA27
D	342	HIS	-	EXPRESSION TAG	UNP P0AA27
D	343	PRO	-	EXPRESSION TAG	UNP P0AA27
D	453	GLY	-	LINKER	UNP Q9UHX1
D	454	SER	-	LINKER	UNP Q9UHX1
D	455	ALA	-	LINKER	UNP Q9UHX1
D	456	MET	-	LINKER	UNP Q9UHX1
E	335	MET	-	EXPRESSION TAG	UNP P0AA27
E	336	LYS	-	EXPRESSION TAG	UNP P0AA27
E	337	HIS	-	EXPRESSION TAG	UNP P0AA27

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
E	338	HIS	-	EXPRESSION TAG	UNP P0AA27
E	339	HIS	-	EXPRESSION TAG	UNP P0AA27
E	340	HIS	-	EXPRESSION TAG	UNP P0AA27
E	341	HIS	-	EXPRESSION TAG	UNP P0AA27
E	342	HIS	-	EXPRESSION TAG	UNP P0AA27
E	343	PRO	-	EXPRESSION TAG	UNP P0AA27
E	453	GLY	-	LINKER	UNP Q9UHX1
E	454	SER	-	LINKER	UNP Q9UHX1
E	455	ALA	-	LINKER	UNP Q9UHX1
E	456	MET	-	LINKER	UNP Q9UHX1
F	335	MET	-	EXPRESSION TAG	UNP P0AA27
F	336	LYS	-	EXPRESSION TAG	UNP P0AA27
F	337	HIS	-	EXPRESSION TAG	UNP P0AA27
F	338	HIS	-	EXPRESSION TAG	UNP P0AA27
F	339	HIS	-	EXPRESSION TAG	UNP P0AA27
F	340	HIS	-	EXPRESSION TAG	UNP P0AA27
F	341	HIS	-	EXPRESSION TAG	UNP P0AA27
F	342	HIS	-	EXPRESSION TAG	UNP P0AA27
F	343	PRO	-	EXPRESSION TAG	UNP P0AA27
F	453	GLY	-	LINKER	UNP Q9UHX1
F	454	SER	-	LINKER	UNP Q9UHX1
F	455	ALA	-	LINKER	UNP Q9UHX1
F	456	MET	-	LINKER	UNP Q9UHX1
G	335	MET	-	EXPRESSION TAG	UNP P0AA27
G	336	LYS	-	EXPRESSION TAG	UNP P0AA27
G	337	HIS	-	EXPRESSION TAG	UNP P0AA27
G	338	HIS	-	EXPRESSION TAG	UNP P0AA27
G	339	HIS	-	EXPRESSION TAG	UNP P0AA27
G	340	HIS	-	EXPRESSION TAG	UNP P0AA27
G	341	HIS	-	EXPRESSION TAG	UNP P0AA27
G	342	HIS	-	EXPRESSION TAG	UNP P0AA27
G	343	PRO	-	EXPRESSION TAG	UNP P0AA27
G	453	GLY	-	LINKER	UNP Q9UHX1
G	454	SER	-	LINKER	UNP Q9UHX1
G	455	ALA	-	LINKER	UNP Q9UHX1
G	456	MET	-	LINKER	UNP Q9UHX1
H	335	MET	-	EXPRESSION TAG	UNP P0AA27
H	336	LYS	-	EXPRESSION TAG	UNP P0AA27
H	337	HIS	-	EXPRESSION TAG	UNP P0AA27
H	338	HIS	-	EXPRESSION TAG	UNP P0AA27
H	339	HIS	-	EXPRESSION TAG	UNP P0AA27
H	340	HIS	-	EXPRESSION TAG	UNP P0AA27

*Continued on next page...*


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
H	341	HIS	-	EXPRESSION TAG	UNP P0AA27
H	342	HIS	-	EXPRESSION TAG	UNP P0AA27
H	343	PRO	-	EXPRESSION TAG	UNP P0AA27
H	453	GLY	-	LINKER	UNP Q9UHX1
H	454	SER	-	LINKER	UNP Q9UHX1
H	455	ALA	-	LINKER	UNP Q9UHX1
H	456	MET	-	LINKER	UNP Q9UHX1

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	H	1	Total Cl 1 1	0	0
2	A	1	Total Cl 1 1	0	0
2	F	1	Total Cl 1 1	0	0
2	E	1	Total Cl 1 1	0	0

- Molecule 3 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	E	1	Total C O 4 2 2	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	148	Total 148	O 148	0	0
4	B	132	Total 132	O 132	0	0
4	C	173	Total 173	O 173	0	0
4	D	123	Total 123	O 123	0	0
4	E	149	Total 149	O 149	0	0
4	F	156	Total 156	O 156	0	0
4	G	141	Total 141	O 141	0	0
4	H	169	Total 169	O 169	0	0




### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $\text{RSRZ} > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)




- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)




- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)




- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)




- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)




- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)



- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)



- Molecule 1: thioredoxin N-terminally fused to Puf60(UHM)



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	75.12Å 89.43Å 299.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.75 – 2.20 46.82 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.8 (49.75-2.20) 99.8 (46.82-2.20)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	0.11	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.39 (at 2.20Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.211 , 0.271 0.232 , 0.276	Depositor DCC
$R_{free}$ test set	5174 reflections (5.03%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.5	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 50.2	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.28$	Xtriage
Outliers	3 of 102913 reflections (0.003%)	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	14337	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.23 % of the origin peak, indicating pseudo translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo translational symmetry is equal to 8.8621e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1682	0.50	0/2272
1	B	0.32	0/1650	0.49	0/2231
1	C	0.33	0/1683	0.51	0/2275
1	D	0.33	0/1712	0.52	0/2314
1	E	0.32	0/1648	0.50	0/2228
1	F	0.34	0/1662	0.51	0/2247
1	G	0.32	0/1656	0.49	0/2239
1	H	0.33	0/1656	0.50	0/2239
All	All	0.33	0/13349	0.50	0/18045

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	2
1	C	0	2
1	E	0	2
All	All	0	8

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 8 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	343	PRO	Peptide

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Group
1	A	534	PHE	Peptide
1	B	502	GLU	Peptide
1	B	504	GLU	Peptide
1	C	451	LEU	Peptide

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1655	0	1631	14	0
1	B	1624	0	1594	33	0
1	C	1657	0	1615	25	0
1	D	1684	0	1645	28	0
1	E	1622	0	1594	33	0
1	F	1636	0	1601	17	0
1	G	1630	0	1598	19	0
1	H	1630	0	1598	19	0
2	A	1	0	0	1	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	H	1	0	0	0	0
3	E	4	0	6	0	0
4	A	148	0	0	3	0
4	B	132	0	0	5	0
4	C	173	0	0	7	0
4	D	123	0	0	3	0
4	E	149	0	0	3	0
4	F	156	0	0	3	0
4	G	141	0	0	1	0
4	H	169	0	0	2	0
All	All	14337	0	12882	176	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 176 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:346:ASP:N	1:G:347:LYS:HB2	1.36	1.41
1:C:452[B]:ALA:CB	1:C:453[B]:GLY:HA2	1.55	1.34
1:C:452[B]:ALA:HB1	1:C:453[B]:GLY:CA	1.64	1.27
1:E:504:GLU:N	1:E:505:ASP:HB3	1.52	1.24
1:B:502:GLU:HG2	1:D:427:ASN:O	1.40	1.18

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	212/222 (96%)	209 (99%)	3 (1%)	0	100	100
1	B	210/222 (95%)	202 (96%)	5 (2%)	3 (1%)	14	10
1	C	213/222 (96%)	206 (97%)	7 (3%)	0	100	100
1	D	216/222 (97%)	208 (96%)	7 (3%)	1 (0%)	34	35
1	E	208/222 (94%)	203 (98%)	4 (2%)	1 (0%)	34	35
1	F	210/222 (95%)	208 (99%)	2 (1%)	0	100	100
1	G	209/222 (94%)	205 (98%)	3 (1%)	1 (0%)	34	35
1	H	209/222 (94%)	204 (98%)	4 (2%)	1 (0%)	34	35
All	All	1687/1776 (95%)	1645 (98%)	35 (2%)	7 (0%)	39	42

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	502	GLU
1	B	504	GLU
1	B	506	ALA
1	H	504	GLU
1	E	503	GLU

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	177/185 (96%)	171 (97%)	6 (3%)	44	54
1	B	170/185 (92%)	160 (94%)	10 (6%)	24	27
1	C	175/185 (95%)	168 (96%)	7 (4%)	38	47
1	D	179/185 (97%)	162 (90%)	17 (10%)	11	10
1	E	172/185 (93%)	167 (97%)	5 (3%)	50	62
1	F	173/185 (94%)	164 (95%)	9 (5%)	29	33
1	G	173/185 (94%)	168 (97%)	5 (3%)	50	62
1	H	173/185 (94%)	166 (96%)	7 (4%)	38	47
All	All	1392/1480 (94%)	1326 (95%)	66 (5%)	34	39

5 of 66 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	447	LEU
1	D	490	VAL
1	H	447	LEU
1	D	468	ASP
1	D	483[A]	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 19 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	546	GLN
1	D	523	HIS
1	F	546	GLN
1	C	350	HIS
1	G	500	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
3	EDO	E	557	-	3,3,3	0.52	0	2,2,2	0.28	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	E	557	-	-	0/1/1/1	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	214/222 (96%)	-0.05	6 (2%) 56 55	12, 18, 25, 31	0
1	B	211/222 (95%)	0.09	6 (2%) 56 55	12, 23, 30, 32	0
1	C	211/222 (95%)	-0.09	2 (0%) 85 85	12, 18, 24, 27	0
1	D	216/222 (97%)	0.17	10 (4%) 36 35	15, 22, 28, 32	0
1	E	210/222 (94%)	0.10	7 (3%) 50 49	12, 20, 29, 32	0
1	F	211/222 (95%)	0.07	2 (0%) 85 85	10, 19, 28, 36	0
1	G	211/222 (95%)	0.01	1 (0%) 91 91	12, 22, 28, 32	0
1	H	211/222 (95%)	0.01	1 (0%) 91 91	11, 19, 27, 32	0
All	All	1695/1776 (95%)	0.04	35 (2%) 67 65	10, 20, 28, 36	0

The worst 5 of 35 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	341	HIS	10.1
1	B	346	ASP	5.6
1	F	505	ASP	4.7
1	F	346	ASP	4.7
1	D	534	PHE	3.4

### 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	EDO	E	557	4/4	0.72	0.26	19.27	32,33,33,33	0
2	CL	E	1	1/1	0.88	0.10	-	46,46,46,46	0
2	CL	F	1	1/1	0.81	0.12	-	45,45,45,45	0
2	CL	H	1	1/1	0.96	0.11	-	42,42,42,42	0
2	CL	A	1	1/1	0.94	0.08	-	37,37,37,37	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.