



Full wwPDB X-ray Structure Validation Report i

Feb 1, 2016 – 08:19 AM GMT

PDB ID : 3E5X
Title : OCPA complexed CprK
Authors : Levy, C.
Deposited on : 2008-08-14
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix)	:	1.9-1692
EDS	:	rb-20026688
Percentile statistics	:	20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac	:	5.8.0135
CCP4	:	6.5.0
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	91344	6249 (2.00-2.00)
Clashscore	102246	7340 (2.00-2.00)
Ramachandran outliers	100387	7248 (2.00-2.00)
Sidechain outliers	100360	7247 (2.00-2.00)
RSRZ outliers	91569	6262 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-

ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	3C4	B	505	-	-	-	X

2 Entry composition [\(i\)](#)


There are 3 unique types of molecules in this entry. The entry contains 7584 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cyclic nucleotide-binding protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	214	Total	C	N	O	S	0	0	0
			1719	1108	283	317	11			
1	C	214	Total	C	N	O	S	0	0	0
			1719	1108	283	317	11			
1	B	219	Total	C	N	O	S	0	0	0
			1758	1133	291	323	11			
1	D	220	Total	C	N	O	S	0	1	0
			1775	1143	294	327	11			

- Molecule 2 is (3-CHLORO-4-HYDROXYPHENYL)ACETIC ACID (three-letter code: 3C4) (formula: C₈H₇ClO₃).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	Cl	O	0	0
			12	8	1	3		
2	C	1	Total	C	Cl	O	0	0
			12	8	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total C Cl O 12 8 1 3	0	0
2	B	1	Total C Cl O 12 8 1 3	0	0
2	D	1	Total C Cl O 12 8 1 3	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	121	Total O 121 121	0	0
3	C	130	Total O 130 130	0	0
3	B	151	Total O 151 151	0	0
3	D	151	Total O 151 151	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cyclic nucleotide-binding protein



- Molecule 1: Cyclic nucleotide-binding protein




- Molecule 1: Cyclic nucleotide-binding protein



TYR	
SER	
ASP	
PRO	
ASN	
SER	
SER	
VAL	
GLY	
LEU	
GLY	
LYS	
D9	

- Molecule 1: Cyclic nucleotide-binding protein



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.70 Å 118.38 Å 87.46 Å 90.00° 95.68° 90.00°	Depositor
Resolution (Å)	35.94 – 2.00 32.92 – 2.00	Depositor EDS
% Data completeness (in resolution range)	96.3 (35.94-2.00) 96.3 (32.92-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ ¹	3.41 (at 2.00 Å)	Xtriage
Refinement program	REFMAC 5.2	Depositor
R , R_{free}	0.193 , 0.238 0.196 , 0.239	Depositor DCC
R_{free} test set	3754 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 36.7	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$< L > = 0.51$, $< L^2 > = 0.35$	Xtriage
Outliers	0 of 74673 reflections	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7584	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
3C4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	1.12	4/1752 (0.2%)	0.91	7/2362 (0.3%)
1	B	1.16	7/1792 (0.4%)	0.97	5/2415 (0.2%)
1	C	1.13	5/1752 (0.3%)	0.99	3/2362 (0.1%)
1	D	1.14	7/1813 (0.4%)	0.93	5/2444 (0.2%)
All	All	1.14	23/7109 (0.3%)	0.95	20/9583 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	C	0	1
All	All	0	2

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	171	THR	CB-OG1	17.68	1.78	1.43
1	C	200	CYS	CB-SG	-11.46	1.62	1.82
1	B	200	CYS	CB-SG	-10.38	1.64	1.82
1	B	221	GLU	C-N	10.38	1.57	1.34
1	A	171	THR	CB-CG2	10.35	1.86	1.52
1	C	68	GLU	CG-CD	8.32	1.64	1.51
1	D	200	CYS	CB-SG	-8.16	1.68	1.82
1	D	23	GLU	CG-CD	7.99	1.64	1.51
1	D	49	GLU	CG-CD	7.59	1.63	1.51
1	A	68	GLU	CG-CD	7.35	1.62	1.51
1	B	135	ALA	CA-CB	6.89	1.67	1.52
1	A	49	GLU	CG-CD	6.28	1.61	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	49	GLU	CD-OE1	6.15	1.32	1.25
1	D	58	GLU	CG-CD	5.86	1.60	1.51
1	B	68	GLU	CG-CD	5.75	1.60	1.51
1	C	68	GLU	CB-CG	5.62	1.62	1.52
1	B	23	GLU	CG-CD	5.58	1.60	1.51
1	D	111	SER	CB-OG	-5.47	1.35	1.42
1	B	49	GLU	CG-CD	5.21	1.59	1.51
1	B	57	VAL	CB-CG2	5.13	1.63	1.52
1	D	135	ALA	CA-CB	5.12	1.63	1.52
1	C	135	ALA	CA-CB	5.09	1.63	1.52
1	C	195	SER	CB-OG	5.01	1.48	1.42

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	139	ARG	NE-CZ-NH2	-8.93	115.84	120.30
1	C	139	ARG	NE-CZ-NH1	8.81	124.70	120.30
1	A	139	ARG	NE-CZ-NH1	8.12	124.36	120.30
1	D	113	ARG	NE-CZ-NH1	7.73	124.17	120.30
1	A	56	LEU	CB-CG-CD1	-7.18	98.80	111.00
1	B	26	ARG	NE-CZ-NH2	-7.11	116.74	120.30
1	D	207	LEU	CA-CB-CG	6.62	130.52	115.30
1	A	139	ARG	NE-CZ-NH2	-6.56	117.02	120.30
1	C	196	ARG	NE-CZ-NH1	6.52	123.56	120.30
1	D	56	LEU	CB-CG-CD2	-6.10	100.64	111.00
1	A	127	PHE	CB-CG-CD1	6.08	125.06	120.80
1	B	10	PHE	N-CA-C	-5.62	95.81	111.00
1	D	139	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	196	ARG	NE-CZ-NH2	-5.40	117.60	120.30
1	B	113	ARG	NE-CZ-NH1	5.31	122.95	120.30
1	B	64	ASP	CB-CG-OD2	-5.20	113.62	118.30
1	A	121	ASP	CB-CG-OD1	5.18	122.96	118.30
1	A	113	ARG	NE-CZ-NH1	5.07	122.84	120.30
1	D	139	ARG	NE-CZ-NH2	-5.06	117.77	120.30
1	A	155	ARG	NE-CZ-NH1	5.06	122.83	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	9	ASP	Peptide
1	C	9	ASP	Peptide

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1719	0	1748	13	0
1	B	1758	0	1788	12	0
1	C	1719	0	1748	14	0
1	D	1775	0	1798	23	0
2	A	12	0	6	0	0
2	B	24	0	12	0	0
2	C	12	0	6	0	0
2	D	12	0	6	0	0
3	A	121	0	0	2	0
3	B	151	0	0	3	0
3	C	130	0	0	4	0
3	D	151	0	0	6	0
All	All	7584	0	7112	61	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (61) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:171:THR:CG2	1:A:171:THR:CB	1.86	1.50
1:A:171:THR:CB	1:A:171:THR:OG1	1.78	1.30
1:B:121:ASP:OD2	3:B:1594:HOH:O	1.90	0.89
1:D:23:GLU:O	1:D:26:ARG:HG2	1.78	0.82
1:C:151:ILE:HD12	3:C:1099:HOH:O	1.80	0.80
1:B:139:ARG:NH1	3:B:1318:HOH:O	2.21	0.72
1:D:152:ARG:NH1	1:D:187:THR:O	2.33	0.61
1:D:157:PHE:CZ	1:D:198:LEU:HD21	2.38	0.58
1:D:150:THR:HG21	3:D:1116:HOH:O	2.04	0.57
1:D:49:GLU:HG2	3:D:1305:HOH:O	2.06	0.56
1:B:56:LEU:HD12	3:B:1575:HOH:O	2.06	0.56
1:D:174:ILE:HG22	1:D:176:MET:HG2	1.89	0.54
1:C:9:ASP:N	1:C:9:ASP:OD1	2.36	0.54
1:A:173:GLU:HG2	1:A:215:ILE:HD12	1.89	0.53
1:D:204:GLU:HB2	1:D:206:ILE:HD12	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:209:LYS:O	1:D:213:LYS:O	2.26	0.53
1:A:151:ILE:HD12	3:A:1258:HOH:O	2.09	0.53
1:D:157:PHE:HZ	1:D:198:LEU:HD21	1.74	0.53
1:C:68:GLU:CD	1:C:68:GLU:H	2.13	0.52
1:D:154:LEU:CD2	1:D:201:LEU:HD21	2.40	0.52
1:C:58:GLU:HG2	1:C:103:ARG:HB3	1.92	0.51
1:B:19:PHE:O	1:B:20:PHE:C	2.49	0.51
1:D:46:PRO:HB3	1:D:66:ILE:HD12	1.93	0.51
1:C:182:SER:HA	1:C:185:GLU:HG2	1.93	0.51
1:A:171:THR:CG2	1:A:171:THR:CA	2.84	0.50
1:D:180:GLN:HG2	1:D:198:LEU:HD12	1.94	0.48
1:D:228:GLN:HB2	3:D:1583:HOH:O	2.13	0.47
1:C:191:HIS:HD2	3:C:1133:HOH:O	1.96	0.47
1:D:228:GLN:C	3:D:1130:HOH:O	2.53	0.47
1:D:200:CYS:O	1:D:204:GLU:HG3	2.15	0.46
1:D:206:ILE:HD13	1:D:225:LEU:CD1	2.46	0.45
1:A:209:LYS:HG2	1:A:214:ILE:CD1	2.46	0.45
1:A:15:ILE:HD11	1:A:156:LEU:HD11	1.98	0.45
1:B:26:ARG:NH2	1:B:57:VAL:O	2.50	0.44
1:B:31:MET:O	1:B:111:SER:HB2	2.16	0.44
1:C:181:LYS:O	1:C:185:GLU:HG2	2.17	0.44
1:B:150:THR:HG22	1:B:197:VAL:HG21	1.99	0.44
1:C:46:PRO:HB3	1:C:66:ILE:HD12	2.01	0.43
1:C:182:SER:OG	3:C:1534:HOH:O	2.21	0.43
1:B:149:PRO:HB3	1:B:189:VAL:HG22	2.00	0.43
1:D:139:ARG:CZ	3:D:1502:HOH:O	2.67	0.43
1:C:31:MET:O	1:C:111:SER:HB2	2.19	0.42
1:D:151:ILE:HD13	1:D:151:ILE:HA	1.86	0.42
1:B:64:ASP:OD2	1:B:72:GLU:OE2	2.37	0.42
1:C:64:ASP:OD2	1:C:72:GLU:OE2	2.37	0.42
1:D:53:MET:HB3	1:D:83:LEU:HD11	2.01	0.42
1:B:15:ILE:HD12	1:B:17:ASP:HB3	2.00	0.42
1:A:95:TYR:CZ	1:A:97:THR:CG2	3.02	0.42
1:B:204:GLU:HB2	1:B:206:ILE:HD12	2.01	0.42
1:D:150:THR:HG23	3:D:1208:HOH:O	2.19	0.41
1:C:192:VAL:HG12	3:C:1343:HOH:O	2.20	0.41
1:D:26:ARG:HA	1:D:29:THR:HG23	2.03	0.41
1:A:140:GLN:HG3	3:A:1263:HOH:O	2.21	0.41
1:B:54:ILE:HB	1:B:84:ILE:HB	2.03	0.41
1:A:10:PHE:CE2	1:A:12:GLY:HA3	2.56	0.41
1:A:118:THR:O	1:C:166:LYS:HB3	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:221:GLU:HA	1:C:221:GLU:OE1	2.20	0.41
1:A:209:LYS:HE2	1:A:209:LYS:O	2.21	0.41
1:D:54:ILE:HB	1:D:84:ILE:HB	2.03	0.40
1:D:64:ASP:OD2	1:D:72:GLU:OE2	2.38	0.40
1:A:160:LEU:HD12	1:A:160:LEU:HA	1.95	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	212/250 (85%)	206 (97%)	5 (2%)	1 (0%)	34 26
1	B	217/250 (87%)	211 (97%)	6 (3%)	0	100 100
1	C	212/250 (85%)	204 (96%)	6 (3%)	2 (1%)	21 13
1	D	219/250 (88%)	214 (98%)	4 (2%)	1 (0%)	34 26
All	All	860/1000 (86%)	835 (97%)	21 (2%)	4 (0%)	34 26

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	210	LYS
1	C	10	PHE
1	C	20	PHE
1	A	20	PHE

5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	189/220 (86%)	183 (97%)	6 (3%)	46 44
1	B	193/220 (88%)	190 (98%)	3 (2%)	70 73
1	C	189/220 (86%)	183 (97%)	6 (3%)	46 44
1	D	195/220 (89%)	189 (97%)	6 (3%)	47 46
All	All	766/880 (87%)	745 (97%)	21 (3%)	52 52

All (21) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	143	GLU
1	A	160	LEU
1	A	175	THR
1	A	176	MET
1	A	207	LEU
1	A	209	LYS
1	C	9	ASP
1	C	10	PHE
1	C	56	LEU
1	C	181	LYS
1	C	200	CYS
1	C	209	LYS
1	B	25	LEU
1	B	30	GLN
1	B	209	LYS
1	D	21	PRO
1	D	25	LEU
1	D	49	GLU
1	D	56	LEU
1	D	181	LYS
1	D	207	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [\(i\)](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	3C4	A	501	-	9,12,12	5.13	4 (44%)	13,16,16	2.50	6 (46%)
2	3C4	B	504	-	9,12,12	4.65	4 (44%)	13,16,16	2.35	3 (23%)
2	3C4	B	505	-	9,12,12	4.55	6 (66%)	13,16,16	3.39	5 (38%)
2	3C4	C	502	-	9,12,12	4.73	3 (33%)	13,16,16	2.23	2 (15%)
2	3C4	D	503	-	9,12,12	5.17	4 (44%)	13,16,16	2.39	3 (23%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	3C4	A	501	-	-	0/2/4/4	0/1/1/1
2	3C4	B	504	-	-	0/2/4/4	0/1/1/1
2	3C4	B	505	-	-	0/2/4/4	0/1/1/1
2	3C4	C	502	-	-	0/2/4/4	0/1/1/1
2	3C4	D	503	-	-	0/2/4/4	0/1/1/1

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	505	3C4	CE2-CD2	-7.42	1.25	1.38
2	D	503	3C4	CE2-CD2	-6.64	1.26	1.38
2	A	501	3C4	CE2-CD2	-6.40	1.27	1.38
2	B	504	3C4	CE2-CD2	-5.14	1.29	1.38
2	C	502	3C4	CE2-CD2	-4.64	1.30	1.38
2	B	505	3C4	CD1-CA	-4.21	1.31	1.39
2	D	503	3C4	CD1-CA	-4.10	1.32	1.39
2	B	505	3C4	CD1-CE1	-3.09	1.33	1.38
2	B	504	3C4	CD1-CA	-2.94	1.34	1.39
2	B	505	3C4	OH-CZ	2.91	1.42	1.36
2	B	505	3C4	CE2-CZ	3.37	1.45	1.39
2	A	501	3C4	CE2-CZ	3.60	1.46	1.39
2	D	503	3C4	CE2-CZ	3.63	1.46	1.39
2	A	501	3C4	OH-CZ	4.17	1.45	1.36
2	B	504	3C4	CE2-CZ	4.76	1.48	1.39
2	C	502	3C4	CE2-CZ	6.08	1.50	1.39
2	B	505	3C4	CZ-CE1	8.97	1.48	1.39
2	B	504	3C4	CZ-CE1	11.41	1.50	1.39
2	C	502	3C4	CZ-CE1	11.45	1.50	1.39
2	D	503	3C4	CZ-CE1	12.55	1.51	1.39
2	A	501	3C4	CZ-CE1	12.60	1.51	1.39

All (19) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	505	3C4	CE2-CZ-CE1	-9.07	109.37	118.57
2	D	503	3C4	CE2-CZ-CE1	-7.05	111.42	118.57
2	B	504	3C4	CE2-CZ-CE1	-6.77	111.70	118.57
2	C	502	3C4	CE2-CZ-CE1	-6.63	111.85	118.57
2	A	501	3C4	CE2-CZ-CE1	-6.61	111.87	118.57
2	A	501	3C4	CE1-CD1-CA	-2.59	118.69	120.36
2	B	505	3C4	CE1-CD1-CA	-2.12	118.99	120.36
2	D	503	3C4	CD2-CA-CD1	2.12	121.68	118.55
2	B	504	3C4	OH-CZ-CE2	2.16	125.29	119.35
2	A	501	3C4	CD2-CA-CD1	2.28	121.92	118.55
2	B	504	3C4	CD2-CA-CD1	2.33	121.99	118.55
2	A	501	3C4	CD1-CE1-CL	2.34	122.10	118.50
2	D	503	3C4	OH-CZ-CE2	2.39	125.92	119.35
2	A	501	3C4	CD2-CE2-CZ	2.67	123.24	120.49
2	C	502	3C4	OH-CZ-CE2	2.73	126.85	119.35
2	A	501	3C4	OH-CZ-CE2	2.90	127.32	119.35
2	B	505	3C4	CE2-CD2-CA	2.91	125.02	121.04
2	B	505	3C4	OH-CZ-CE2	3.67	129.43	119.35

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	505	3C4	CD1-CE1-CZ	5.47	124.27	121.22

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

6 Fit of model and data i

6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	214/250 (85%)	2.02	85 (39%) 0 1	20, 29, 62, 69	0
1	B	219/250 (87%)	1.20	45 (20%) 1 1	18, 27, 44, 49	0
1	C	214/250 (85%)	1.27	43 (20%) 1 2	22, 29, 44, 50	0
1	D	220/250 (88%)	1.17	41 (18%) 2 2	19, 27, 45, 57	0
All	All	867/1000 (86%)	1.41	214 (24%) 1 1	18, 28, 55, 69	0

All (214) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	199	ALA	10.9
1	A	212	ASN	8.5
1	A	134	VAL	7.4
1	B	134	VAL	6.8
1	A	167	ARG	6.7
1	D	134	VAL	6.6
1	A	175	THR	6.6
1	A	217	TYR	6.5
1	A	219	LEU	6.3
1	B	141	VAL	6.3
1	A	192	VAL	6.2
1	A	207	LEU	6.2
1	A	211	LYS	5.9
1	B	168	VAL	5.8
1	A	191	HIS	5.7
1	A	171	THR	5.6
1	C	75	LEU	5.6
1	A	190	HIS	5.6
1	B	137	TYR	5.6
1	D	212	ASN	5.6
1	D	170	ASP	5.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	206	ILE	5.3
1	A	137	TYR	5.3
1	C	134	VAL	5.3
1	A	75	LEU	5.2
1	A	138	ALA	5.2
1	A	215	ILE	5.1
1	D	137	TYR	5.1
1	A	209	LYS	5.1
1	B	212	ASN	5.0
1	A	172	TYR	5.0
1	A	205	ASN	5.0
1	A	166	LYS	4.9
1	A	198	LEU	4.9
1	A	141	VAL	4.8
1	A	170	ASP	4.8
1	D	211	LYS	4.7
1	A	173	GLU	4.7
1	B	167	ARG	4.6
1	A	126	ILE	4.6
1	D	138	ALA	4.6
1	C	212	ASN	4.5
1	D	136	TYR	4.5
1	A	136	TYR	4.5
1	A	214	ILE	4.4
1	A	174	ILE	4.4
1	C	137	TYR	4.3
1	D	131	LEU	4.3
1	B	131	LEU	4.2
1	C	192	VAL	4.2
1	B	136	TYR	4.1
1	A	210	LYS	4.1
1	B	138	ALA	4.1
1	A	203	ARG	4.0
1	D	135	ALA	4.0
1	A	196	ARG	4.0
1	C	130	TYR	4.0
1	A	201	LEU	4.0
1	A	135	ALA	4.0
1	B	135	ALA	4.0
1	A	168	VAL	4.0
1	C	76	TYR	3.9
1	A	213	LYS	3.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	130	TYR	3.8
1	A	65	ILE	3.8
1	A	204	GLU	3.8
1	D	141	VAL	3.8
1	C	190	HIS	3.7
1	B	75	LEU	3.7
1	B	130	TYR	3.7
1	D	166	LYS	3.7
1	B	186	ILE	3.6
1	A	19	PHE	3.6
1	B	127	PHE	3.6
1	A	165	GLY	3.6
1	D	130	TYR	3.6
1	A	164	GLN	3.6
1	A	63	LEU	3.5
1	C	84	ILE	3.5
1	D	186	ILE	3.5
1	C	131	LEU	3.5
1	A	218	ASN	3.5
1	C	65	ILE	3.5
1	C	211	LYS	3.4
1	A	142	ALA	3.4
1	A	131	LEU	3.4
1	A	221	GLU	3.4
1	A	208	ASP	3.4
1	C	210	LYS	3.4
1	A	194	VAL	3.4
1	A	197	VAL	3.4
1	D	19	PHE	3.3
1	B	74	LEU	3.3
1	B	166	LYS	3.3
1	C	136	TYR	3.3
1	B	172	TYR	3.3
1	C	19	PHE	3.3
1	A	186	ILE	3.3
1	C	191	HIS	3.2
1	C	85	GLY	3.2
1	C	135	ALA	3.2
1	B	211	LYS	3.2
1	B	132	THR	3.2
1	D	142	ALA	3.2
1	A	83	LEU	3.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	133	LYS	3.2
1	C	141	VAL	3.2
1	A	177	PRO	3.2
1	A	158	TYR	3.1
1	B	170	ASP	3.1
1	C	203	ARG	3.1
1	D	209	LYS	3.1
1	C	133	LYS	3.1
1	C	63	LEU	3.1
1	B	84	ILE	3.1
1	D	15	ILE	3.1
1	B	169	GLY	3.1
1	D	83	LEU	3.1
1	B	171	THR	3.0
1	A	76	TYR	3.0
1	C	38	ALA	3.0
1	D	213	LYS	3.0
1	B	76	TYR	3.0
1	C	138	ALA	2.9
1	C	112	LEU	2.9
1	A	157	PHE	2.9
1	B	16	PRO	2.9
1	A	216	VAL	2.9
1	A	127	PHE	2.9
1	D	167	ARG	2.9
1	C	177	PRO	2.8
1	B	19	PHE	2.8
1	C	196	ARG	2.8
1	C	186	ILE	2.8
1	A	195	SER	2.8
1	A	85	GLY	2.8
1	B	140	GLN	2.8
1	B	123	ILE	2.8
1	A	193	THR	2.7
1	A	222	LEU	2.7
1	B	126	ILE	2.7
1	C	101	PRO	2.7
1	D	22	ILE	2.7
1	C	178	LEU	2.7
1	D	219	LEU	2.7
1	D	123	ILE	2.7
1	A	202	LYS	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	63	LEU	2.6
1	B	142	ALA	2.6
1	D	127	PHE	2.6
1	C	74	LEU	2.6
1	A	13	ALA	2.6
1	D	14	ILE	2.6
1	C	132	THR	2.6
1	D	228	GLN	2.6
1	C	14	ILE	2.6
1	D	169	GLY	2.6
1	D	210	LYS	2.6
1	D	75	LEU	2.5
1	D	126	ILE	2.5
1	D	177	PRO	2.5
1	D	172	TYR	2.5
1	B	133	LYS	2.5
1	D	156	LEU	2.5
1	A	22	ILE	2.5
1	A	101	PRO	2.4
1	A	116	PHE	2.4
1	D	168	VAL	2.4
1	A	178	LEU	2.4
1	C	205	ASN	2.4
1	A	180	GLN	2.3
1	C	83	LEU	2.3
1	A	150	THR	2.3
1	A	139	ARG	2.3
1	B	116	PHE	2.3
1	B	25	LEU	2.3
1	A	163	SER	2.3
1	A	179	SER	2.3
1	C	193	THR	2.3
1	A	38	ALA	2.3
1	C	15	ILE	2.2
1	A	140	GLN	2.2
1	B	115	VAL	2.2
1	C	142	ALA	2.2
1	B	15	ILE	2.2
1	B	22	ILE	2.2
1	A	74	LEU	2.2
1	A	200	CYS	2.2
1	C	87	LEU	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	63	LEU	2.2
1	C	53	MET	2.2
1	D	139	ARG	2.2
1	A	112	LEU	2.2
1	B	165	GLY	2.1
1	A	84	ILE	2.1
1	B	66	ILE	2.1
1	D	84	ILE	2.1
1	A	132	THR	2.1
1	B	69	ASP	2.1
1	D	69	ASP	2.1
1	B	161	CYS	2.1
1	C	57	VAL	2.1
1	B	213	LYS	2.0
1	A	129	ASN	2.0
1	A	94	ILE	2.0
1	A	107	PHE	2.0
1	D	225	LEU	2.0
1	B	85	GLY	2.0
1	B	188	GLY	2.0
1	D	129	ASN	2.0
1	C	220	GLY	2.0
1	D	132	THR	2.0
1	B	139	ARG	2.0
1	C	199	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors

of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
2	3C4	B	505	12/12	0.68	0.20	2.91	47,48,49,54	0
2	3C4	B	504	12/12	0.98	0.11	-1.66	17,20,21,22	0
2	3C4	D	503	12/12	0.98	0.10	-1.99	17,18,22,22	0
2	3C4	A	501	12/12	0.98	0.07	-2.81	17,19,21,21	0
2	3C4	C	502	12/12	0.98	0.07	-3.33	16,19,21,22	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.