



wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 27, 2016 – 04:57 PM EDT

PDB ID : 5E6K
Title : Ketosynthase from module 6 of the bacillaene synthase from *Bacillus subtilis* 168 (C167S mutant, crystal form 2)
Authors : Wagner, D.T.; Gay, D.C.; Keatinge-Clay, A.T.
Deposited on : 2015-10-10
Resolution : 2.16 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.1 (RC1), CSD as537be (2016)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20027939
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027939


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.16 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1045 (2.16-2.16)
Clashscore	102246	1152 (2.16-2.16)
Ramachandran outliers	100387	1131 (2.16-2.16)
Sidechain outliers	100360	1131 (2.16-2.16)
RSRZ outliers	91569	1050 (2.16-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	617	<div> <div>4%</div> <div>75%</div> <div>10%</div> <div>15%</div> </div>
1	B	617	<div> <div>7%</div> <div>75%</div> <div>10%</div> <div>14%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	601	-	-	-	X
2	SO4	A	602	-	-	-	X

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 8408 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyketide synthase PksL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4149	2637	698	790	24			
1	B	531	Total	C	N	O	S	0	0	0
			4171	2652	702	793	24			

There are 42 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-21	MET	-	initiating methionine	UNP Q05470
A	-20	GLY	-	expression tag	UNP Q05470
A	-19	SER	-	expression tag	UNP Q05470
A	-18	SER	-	expression tag	UNP Q05470
A	-17	HIS	-	expression tag	UNP Q05470
A	-16	HIS	-	expression tag	UNP Q05470
A	-15	HIS	-	expression tag	UNP Q05470
A	-14	HIS	-	expression tag	UNP Q05470
A	-13	HIS	-	expression tag	UNP Q05470
A	-12	HIS	-	expression tag	UNP Q05470
A	-11	SER	-	expression tag	UNP Q05470
A	-10	SER	-	expression tag	UNP Q05470
A	-9	GLY	-	expression tag	UNP Q05470
A	-8	LEU	-	expression tag	UNP Q05470
A	-7	VAL	-	expression tag	UNP Q05470
A	-6	PRO	-	expression tag	UNP Q05470
A	-5	ARG	-	expression tag	UNP Q05470
A	-4	GLY	-	expression tag	UNP Q05470
A	-3	SER	-	expression tag	UNP Q05470
A	-2	SER	-	expression tag	UNP Q05470
A	169	SER	CYS	engineered mutation	UNP Q05470
B	-21	MET	-	initiating methionine	UNP Q05470
B	-20	GLY	-	expression tag	UNP Q05470
B	-19	SER	-	expression tag	UNP Q05470
B	-18	SER	-	expression tag	UNP Q05470

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-17	HIS	-	expression tag	UNP Q05470
B	-16	HIS	-	expression tag	UNP Q05470
B	-15	HIS	-	expression tag	UNP Q05470
B	-14	HIS	-	expression tag	UNP Q05470
B	-13	HIS	-	expression tag	UNP Q05470
B	-12	HIS	-	expression tag	UNP Q05470
B	-11	SER	-	expression tag	UNP Q05470
B	-10	SER	-	expression tag	UNP Q05470
B	-9	GLY	-	expression tag	UNP Q05470
B	-8	LEU	-	expression tag	UNP Q05470
B	-7	VAL	-	expression tag	UNP Q05470
B	-6	PRO	-	expression tag	UNP Q05470
B	-5	ARG	-	expression tag	UNP Q05470
B	-4	GLY	-	expression tag	UNP Q05470
B	-3	SER	-	expression tag	UNP Q05470
B	-2	SER	-	expression tag	UNP Q05470
B	169	SER	CYS	engineered mutation	UNP Q05470

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	33	Total	O	0	0
			33	33		
3	B	35	Total	O	0	0
			35	35		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	76.61Å 105.45Å 89.33Å 90.00° 96.76° 90.00°	Depositor
Resolution (Å)	36.55 – 2.16 36.55 – 2.16	Depositor EDS
% Data completeness (in resolution range)	99.2 (36.55-2.16) 99.2 (36.55-2.16)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.82 (at 2.16Å)	Xtriage
Refinement program	REFMAC 5.8.0107	Depositor
R, R_{free}	0.221 , 0.254 0.225 , 0.257	Depositor DCC
R_{free} test set	3769 reflections (5.29%)	DCC
Wilson B-factor (Å ²)	35.1	Xtriage
Anisotropy	0.102	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 32.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	8408	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/4237	0.73	7/5724 (0.1%)
1	B	0.45	0/4260	0.71	6/5752 (0.1%)
All	All	0.46	0/8497	0.72	13/11476 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

The worst 5 of 13 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	243	ARG	NE-CZ-NH1	-8.43	116.09	120.30
1	B	584	ALA	CB-CA-C	7.08	120.72	110.10
1	B	488	ARG	NE-CZ-NH2	7.03	123.81	120.30
1	B	488	ARG	NE-CZ-NH1	-6.23	117.19	120.30
1	A	243	ARG	NE-CZ-NH2	5.97	123.29	120.30

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	243	ARG	Sidechain
1	A	368	PRO	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4149	0	4071	53	0
1	B	4171	0	4105	60	0
2	A	15	0	0	1	0
2	B	5	0	0	0	0
3	A	33	0	0	0	0
3	B	35	0	0	1	0
All	All	8408	0	8176	111	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 111 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:63:MET:CE	1:B:65:MET:HG2	1.39	1.48
1:B:209:CYS:SG	1:B:230:PHE:O	2.03	1.17
1:B:63:MET:CE	1:B:65:MET:CG	2.21	1.17
1:B:63:MET:HE2	1:B:65:MET:CG	1.78	1.09
1:A:446:ALA:HA	1:A:485:MET:CE	1.81	1.08

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	513/617 (83%)	499 (97%)	14 (3%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	517/617 (84%)	497 (96%)	20 (4%)	0	100	100
All	All	1030/1234 (84%)	996 (97%)	34 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	439/515 (85%)	436 (99%)	3 (1%)	88	93
1	B	442/515 (86%)	436 (99%)	6 (1%)	74	80
All	All	881/1030 (86%)	872 (99%)	9 (1%)	82	87

5 of 9 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	151	ARG
1	B	591	PRO
1	B	399	ASP
1	A	496	ARG
1	B	397	THR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 23 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	512	ASN
1	B	90	GLN
1	B	389	ASN
1	B	29	HIS
1	B	96	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry ⓘ

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	601	-	4,4,4	0.37	0	6,6,6	0.37	0
2	SO4	A	602	-	4,4,4	0.44	0	6,6,6	0.13	0
2	SO4	A	603	-	4,4,4	0.34	0	6,6,6	0.12	0
2	SO4	B	601	-	4,4,4	0.38	0	6,6,6	0.14	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	601	-	-	0/0/0/0	0/0/0/0
2	SO4	A	602	-	-	0/0/0/0	0/0/0/0
2	SO4	A	603	-	-	0/0/0/0	0/0/0/0
2	SO4	B	601	-	-	0/0/0/0	0/0/0/0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	603	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/617 (85%)	0.08	25 (4%) 35 46	22, 35, 72, 107	0
1	B	531/617 (86%)	0.15	41 (7%) 16 22	23, 36, 79, 118	0
All	All	1058/1234 (85%)	0.11	66 (6%) 24 33	22, 36, 77, 118	0

The worst 5 of 66 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	571	TYR	6.6
1	A	513	GLY	5.7
1	B	60	CYS	5.6
1	A	47	TYR	5.5
1	B	132	LEU	5.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
2	SO4	A	602	5/5	0.90	0.22	3.94	72,80,90,90	0
2	SO4	A	601	5/5	0.95	0.22	3.58	61,63,64,68	0
2	SO4	A	603	5/5	0.92	0.21	-	80,90,93,101	0
2	SO4	B	601	5/5	0.96	0.23	-	64,65,70,71	0

6.5 Other polymers [i](#)

There are no such residues in this entry.