



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 10, 2016 – 05:55 PM EDT

PDB ID : 5E76  
Title : Crystal structure of Bacova\_02651 with xylogluco-oligosaccharide  
Authors : Koropatkin, N.M.  
Deposited on : 2015-10-11  
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.1 (RC1), CSD as537be (2016)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	511	<div> <div>7%</div> <div>85%</div> <div>10%</div> <div>...</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	610	-	-	-	X
4	EDO	A	611	-	-	-	X

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	EDO	A	613	-	-	-	X
4	EDO	A	614	-	-	-	X

## 2 Entry composition [i](#)


There are 7 unique types of molecules in this entry. The entry contains 8216 atoms, of which 3895 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SusD-like protein BACOVA\_02651.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	492	Total	C	H	N	O	S	0	0	0
			7751	2527	3777	658	772	17			

- Molecule 2 is BETA-D-GLUCOSE (three-letter code: BGC) (formula:  $C_6H_{12}O_6$ ).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			22	6	11	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			20	6	9	5		
2	A	1	Total	C	H	O	0	0
			20	6	9	5		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			21	6	10	5		
2	A	1	Total	C	H	O	0	0
			23	6	11	6		

- Molecule 3 is XYLOPYRANOSE (three-letter code: XYS) (formula: C<sub>5</sub>H<sub>10</sub>O<sub>5</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	H	O	0	0
			18	5	9	4		
3	A	1	Total	C	H	O	0	0
			18	5	9	4		

- Molecule 4 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		
4	A	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Cl	0	0
			1	1		

- Molecule 6 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Na	0	0
			1	1		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	229	Total 229	O 229	0	0

### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: SusD-like protein BACOVA\_02651





## 4 Data and refinement statistics

Property	Value	Source
Space group	I 4 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	131.50Å 131.50Å 188.04Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	56.13 – 2.30 56.58 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.6 (56.13-2.30) 96.0 (56.58-2.30)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.37 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.217 , 0.253 0.231 , 0.252	Depositor DCC
$R_{free}$ test set	2004 reflections (5.95%)	DCC
Wilson B-factor (Å <sup>2</sup> )	30.7	Xtriage
Anisotropy	0.793	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 59.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.033 for -1/2*h-1/2*k-1/2*l,-1/2*h-1/2*k+1/2*l,-h+k 0.018 for -1/2*h+1/2*k-1/2*l,1/2*h-1/2*k-1/2*l,-h-k	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	8216	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	43.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, BGC, YYS, EDO, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.69	2/4071 (0.0%)	0.70	1/5529 (0.0%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	518	GLU	CD-OE1	6.11	1.32	1.25
1	A	146	CYS	CB-SG	5.16	1.91	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	78	MET	CG-SD-CE	5.22	108.56	100.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3974	3777	3781	26	0
2	A	78	70	63	2	0
3	A	18	18	16	2	0
4	A	20	30	30	2	0
5	A	1	0	0	0	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	1	0	0	0	0
7	A	229	0	0	3	0
All	All	4321	3895	3890	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:121:GLU:OE1	1:A:121:GLU:N	2.24	0.70
1:A:226:ASN:OD1	7:A:701:HOH:O	2.13	0.67
3:A:604:XYS:H52	2:A:606:BGC:C6	2.30	0.61
1:A:165:GLN:O	1:A:167:LEU:N	2.29	0.58
1:A:72:SER:HB3	1:A:78:MET:CE	2.34	0.58
3:A:604:XYS:H52	2:A:606:BGC:H6C1	1.89	0.53
1:A:295:PHE:HE1	1:A:299:VAL:HG21	1.75	0.51
1:A:72:SER:HB3	1:A:78:MET:HE1	1.94	0.50
1:A:270:GLU:OE1	1:A:420:ARG:NH1	2.42	0.50
1:A:293:LEU:HD13	1:A:360:GLY:HA3	1.94	0.49
1:A:320:GLU:OE1	1:A:418:ARG:NH2	2.42	0.49
1:A:264:LYS:HE3	1:A:368:ASP:HB2	1.94	0.49
1:A:78:MET:HE3	1:A:444:MET:SD	2.53	0.49
1:A:122:THR:O	1:A:123:GLN:CB	2.61	0.48
1:A:355:SER:N	1:A:356:PRO:CD	2.78	0.47
1:A:308:SER:O	7:A:702:HOH:O	2.20	0.47
1:A:461:ASP:HB2	4:A:614:EDO:H11	1.98	0.46
1:A:84:TYR:OH	1:A:97:GLU:OE1	2.28	0.45
1:A:151:MET:HB3	1:A:158:VAL:HG11	1.98	0.45
1:A:460:PRO:HG2	4:A:614:EDO:H21	2.00	0.43
1:A:78:MET:CE	1:A:444:MET:SD	3.07	0.43
1:A:263:TYR:CD1	1:A:379:PRO:HB2	2.53	0.43
1:A:482:ASP:HB2	1:A:486:LYS:O	2.19	0.42
1:A:47:LEU:O	1:A:50:ALA:N	2.52	0.41
1:A:503:PRO:HA	1:A:506:TYR:CD2	2.56	0.41
1:A:81:PRO:CG	1:A:475:ARG:NH1	2.83	0.41
1:A:179:GLU:OE2	7:A:703:HOH:O	2.22	0.40
1:A:74:ARG:NH2	1:A:519:ASP:O	2.54	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	484/511 (95%)	454 (94%)	27 (6%)	3 (1%)	30	36

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	65	ARG
1	A	166	LYS
1	A	202	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	424/439 (97%)	409 (96%)	15 (4%)	43	58

All (15) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	45	GLU
1	A	56	ASN
1	A	78	MET
1	A	100	SER
1	A	104	SER
1	A	121	GLU
1	A	264	LYS
1	A	321	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	424	ASN
1	A	446	TYR
1	A	476	THR
1	A	482	ASP
1	A	493	ASP
1	A	498	THR
1	A	500	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	BGC	A	601	2	11,11,12	2.59	5 (45%)	15,15,17	0.99	1 (6%)
2	BGC	A	602	2	11,11,12	2.34	4 (36%)	15,15,17	1.60	2 (13%)
2	BGC	A	603	2	11,11,12	2.48	5 (45%)	15,15,17	1.16	1 (6%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	XYS	A	604	2	9,9,10	0.22	0	12,12,14	0.57	0
3	XYS	A	605	2	9,9,10	1.64	1 (11%)	12,12,14	1.99	5 (41%)
2	BGC	A	606	3,2	11,11,12	0.26	0	15,15,17	0.59	0
2	BGC	A	607	3,2	11,11,12	2.53	4 (36%)	15,15,17	1.49	3 (20%)
2	BGC	A	608	2	11,11,12	2.59	4 (36%)	15,15,17	1.76	4 (26%)
2	BGC	A	609	2	12,12,12	2.24	3 (25%)	17,17,17	0.93	0
4	EDO	A	610	-	3,3,3	0.45	0	2,2,2	0.67	0
4	EDO	A	611	-	3,3,3	0.41	0	2,2,2	0.78	0
4	EDO	A	612	-	3,3,3	0.51	0	2,2,2	0.47	0
4	EDO	A	613	6	3,3,3	0.30	0	2,2,2	0.74	0
4	EDO	A	614	-	3,3,3	0.65	0	2,2,2	0.44	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	A	601	2	-	0/2/19/22	0/1/1/1
2	BGC	A	602	2	-	0/2/19/22	0/1/1/1
2	BGC	A	603	2	-	0/2/19/22	0/1/1/1
3	XYS	A	604	2	-	0/0/14/17	0/1/1/1
3	XYS	A	605	2	-	0/0/14/17	0/1/1/1
2	BGC	A	606	3,2	-	0/2/19/22	0/1/1/1
2	BGC	A	607	3,2	-	0/2/19/22	0/1/1/1
2	BGC	A	608	2	-	0/2/19/22	0/1/1/1
2	BGC	A	609	2	-	0/2/22/22	0/1/1/1
4	EDO	A	610	-	-	0/1/1/1	0/0/0/0
4	EDO	A	611	-	-	0/1/1/1	0/0/0/0
4	EDO	A	612	-	-	0/1/1/1	0/0/0/0
4	EDO	A	613	6	-	0/1/1/1	0/0/0/0
4	EDO	A	614	-	-	0/1/1/1	0/0/0/0

All (26) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	605	XYS	O5-C1	-3.59	1.35	1.42
2	A	607	BGC	C2-C3	-3.29	1.48	1.52
2	A	603	BGC	C2-C3	-3.23	1.48	1.52
2	A	608	BGC	C2-C3	-3.17	1.48	1.52
2	A	602	BGC	C2-C3	-2.74	1.48	1.52

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	609	BGC	O1-C1	-2.65	1.30	1.39
2	A	607	BGC	C4-C5	-2.64	1.47	1.53
2	A	601	BGC	C2-C3	-2.59	1.49	1.52
2	A	609	BGC	C4-C3	-2.51	1.45	1.52
2	A	603	BGC	C4-C5	-2.42	1.47	1.53
2	A	607	BGC	C4-C3	-2.35	1.46	1.52
2	A	603	BGC	C4-C3	-2.30	1.46	1.52
2	A	601	BGC	C4-C3	-2.24	1.46	1.52
2	A	608	BGC	C4-C3	-2.23	1.46	1.52
2	A	601	BGC	C4-C5	-2.19	1.48	1.53
2	A	608	BGC	O3-C3	2.00	1.47	1.43
2	A	603	BGC	O3-C3	2.03	1.47	1.43
2	A	602	BGC	O4-C4	2.03	1.47	1.43
2	A	602	BGC	O3-C3	2.06	1.47	1.43
2	A	601	BGC	O3-C3	2.13	1.48	1.43
2	A	609	BGC	O5-C1	5.55	1.53	1.43
2	A	602	BGC	O5-C1	5.74	1.53	1.43
2	A	603	BGC	O5-C1	6.01	1.53	1.43
2	A	607	BGC	O5-C1	6.12	1.53	1.43
2	A	608	BGC	O5-C1	6.59	1.54	1.43
2	A	601	BGC	O5-C1	6.68	1.54	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	607	BGC	O5-C1-C2	-4.29	104.04	110.89
2	A	603	BGC	O5-C1-C2	-3.27	105.67	110.89
2	A	608	BGC	O5-C1-C2	-2.72	106.54	110.89
3	A	605	XYS	C5-C4-C3	-2.66	106.33	109.55
3	A	605	XYS	O3-C3-C4	-2.65	105.16	110.01
3	A	605	XYS	C1-C2-C3	-2.64	106.35	109.55
2	A	602	BGC	C1-O5-C5	-2.55	108.39	112.14
3	A	605	XYS	C4-C3-C2	-2.54	108.19	111.02
2	A	607	BGC	C1-O5-C5	-2.48	108.50	112.14
3	A	605	XYS	O2-C2-C3	-2.44	105.27	110.19
2	A	608	BGC	O4-C4-C3	-2.15	105.51	110.36
2	A	601	BGC	O5-C1-C2	-2.05	107.62	110.89
2	A	607	BGC	O5-C5-C6	2.18	112.00	107.34
2	A	608	BGC	C2-C3-C4	2.54	115.48	111.05
2	A	608	BGC	C1-C2-C3	4.01	114.41	109.55
2	A	602	BGC	C1-C2-C3	4.07	114.49	109.55

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	604	XYS	2	0
2	A	606	BGC	2	0
4	A	614	EDO	2	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data ⓘ

### 6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	492/511 (96%)	0.58	37 (7%) 17 24	25, 50, 78, 92	0

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	284	TYR	3.9
1	A	347	GLY	3.6
1	A	133	LYS	3.4
1	A	500	LEU	3.4
1	A	167	LEU	3.3
1	A	47	LEU	3.2
1	A	122	THR	3.1
1	A	123	GLN	3.1
1	A	282	GLU	3.0
1	A	137	ILE	3.0
1	A	300	THR	3.0
1	A	140	ALA	3.0
1	A	480	ILE	2.8
1	A	46	ALA	2.8
1	A	48	GLU	2.8
1	A	50	ALA	2.8
1	A	49	ALA	2.7
1	A	493	ASP	2.7
1	A	201	ASP	2.7
1	A	202	LYS	2.7
1	A	130	GLU	2.6
1	A	199	GLN	2.6
1	A	374	LYS	2.5
1	A	191	TYR	2.5
1	A	372	LYS	2.5
1	A	166	LYS	2.4
1	A	119	ALA	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	341	ILE	2.3
1	A	117	ILE	2.3
1	A	200	SER	2.2
1	A	366	ASP	2.2
1	A	57	ARG	2.2
1	A	290	LEU	2.1
1	A	546	GLU	2.1
1	A	142	LEU	2.0
1	A	494	THR	2.0
1	A	263	TYR	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
4	EDO	A	614	4/4	0.77	0.27	9.71	5,5,30,31	0
4	EDO	A	610	4/4	0.86	0.31	9.13	2,2,54,55	0
4	EDO	A	611	4/4	0.88	0.19	7.89	0,0,35,35	0
4	EDO	A	613	4/4	0.84	0.23	5.89	3,3,39,40	0
3	XYS	A	604	9/10	0.81	0.18	0.22	34,54,63,65	0
2	BGC	A	603	11/12	0.94	0.16	-0.27	16,60,68,78	0
2	BGC	A	607	11/12	0.92	0.12	-3.34	32,64,81,83	0
5	CL	A	615	1/1	0.99	0.07	-4.49	34,34,34,34	0
2	BGC	A	602	11/12	0.91	0.13	-	22,42,67,69	0
6	NA	A	616	1/1	0.95	0.10	-	35,35,35,35	0
2	BGC	A	608	11/12	0.90	0.13	-	42,73,77,79	0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
2	BGC	A	601	11/12	0.88	0.13	-	54,65,68,69	0
2	BGC	A	606	11/12	0.93	0.10	-	23,61,73,76	0
2	BGC	A	609	12/12	0.85	0.16	-	5,5,77,80	0
4	EDO	A	612	4/4	0.81	0.21	-	0,0,41,42	0
3	XYS	A	605	9/10	0.71	0.29	-	5,5,87,88	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.